


表 4.17 Distribution of total AF metabolites

	Control (AF only)	Gypenosides 24h + AF	Gypenosides + AF
Bladder	1.070 ± 0.135	0.764 ± 0.087	0.992 ± 0.074
Blood	3.649 ± 0.499	2.970 ± 0.518	4.140 ± 0.475
Colon	0.914 ± 0.150	0.860 ± 0.210	1.252 ± 0.220
Kidney	5.527 ± 1.362	1.859 ± 0.135*	2.554 ± 0.541*
Liver	61.590 ± 8.600	62.200 ± 4.980	128.090 ± 13.360
Stomach	1.162 ± 0.220	0.048 ± 0.040	0.423 ± 0.158*
Uterus	4.767 ± 0.970	9.064 ± 0.334	2.780 ± 0.556*
Urine	69.026 ± 7.912	38.030 ± 6.490	56.750 ± 7.289
Stool	512.390 ± 35.752	472.540 ± 36.970	278.470 ± 25.550*

Values are mean ± SD; n = 6.

Data were analyzed by an unpaired Student's *t*-test.

\* Difference between AF treated (control) and gypenosides treated 24 hours with AF or AF with gypenosides co-treated.  $p < 0.05$






圖 4.18 Distribution of total AF metabolites




圖 4.19 Distribution of total AAF.