

附圖目錄

- Fig. 1-1** Chromatograms of fisetin (F) and ethyl paraben (internal standard; IS) in rat serum: (a) blank serum ; (b) fisetin and IS spiked in blank serum , F: 21.8 nmol/mL ; (c) serum sample obtained after oral administration of 50 mg/kg fisetin , F: 19.3 nmol/mL.....52
- Fig. 1-2** (a) Mean (\pm S.E.) serum concentration-time profiles of fisetin (), its sulfates () and glucuronides (\blacktriangledown) after intravenous administration of fisetin (10 mg/kg) to six rats and (b) the semi-log diagram of (a).....53
- Fig. 1-3** (a) Mean (\pm S.E.) serum concentration-time profiles of fisetin free form (), its sulfates () and glucuronides (\blacktriangledown) after oral administration of fisetin (50 mg/kg) to six rats and (b) the semi-log diagram of (a).....54
- Fig. 2-1** (a) Mean (\pm S.E.) blood concentration-time profiles of cyclosporine after oral administration of cyclosporine alone (\bullet) and coadministration with 5-hydroxyflavone at doses of 20 mg/kg (\circ) and 40 mg/kg (\blacktriangledown) and (b) the semi-log diagram of (a).....55
- Fig. 2-2** (a) Mean (\pm S.E.) blood concentration-time profiles of cyclosporine after oral administration of cyclosporine alone (\bullet) and coadministration with flavone at doses of 20 mg/kg (\circ) and 40 mg/kg (\blacktriangledown) and (b) the semi-log diagram of (a).....56
- Fig. 2-3** (a) Mean (\pm S.E.) blood concentration-time profiles of cyclosporine after oral administration of cyclosporine alone (\bullet) and coadministration with fisetin at doses of 20 mg/kg (\circ) and 40 mg/kg (\blacktriangledown) and (b) the semi-log diagram of (a).....57

Fig. 2-4	(a) Mean (\pm S.E.) blood concentration-time profiles of cyclosporine after oral administration of cyclosporine alone (●) and coadministration with quercetin at doses of 20 mg/kg (○) and (b) the semi-log diagram of (a).....	58
Fig. 2-5	(a) Mean (\pm S.E.) blood concentration-time profiles of cyclosporine after oral administration of cyclosporine alone (●) and coadministration with morin at doses of 20 mg/kg (○), 37.5 mg/kg (▼) and 50 mg/kg (∇) and (b) the semi-log diagram of (a).....	59
Fig. 3-1	Average transport of rhodamine 123 (ng/mL) across jejunum in the absence (●) or presence of 420 μ M (○) and 840 μ M (▼) 5-hydroxyflavone, respectively.....	60
Fig. 3-2	Average transport of rhodamine 123 (ng/mL) across ileum in the absence (●) or presence of 420 μ M (○) and 840 μ M (▼) 5-hydroxyflavone, respectively.....	60
Fig. 3-3	Average transport of rhodamine 123 (ng/mL) across jejunum in the absence (●) or presence of flavone (○).....	61
Fig. 3-4	Average transport of rhodamine 123 (ng/mL) across ileum in the absence (●) or presence of flavone (○).....	61
Fig. 3-5	Average transport of rhodamine 123 (ng/mL) across jejunum in the absence (●) or presence of morin (○).....	62
Fig. 3-6	Average transport of rhodamine 123 (ng/mL) across ileum in the absence (●) or presence of morin (○).....	62

附表目錄

Table 1-1	Intra-run and inter-run analytical precision and accuracy of fisetin.....	63
Table 1-2	Recovery (%) of fisetin from rat serum (n=3).....	63
Table 1-3	The serum concentrations (nmol/mL) of fisetin in six rats after intravenous administration of fisetin (10 mg/kg).....	64
Table 1-4	The serum concentrations (nmol/mL) of fisetin sulfates in six rats after intravenous administration of fisetin (10 mg/kg).....	64
Table 1-5	The serum concentrations (nmol/mL) of fisetin glucuronides in six rats after intravenous administration of fisetin (10 mg/kg).....	65
Table 1-6	The serum concentrations (nmol/mL) of fisetin in six rats after oral administration of fisetin (50 mg/kg).....	65
Table 1-7	The serum concentrations (nmol/mL) of fisetin sulfates in six rats after oral administration of fisetin (50 mg/kg).....	66
Table 1-8	The serum concentrations (nmol/mL) of fisetin glucuronides in six rats after oral administration of fisetin (50 mg/kg).....	66
Table 1-9	Individual pharmacokinetic parameters of fisetin after intravenous administration of fisetin (10 mg/kg) to six rats.....	67
Table 1-10	Individual pharmacokinetic parameters of fisetin sulfates after intravenous administration of fisetin (10 mg/kg) to six rats.....	67
Table 1-11	Individual pharmacokinetic parameters of fisetin glucuronides after intravenous administration of fisetin (10 mg/kg) to six rats.....	68

Table 1-12	Individual pharmacokinetic parameters of fisetin after oral administration of fisetin (50 mg/kg) to six rats.....	68
Table 1-13	Individual pharmacokinetic parameters of fisetin sulfates after oral administration of fisetin (50 mg/kg) to six rats.....	69
Table 1-14	Individual pharmacokinetic parameters of fisetin glucuronides after oral administration of fisetin (50 mg/kg) to six rats.....	69
Table 1-15	Comparison of pharmacokinetic parameters of fisetin sulfates and glucuronides after intravenous administration of fisetin (10 mg/kg) to six rats.....	70
Table 1-16	Comparison of pharmacokinetic parameters of fisetin sulfates and glucuronides after oral administration of fisetin (50 mg/kg) to six rats.....	70
Table 1-17	Individual pharmacokinetic parameters of fisetin after intravenous administration of fisetin (10 mg/kg) to six rats.....	71
Table 1-18	Individual pharmacokinetic parameters of fisetin sulfates after intravenous administration of fisetin (10 mg/kg) to six rats.....	71
Table 1-19	Individual pharmacokinetic parameters of fisetin glucuronides after intravenous administration of fisetin (10 mg/kg) to six rats.....	72
Table 2-1	Cyclosporine blood concentrations (ng/mL) of six rats after oral administration of cyclosporine (2.5 mg/kg).....	73
Table 2-2	Cyclosporine blood concentrations (ng/mL) of six rats after oral administrations of cyclosporine (2.5 mg/kg) and 5-hydroxyflavone (20 mg/kg).....	73
Table 2-3	Cyclosporine blood concentrations (ng/mL) of seven rats after oral administrations of cyclosporine (2.5 mg/kg) and 5-hydroxyflavone (40 mg/kg).....	74

Table 2-4 Comparison of pharmacokinetic parameters of cyclosporine between rats receiving cyclosporine (2.5 mg/kg) (control) and cyclosporine with 5-hydroxyflavone (20 mg/kg) (treatment)..74

Table 2-5 Comparison of pharmacokinetic parameters of cyclosporine between rats receiving cyclosporine (2.5 mg/kg) (control) and cyclosporine with 5-hydroxyflavone (40 mg/kg) (treatment)..75

Table 2-6 Cyclosporine blood concentrations (ng/mL) of six rats after oral administrations of cyclosporine (2.5 mg/kg) and flavone (20 mg/kg).....75

Table 2-7 Cyclosporine blood concentrations (ng/mL) of six rats after oral administrations of cyclosporine (2.5 mg/kg) and flavone (40 mg/kg).....76

Table 2-8 Comparison of pharmacokinetic parameters of cyclosporine between rats receiving cyclosporine (2.5 mg/kg) (control) and cyclosporine with flavone (20 mg/kg) (treatment).....76

Table 2-9 Comparison of pharmacokinetic parameters of cyclosporine between rats receiving cyclosporine (2.5 mg/kg) (control) and cyclosporine with flavone (40 mg/kg) (treatment).....77

Table 2-10 Cyclosporine blood concentrations (ng/mL) of six rats after oral administrations of cyclosporine (2.5 mg/kg) and fisetin (20 mg/kg).....77

Table 2-11 Cyclosporine blood concentrations (ng/mL) of seven rats after oral administrations of cyclosporine (2.5 mg/kg) and fisetin (40 mg/kg).....78

Table 2-12 Comparison of pharmacokinetic parameters of cyclosporine between rats receiving cyclosporine (2.5 mg/kg) (control) and cyclosporine with fisetin (20 mg/kg) (treatment).....78

Table 2-13 Comparison of pharmacokinetic parameters of cyclosporine between rats receiving cyclosporine (2.5 mg/kg) (control) and cyclosporine with fisetin (40 mg/kg) (treatment).....79

Table 2-14 Cyclosporine blood concentrations (ng/mL) of five rats after oral administrations of cyclosporine (2.5 mg/kg) and quercetin (20 mg/kg).....79

Table 2-15 Comparison of pharmacokinetic parameters of cyclosporine between rats receiving cyclosporine (2.5 mg/kg) (control) and cyclosporine with quercetin (20 mg/kg) (treatment).....80

Table 2-16 Cyclosporine blood concentrations (ng/mL) of six rats after oral administrations of cyclosporine (2.5 mg/kg) and morin (20 mg/kg).....80

Table 2-17 Cyclosporine blood concentrations (ng/mL) of seven rats after oral administrations of cyclosporine (2.5 mg/kg) and morin (37.5 mg/kg).....81

Table 2-18 Cyclosporine blood concentrations (ng/mL) of six rats after oral administrations of cyclosporine (2.5 mg/kg) and morin (50 mg/kg).....81

Table 2-19 Comparison of pharmacokinetic parameters of cyclosporine between rats receiving cyclosporine (2.5 mg/kg) (control) and cyclosporine with morin (20 mg/kg) (treatment).....82

Table 2-20 Comparison of pharmacokinetic parameters of cyclosporine between rats receiving cyclosporine (2.5 mg/kg) (control) and cyclosporine with morin (37.5 mg/kg) (treatment).....82

Table 2-21 Comparison of pharmacokinetic parameters of cyclosporine between rats receiving cyclosporine (2.5 mg/kg) (control) and cyclosporine with morin (50 mg/kg) (treatment).....83

Table 3-1 Transport of rhodamine 123 (ng/mL) from serosal to mucosal side across the everted jejunum of rats (n=3, control).....84

Table 3-2 Transport of rhodamine 123 (ng/mL) from serosal to mucosal side across the everted jejunum of rats (n=3) in the presence of 5-hydroxyflavone at concentration of 420 μ M.....84

Table 3-3 Transport of rhodamine 123 (ng/mL) from serosal to mucosal side across the everted jejunum of rats (n=3) in the presence of 5-hydroxyflavone at concentration of 840 μ M.....84

Table 3-4 Transport of rhodamine 123 (ng/mL) from serosal to mucosal side across the everted jejunum of rats (n=3) in the presence of flavone at concentration of 900 μ M.....85

Table 3-5 Transport of rhodamine 123 (ng/mL) from serosal to mucosal side across the everted jejunum of rats (n=3) in the presence of morin at concentration of 830 μ M.....85

Table 3-6 Transport of rhodamine 123 (ng/mL) from serosal to mucosal side across the everted ileum of rats (n=3, control).....85

Table 3-7 Transport of rhodamine 123 (ng/mL) from serosal to mucosal side across the everted ileum of rats (n=3) in the presence of 5-hydroxyflavone at concentration of 420 μ M.....86

Table 3-8 Transport of rhodamine 123 (ng/mL) from serosal to mucosal side across the everted ileum of rats (n=3) in the presence of 5-hydroxyflavone at concentration of 840 μ M.....86

Table 3-9 Transport of rhodamine 123 (ng/mL) from serosal to mucosal side across the everted ileum of rats (n=3) in the presence of flavone at concentration of 900 μ M.....86

Table 3-10 Transport of rhodamine 123 (ng/mL) from serosal to mucosal side across the everted ileum of rats (n=3) in the presence of morin at concentration of 820 μ M.....87