

表 目 錄

Table 1.	Retention time of flavonoid aglycones and internal standard in various feces by HPLC analysis	117
Table 2.	Concentrations of hesperetin after incubation of hesperidin with rabbit, rat and human feces at 37	118
Table 3.	Concentrations of naringenin after incubation of naringin with rabbit, rat and human feces at 37	118
Table 4.	Concentrations of quercetin after incubation of rutin with rabbit, rat and human feces at 37	119
Table 5.	Concentrations of neophellamuretin after incubation of phellamurin with rabbit, rat and human feces at 37	119
Table 6.	Contents ($\mu\text{g/mL}$) of metabolites after incubation of daidzin and genistin with rabbit and rat feces at 37	120
Table 7.	Remaining percentages (%) of flavonoid aglycones incubated at 37 with artificial intestinal juice	121
Table 8.	Remaining percentages (%) of flavonoid aglycones incubated at 37 with rabbit feces	122
Table 9.	Remaining percentages (%) of flavonoid aglycones incubated at 37 with rat feces	123
Table 10.	Remaining percentages (%) of flavonoid aglycones incubated at 37 with human feces	124
Table 11.	Degradation slope (r) of flavonoid aglycones incubated at 37 with rabbit, rat, human feces and artificial intestinal juice	125
Table 12.	Remanence (peak area ratio) of catalpol with heating or with artificial gastric juice	126
Table 13.	Intraday and interday analytical precision and accuracy of paeoniflorogenin assay in serum (n=3)	127
Table 14.	Recovery (%) of paeoniflorogenin from serum (n=3)	127

Table 15.	Paeoniflorgenin concentrations ($\mu\text{g}/\text{mL}$) in serum after oral administration of Paeoniae radix decoction to rats (n=6)	128
Table 16.	Pharmacokinetic parameters of paeoniflorgenin after oral administration of Paeoniae radix decoction to rats (n=6)	128
Table 17.	Remanence (peak area ratio) of paeoniflorgenin after incubation of paeoniflorin with rabbit, rat, pig and human feces at 37	129
Table 18.	Quercetin contents after hydrolysis of onion and Huaimi with 1.2 N HCl in the presence of various amount of vitamin C at 80 for 1 h or 2 h (n=3)	130
Table 19.	Intraday and interday analytical precision and accuracy of quercetin in human feces (n=3)	131
Table 20.	Intraday and interday analytical precision and accuracy of quercetin in rat feces (n=3)	132
Table 21.	Concentrations ($\mu\text{g}/\text{mL}$) of quercetin after incubation of onion juice and Huaimi infusion with human and rat feces	133
Table 22.	Peak area ratios of quercetin (to internal standard) after incubation of quercetin sulfates with various substances	133
Table 23.	The concentration ranges, regression equations and their correlation coefficients of anthraquinones in rubarb	134
Table 24.	Intraday and interday analytical precision and accuracy of aloe-emodin in rat feces (n=3)	134
Table 25.	Intraday and interday analytical precision and accuracy of rhein in rat feces (n=3)	135
Table 26.	Intraday and interday analytical precision and accuracy of emodin in rat feces (n=3)	136
Table 27.	Intraday and interday analytical precision and accuracy of chrysophanol in rat feces (n=3)	137
Table 28.	Concentrations ($\mu\text{g}/\text{mL}$) of aloe-emodin, rhein, emodin and chrysophanol after incubation of rubarb decoction with rat feces	138

Table 29. Remaining percentages (%) of aloe-emodin, rhein, emodin and chrysophanol after incubation with rat feces at 37 , respectively.....	138
Table 30. Remaining percentages (%) of aloe-emodin, rhein, emodin and chrysophanol after incubation of their mixture at 37 with rat feces	139
Table 31. Intraday and interday analytical precision and accuracy of rhein in rabbit feces (n=3)	140
Table 32. Intraday and interday analytical precision and accuracy of rhein in rat feces (n=3)	141
Table 33. Concentrations (µg/mL) of rhein after incubation of sennosides A (SN-A) and B (SN-B) with rabbit and rat feces	142
Table 34. Remaining percentages (%) of rhein incubated at 37 with rabbit and rat feces	142
Table 35. The concentration ranges, regression equations and their correlation coefficients of flavonoids in <i>Scutellariae Radix</i>	143
Table 36. Intraday and interday analytical precision and accuracy of baicalein in rat feces (n=3)	143
Table 37. Intraday and interday analytical precision and accuracy of wogonin in rat feces (n=3)	144
Table 38. Concentrations (µg/mL) of baicalein and wogonin after incubation of S, SFW or HTS decoctions with rat feces	145
Table 39. Remaining percentages (%) of GA and 3-dehydroGA after incubation of glycyrrhizin or licorice decoction incubated at 37 with rat feces	146
Table 40. Remaining percentages (%) of GA and 3-dehydroGA after incubation of glycyrrhizin or licorice decoction incubated at 37 with pig feces	146

Table 41. Remaining percentages (%) of GA and 3-dehydroGA after incubation of glycyrrhizin or licorice decoction incubated at 37 with human feces	147
Table 42. Remaining percentages (%) of 18 β -glycyrrhetic acid after incubation of licorice decoction alone and with honey, fructose or glucose in rat feces	147
Table 43. Remaining percentages (%) of 3-dehydroglycyrrhetic acid after incubation of licorice decoction alone and with honey, fructose or glucose in rat feces	148