

圖目錄

圖一、研究架構.....	5
圖二、黃酮類配醣體之腸內代謝	8
圖三、芍藥? 的代謝過程	32
圖四、甘草酸的代謝過程	35
圖五、番瀉? 的代謝過程	43
Fig. 1. Mean (\pm S.D.) concentration-time profiles of hesperetin after incubation of hesperidin with human, rabbit, rat feces and artificial gastric juice (n=3).	149
Fig. 2. Mean (\pm S.D.) concentration-time profiles of naringenin after incubation of naringin with human, rabbit, rat feces and artificial gastric juice (n=3).	149
Fig. 3. Mean (\pm S.D.) concentration-time profiles of quercetin after incubation of rutin with human, rabbit, rat feces and artificial gastric juice (n=3).	150
Fig. 4. Mean (\pm S.D.) concentration-time profiles of neophellamuretin after incubation of phellamurin with human, rabbit, rat feces and artificial gastric juice (n=3).	150
Fig. 5. Mean (\pm S.D.) concentration-time profiles of daidzein and genistein after incubation of daidzin and genistin with rabbit and rat feces (n=3).	151
Fig. 6. HPLC chromatogram of flavonoid standards, 1: morin; 2: daidzein; 3: luteolin; 4: quercetin; 5: genistein; 6: naringenin; 7: apigenin; 8: hesperetin; 9: diosmetin; 10: kaempferol; 11: baicalein; 12: 5, 7-dimethoxycoumarin (I.S.); 13: neophellamuretin; 14: wogonin.	151
Fig. 7. Degradation of flavonoid aglycones by artificial intestinal juice (Mean \pm S.D. for each time point, n=3).	152

Fig. 8. Degradation of flavonoid aglycones by rabbit fecal flora (Mean \pm S.D. for each time point, n=3).	152
Fig. 9. Degradation of flavonoid aglycones by rat fecal flora (Mean \pm S.D. for each time point, n=3).	153
Fig. 10. Degradation of flavonoid aglycones by human fecal flora (Mean \pm S.D. for each time point, n=3).	153
Fig. 11. Mean (\pm S.D.) peak area ratio-time profiles of catalpol with heating and with artificial gastric juice (n=3).	154
Fig. 12. HPLC chromatograms of catalpol after boiling with water (A) and incubation with artificial gastric juice at 37 $^{\circ}$ C for 5 min (B). 1: catalpol; 2: 5-hydroxymethyl- 2-furaldehyde (HMF, I.S.).	154
Fig. 13. LC/MS chromatograms of catalpol after incubation with β -glucosidase at 37 $^{\circ}$ C for 2 h.	155
Fig. 14. LC/MS chromatogram (upper) and mass spectrum (lower) of paeoniforgenin.	156
Fig. 15. HPLC chromatograms of traditional decoctions of <i>Paeoniae radix</i> (A) and <i>Paeoniae radix</i> decoction after hydrolysis with 2.4 N HCl (B). 1: paeoniflorin; 2: paeoniforgenin; 3: methylparaben (I.S.).	156
Fig. 16. HPLC chromatograms of paeoniforgenin with internal standard in rat serum: blank serum (A); blank serum spiked with paeoniforgenin and internal standard (B); serum sample obtained 10 min after oral administration of <i>Paeoniae radix</i> decoction (C). 1: paeoniforgenin; 2: methylparaben (I.S.).	157
Fig. 17. Serum concentration-time profiles of paeoniforgenin after oral administration of <i>Paeoniae radix</i> decoction to six rats.	158
Fig. 18. HPLC chromatograms of rabbit (A) and rat (C) blank feces; paeoniforgenin after incubation of paeoniflorin with rabbit (B) and rat (D) feces. 1: paeoniforgenin; 2: methylparaben (IS).	159
Fig. 19. HPLC chromatograms of human (E) and pig (G) blank feces; paeoniforgenin after incubation of paeoniflorin with human (F) and pig (H) feces. 1: paeoniforgenin; 2: methylparaben (IS).	160

Fig. 20. UV spectrums of paeoniforgenin after incubation of paeoniflorin with rat, rabbit, pig and human feces.	161
Fig. 21. Mean (\pm S.D.) peak area ratio-time profiles of paeoniforgenin after incubation of paeoniflorin with rabbit, rat, pig and human feces (n=3).	161
Fig. 22. HPLC chromatograms of 6,7-dimethoxycoumarin (I.S.; 1) and quercetin (2) in methanol solution (A); after hydrolysis of onion juice (B) and Huaimi infusion (C).	162
Fig. 23. HPLC chromatograms of human blank feces (A); quercetin obtained after incubation of onion juice (B) and Huaimi infusion (C) with human feces; rat blank feces (D); quercetin obtained after incubation of onion juice (E) and Huaimi infusion (F) with rat feces. IS: 6,7-dimethoxycoumarin.	163
Fig. 24. Mean (\pm S.D.) concentration-time profiles of quercetin after incubation of onion and Huaimi infusion in rat and human feces (n=3).	164
Fig. 25. HPLC chromatograms of quercetin in standard solution and after incubation of quercetin sulfates with β -glucuronidase and with artificial intestinal juice (A); with human feces and human blank feces (B); with rat feces and rat blank feces (C); with 6,7-dimethoxycoumarin as the internal standard (IS).	165-166
Fig. 26. HPLC chromatograms of blank feces (A); and anthraquinones obtained after incubation of rubarb decoction with rat feces (B) 1: aloe-emodin; 2: rhein; 3: emodin; 4: 2-methylantraquinone (I.S.); 5: chrysophanol.	167
Fig. 27. Mean (\pm S.D.) concentration-time profiles of aloe-emodin, rhein, emodin and chrysophanol after incubation of rubarb decoction with rat feces (n=3).	167
Fig. 28. Mean (\pm S.D.) peak area ratio-time profiles of aloe-emodin, rhein, emodin and chrysophanol after incubation with rat feces (n=3), respectively.	168

Fig. 29. Mean (\pm S.D.) peak area ratio-time profiles of aloe-emodin, rhein, emodin and chrysophanol after incubation of their mixture with rat feces (n=3).	168
Fig. 30. HPLC chromatograms of rabbit blank feces (A); sennoside A incubated with rabbit feces (B); rat blank feces (C) and sennoside B incubated with rat feces (D) 1: rhein; 2: amylyparaben (I.S.).	169
Fig. 31. Mean (\pm S.D.) concentration-time profiles of rhein after incubation of sennosides A and B with rabbit and rat feces (n=3), respectively.	170
Fig. 32. Degradation of rhein by rabbit and rat feces (Mean \pm S.D. for each time point, n=3).	170
Fig. 33. HPLC chromatograms of standard solution (A); blank feces (B); baicalein and wogonin obtained after incubation of S (C), SFW (D) or HTS (E) with rat feces. 1: baicalein; 2: propylparaben (I.S.); 3: wogonin.	171
Fig. 34. Mean (\pm S.D.) concentration-time profiles of baicalein after incubation of S, SFW or HTS with rat feces (n=3). S: Scutellariae Radix, SFW: Scutellariae Radix fried with wine, HTS: honey-treated Scutellariae Radix.	172
Fig. 35. Mean (\pm S.D.) concentration-time profiles of wogonin after incubation of S, SFW or HTS with rat feces (n=3).	172
Fig. 36. HPLC chromatograms of standard solution (A); blank feces (B); GA and 3-dehydroGA obtained after incubation of glycyrrhizin (C) and licorice decoction (D) with pig feces. 1: 2-methylanthraquinone (I.S.); 2: glycyrrhetic acid (GA); 3: 3-dehydroGA.	173
Fig. 37. HPLC chromatograms of blank feces (A); GA and 3-dehydroGA obtained after incubation of glycyrrhizin (B) and licorice decoction (C) with rat feces. 1: 2-methylanthraquinone (I.S.); 2: GA; 3: 3-dehydroGA.	174

Fig. 38. HPLC chromatograms of blank feces (A); GA and 3-dehydroGA obtained after incubation of glycyrrhizin (B) and licorice decoction (C) with human feces. 1: 2-methylanthraquinone (I.S.); 2: GA; 3: 3-dehydroGA.	175
Fig. 39. Mean (\pm S.D.) concentration-time profiles of GA and 3-dehydroGA after incubation of glycyrrhizin or licorice decoction with rabbit feces (n=3).	176
Fig. 40. Mean (\pm S.D.) peak area ratio-time profiles of GA and 3-dehydroGA after incubation of glycyrrhizin or licorice decoction with rat feces (n=3).	176
Fig. 41. Mean (\pm S.D.) peak area ratio-time profiles of GA and 3-dehydroGA after incubation of glycyrrhizin or licorice decoction with pig feces (n=3).	177
Fig. 42. Mean (\pm S.D.) peak area ratio-time profiles of GA and 3-dehydroGA after incubation of glycyrrhizin or licorice decoction with human feces (n=3).	177
Fig. 43. Mean (\pm S.D.) peak area ratio-time profiles of GA and 3-dehydroGA after incubation of licorice decoction (containing 100 μ g/mL GZ) alone or with honey (10 mg/mL) in rat feces (n=3).	178
Fig. 44. Mean (\pm S.D.) peak area ratio-time profiles of GA and 3-dehydroGA after incubation of licorice decoction (containing 100 μ g/mL GZ) alone and with fructose or glucose (10 mg/mL) in rat feces (n=3).	178