

中國醫藥大學藥物化學研究所碩士論文

指導教授：徐素蘭 教授

共同指導教授：李珮端 教授

論文題目

柳橙汁黃酮之代謝動力學
及大黃、黃芩對胺甲葉酸動力學之影響

Metabolic Pharmacokinetics of Flavonoids of Orange
Juice and the Effect of Rhei Rhizoma, Scutellariae
Radix on Methotrexate Pharmacokinetics

研究生 陳仁元

中國醫藥大學藥物化學研究所

中華民國九十四年六月十日

目錄

附圖目錄.....	III
附表目錄.....	VI
中文摘要.....	XI
英文摘要.....	XIII
第一章 緒言.....	1
第二章 總論.....	4
第一節 橙皮苷及其苷元之結構與物化性質.....	4
第二節 橙皮苷及其苷元之藥理活性.....	6
第三節 大黃、黃芩之文獻考察.....	8
第四節 Methotrexate (MTX) 之特性.....	18
第五節 Multidrug resistance protein (MRP) 之特性.....	24
第三章 實驗之部.....	30
第一節 實驗材料.....	30
一、實驗試藥.....	30
二、儀器設備.....	31
三、實驗動物.....	33
四、溶液製備.....	33

第二節 實驗方法.....	35
一、橙皮昔元於大白鼠體內之代謝動力學.....	35
二、橙皮昔於大白鼠體內之代謝動力學.....	40
三、柳橙汁黃酮於人體之尿藥代謝動力學.....	41
四、大黃、黃芩水煎劑於鼠體對MTX動力學之影響.....	48
第四章 結果與討論.....	54
一、橙皮昔元於大白鼠體內之代謝動力學.....	54
二、橙皮昔於大白鼠體內之代謝動力學.....	58
三、柳橙汁黃酮於人體之尿藥代謝動力學.....	59
四、大黃、黃芩水煎劑於鼠體對MTX動力學之影響.....	64
參考文獻.....	116
附錄.....	131

附圖目錄

- Fig. 1-1** Chromatograms of hesperetin and internal standard in rat serum.
(a): blank serum, (b): serum spiked with hesperetin (3.2 µg/mL) and the internal standard, (c): a serum sample treated with glucuronidase, (d): a serum sample treated with sulfatase. H: hesperetin, IS: 5,7-dimethoxycoumarin.....73
- Fig. 1-2** (a): Mean (\pm S.E.) serum concentration-time profiles of hesperetin (●), hesperetin sulfates (○) and hesperetin glucuronides (▼) after intravenous administration of hesperetin (10.0 mg/kg) to six rats (b): the semi-log diagram of (a).....74
- Fig. 1-3** Mean (\pm S.E.) serum concentration-time profiles of hesperetin sulfates (●) and hesperetin glucuronides (○) after oral administration of hesperetin (50.0 mg/kg) to six rats.....75
- Fig. 3-1** Chromatogram of hesperidin, narirutin and internal standard in orange juice. H: hesperidin, N: narirutin, IS: 6,7-dimethoxycoumarin.....76
- Fig. 3-2** Chromatograms of hesperetin, naringenin and the internal standard in human urine. (a): blank urine, (b): urine spiked with hesperetin (10.0 µg/mL), naringenin and internal standard, (c): an urine sample hydrolyzed with glucuronidase, (d): an urine sample hydrolyzed with sulfatase. H: hesperetin, N: naringenin, IS: 5,7-dimethoxycoumarin.....77
- Fig. 3-3** Cumulated urinary excretion of (a): hesperetin sulfates and (b):

hesperetin glucuronides after oral administration of orange juice to eleven volunteers.....	78
Fig. 3-4 Cumulated urinary excretion of (a): naringenin sulfates and (b): naringenin glucuronides after oral administration of orange juice to eleven volunteers.....	79
Fig. 3-5 Individual cumulated urinary recoveries (% of dose) of hesperetin in eleven volunteers after oral administration of orange juice.....	80
Fig. 3-6 Individual cumulated urinary recoveries (% of dose) of naringenin in eleven volunteers after oral administration of orange juice.....	82
Fig. 3-7 Cumulated urinary recoveries (% of dose) of hesperetin sulfates and glucuronides in eleven volunteers after oral administration of orange juice.....	84
Fig. 3-8 Cumulated urinary recoveries (% of dose) of naringenin sulfates and glucuronides in eleven volunteers after oral administration of orange juice.....	85
Fig. 3-9 Mean (\pm S.E.) urinary recovery (% dose /h) of sulfatase and glucuronidase of (a): hesperetin and (b): naringenin in each time interval after administration of orange juice to eleven volunteers.....	86
Fig. 4-1 Mean (\pm S.E.) serum concentration-time profiles of methotrexate after oral methotrexate alone (●) and coadministration with 2.0 g/kg (○), and 1.0 g/kg (▼) of Rhei Rhizoma decoction (n=8).....	87

Fig. 4-2 (a) Mean (\pm S.E.) serum concentration-time profiles of methotrexate after intravenous methotrexate alone (●) and coadministration with Rhei Rhizoma decoction 2.0 g/kg (○) and (b): the semi-log diagram of (a) (n=8).....88

Fig. 4-3 Mean (\pm S.E.) serum concentration-time profiles of methotrexate after oral methotrexate alone (●) and coadministration with 2.0 g/kg (○), 1.0 g/kg (▼) of Scutellariae Radix decoction (n=8).....89

Fig. 4-4 (a) Mean (\pm S.E.) serum concentration-time profiles of methotrexate after intravenous methotrexate alone (●) and coadministration with 2.0 g/kg of Scutellariae Radix decoction (○) and (b): the semi-log diagram of (a) (n=8).....90

附表目錄

Table 1-1 Intraday and interday analytical precision and accuracy of hesperetin in serum.....	91
Table 1-2 Recovery (%) of hesperetin from rat serum.....	91
Table 1-3 Concentrations of hesperetin free form (nmol/mL) in serum after intravenous administration of hesperetin (10.0 mg/kg) to six rats.....	92
Table 1-4 Concentrations of hesperetin sulfates (nmol/mL) in serum after intravenous administration of hesperetin (10.0 mg/kg) to six rats.....	92
Table 1-5 Concentrations of hesperetin glucuronides (nmol/mL) in serum after intravenous administration of hesperetin (10.0 mg/kg) to six rats.....	93
Table 1-6 Pharmacokinetic parameters of hesperetin after intravenous administration of hesperetin to six rats.....	93
Table 1-7 Pharmacokinetic parameters of hesperetin sulfates after intravenous administration of hesperetin to six rats.....	94
Table 1-8 Pharmacokinetic parameters of hesperetin glucuronides after intravenous administration of hesperetin to six rats.....	94
Table 1-9 Concentration of hesperetin sulfates (nmol/mL) in serum after oral administration of hesperetin (50.0 mg/kg) to six rats.....	95
Table 1-10 Concentration of hesperetin glucuronides (nmol/mL) in serum	

after oral administration of hesperetin (50.0 mg/kg) to six rats.....	95
Table 1-11 Pharmacokinetic parameters of hesperetin sulfates after oral administration of hesperetinin to six rats.....	96
Table 1-12 Pharmacokinetic parameters of hesperetin glucuronides after oral administration of hesperetinin to six rats.....	96
Table 3-1 Intraday and interday analytical precision and accuracy of hesperidin in orange juice.....	97
Table 3-2 Intraday and interday analytical precision and accuracy of narirutin in orange juice.....	97
Table 3-3 Recovery (%) of hesperidin from orange juice.....	98
Table 3-4 Recovery (%) of narirutin from orange juice.....	98
Table 3-5 Intraday and interday analytical precision and accuracy of hesperetin in human urine.....	98
Table 3-6 Intraday and interday analytical precision and accuracy of naringenin in human urine.....	99
Table 3-7 Recovery (%) of hesperetin from human urine.....	99
Table 3-8 Recovery (%) of naringenin from human urine.....	99
Table 3-9 Recoveries of hesperetin sulfates ($\mu\text{mol/L}$) in urine after intake of orange juice to eleven subjects.....	100
Table 3-10 Recoveries of hesperetin glucuronides ($\mu\text{mol/L}$) in urine after intake of orange juice to eleven subjects.....	100
Table 3-11 Recoveries of naringenin sulfates ($\mu\text{mol/L}$) in urine after intake of orange juice in eleven subjects.....	101
Table 3-12 Recoveries of naringenin glucuronides ($\mu\text{mol/L}$) in urine after intake of orange juice in eleven subjects.....	101

Table 3-13 Urine recovery of hesperetin sulfates (% of dose/h) in eleven subjects after intake of orange juice during each time interval.....	102
Table 3-14 Urine recovery of hesperetin glucuronides (% of dose/h) in eleven subjects after intake of orange juice during each time interval.....	102
Table 3-15 Urine recovery of naringenin sulfates (% of dose/h) in eleven subjects after intake of orange juice during each time interval.....	103
Table 3-16 Urine recovery of naringenin glucuronides (% of dose/h) in eleven subjects after intake of orange juice during each time interval.....	103
Table 3-17 Half-life of conjugated metabolites of hesperetin and naringenin after oral administration of orange juice to eleven subjects.....	104
Table 4-1 MTX serum concentration ($\mu\text{mol/L}$) of eleven rats after oral administration of MTX (5.0 mg/kg) alone.....	105
Table 4-2 MTX serum concentration ($\mu\text{mol/L}$) of eight rats after oral coadministration of MTX (5.0 mg/kg) with Rhei Rhizoma decoction (2.0 g/kg).....	105
Table 4-3 MTX serum concentration ($\mu\text{mol/L}$) of six rats after oral administration of MTX (5.0 mg/kg) and Rhei Rhizoma decoction (1.0 g/kg).....	106
Table 4-4 Pharmacokinetic parameters of MTX after oral administration of 5.0 mg/kg MTX alone.....	106
Table 4-5 Pharmacokinetic parameters of MTX after oral	

coadministration of 5.0 mg/kg MTX with Rhei Rhizoma decoction (2.0 g/kg).....	107
Table 4-6 Pharmacokinetic parameters of MTX after oral coadministration of 5.0 mg/kg MTX with Rhei Rhizoma decoction (1.0 g/kg).....	107
Table 4-7 Comparison of pharmacokinetic parameters of MTX in rats between oral administration of MTX (5.0 mg/kg) alone and coadministration with decoction of Rhei Rhizoma (1.0 g/kg and 2.0 g/kg).....	108
Table 4-8 MTX serum concentration ($\mu\text{mol/L}$) of seven rats after intravenous administration of MTX alone (1.0 mg/kg)....	109
Table 4-9 MTX serum concentration ($\mu\text{mol/L}$) of six rats after intravenous administration of MTX (1.0 mg/kg) and coadministration with Rhei Rhizoma decoction (2.0 g/kg)	109
Table 4-10 Pharmacokinetic parameters of MTX after intravenous administration of MTX (1.0 mg/kg) to seven rats.....	110
Table 4-11 Pharmacokinetic parameters of MTX between rats after intravenous administration of MTX (1.0 mg/kg) and oral coadministration with Rhei Rhizoma decoction (2.0 g/kg).....	110
Table 4-12 Comparison of pharmacokinetic parameters of MTX in rats between intravenous administration of MTX (1.0 mg/kg) and coadministration with Rhei Rhizoma decoction (2.0 g/kg).....	111
Table 4-13 MTX serum concentration ($\mu\text{mol/L}$) of eleven rats after oral coadministration of MTX (5.0 mg/kg) and Scutellariae	

Radix decoction (2.0 g/kg).....	111
Table 4-14 MTX serum concentration ($\mu\text{mol/L}$) of six rats after oral coadministration of MTX (5.0 mg/kg) and <i>Scutellariae Radix</i> decoction (1.0 g/kg).....	112
Table 4-15 Pharmacokinetic parameters of MTX after oral coadministration of MTX (5.0 mg/kg) with <i>Scutellariae Radix</i> decoction (2.0 mg/kg).....	112
Table 4-16 Pharmacokinetic parameters of MTX after oral coadministration of MTX (5.0 mg/kg) with <i>Scutellariae Radix</i> decoction (1.0 g/kg).....	113
Table 4-17 Comparison of pharmacokinetic parameters of MTX in rats between oral administration of MTX (5.0 mg/kg) alone and coadministration with decoction of <i>Scutellariae Radix</i> (2.0 g/kg and 1.0 g/kg).....	113
Table 4-18 MTX blood concentration ($\mu\text{mol/L}$) of six rats after intravenous administration of MTX (1.0 mg/kg) and oral coadministration with <i>Scutellariae Radix</i> (2.0 g/kg).....	114
Table 4-19 Pharmacokinetic parameters of MTX between rats after intravenous administration of MTX (1.0 mg/kg) and oral coadministration with decoction of <i>Scutellariae Radix</i> (2.0 g/kg).....	114
Table 4-20 Comparison of pharmacokinetic parameters of MTX in rats between receiving MTX (1.0 mg/kg) alone and oral coadministration with decoction of <i>Scutellariae Radix</i> (2.0 g/kg).....	115