



B25 Propofol depresses angiotensin II-induced cell proliferation in rat cardiac fibroblasts

Kar-Lok Wong MD, PhD^{1,2,3}, Yuk-Man Leung PhD^{3,4} Jin-Jer Chen MD⁵,
Tzu-Hung Cheng PhD^{3,5,6}

1. Dept of Anesthesia, China Medical University & Hospital, Taichung
2. Institute of Medical Sciences, China Medical University, Taichung
3. Vascular Biology Research Group, China Medical University, Taichung
4. Graduate Institute of Neural and Cognitive Sciences, CMU, Taichung
5. Cardiovascular Unit, Institute of Biomedical Sciences, Academia Sinica, Taipei
6. Department of Biological Science and Technology, CMU, Taichung

Aim of investigation: Propofol may have beneficial effects on the prevention of angiotensin II (Ang II)-induced cardiac fibroblast proliferation via its anti-oxidative properties. The aim of this study was designed to examine whether propofol may alter Ang II-induced cell proliferation and to identify the putative underlying signaling pathways in rat cardiac fibroblasts.

Methods: The effect of propofol on Ang II-induced NADPH oxidase activity, reactive oxygen species (ROS) formation, ERK1/2 phosphorylation, and activator protein-1 (AP-1)-mediated reporter activity in cultured cardiac fibroblast were examined. In addition, the effect of propofol on nitric oxide (NO) production, protein kinase B (Akt) and eNOS phosphorylations were also tested to elucidate the intracellular mechanism of propofol in proliferation. The p value less than 0.05 were considered significant (ANOVA).

Results: Ang II (100 nM) increased cell proliferation and ET-1 expression which were partially inhibited by propofol (10, 30 μM). Propofol also inhibited Ang II-increased NADPH oxidase activity, ROS formation, ERK phosphorylation, and AP-1-mediated reporter activity. In addition, propofol also increased the nitric oxide generation, Akt and eNOS phosphorylations. L-NAME, an inhibitor of NO synthase, and the short interfering RNA transfection for Akt or eNOS markedly attenuated the inhibitory effect of propofol on Ang II-induced cell proliferation.

Conclusions: we demonstrate for the first time that propofol prevents cardiac fibroblast proliferation by interfering with the generation of ROS and involves the activation of the Akt-eNOS-nitric oxide pathway. Thus, this study delivers important new insight in the molecular pathways that may contribute to the proposed protective effects of propofol in the cardiovascular system (**Acknowledgement: This study was accepted by "Anesthesiology" for publication at July, 2009 Funded by: National Science Council of Taiwan no. NSC-96-2320-B-039-009-MY3 Part of the result presented at the 2009 Annual Meeting of the European Society of Anaesthesiology, Milan, Italy, June 6-9, 2009).