



Available online at www.sciencedirect.com



Journal of Nutritional Biochemistry xx (2011) xxx–xxx

**Journal of  
Nutritional  
Biochemistry**

## A novel mechanism of coenzyme Q10 protects against human endothelial cells from oxidative stress-induced by modulating NO-related pathways☆☆☆

Kun-Ling Tsai<sup>a,c</sup>, Yi-Hsiang Huang<sup>a,c</sup>, Chung-Lan Kao<sup>b</sup>, De-Ming Yang<sup>c</sup>, Hsin-Chen Lee<sup>d</sup>, Hsiang-Yun Chou<sup>e</sup>, Yu-Chih Chen<sup>c</sup>, Guang-Yuh Chiou<sup>c</sup>, Li-Hsin Chen<sup>c,d</sup>, Yi-Ping Yang<sup>a,c</sup>, Tsan-Hung Chiu<sup>f</sup>, Chiou-Sheng Tsai<sup>g</sup>, Hsiu-Chung Ou<sup>h,1,\*</sup>, Shih-Hwa Chiou<sup>a,c,d,\*</sup>

<sup>a</sup>Institute of Clinical Medicine, National Yang-Ming University, Taipei, Taiwan

<sup>b</sup>Department of Physical Medicine and Rehabilitation, Taipei Veterans General Hospital, Taipei, Taiwan

<sup>c</sup>Department of Medical Research and Education, Taipei Veterans General Hospital, Taipei, Taiwan

<sup>d</sup>Department and Institute of Pharmacology, National Yang-Ming University Taipei, Taiwan

<sup>e</sup>Department and Institute of Physiology, National Yang-Ming University, Taipei, Taiwan

<sup>f</sup>Department of Obstetrics and Gynecology, China Medical University Hospital, Taichung, Taiwan

<sup>g</sup>Department of Pathology and Laboratory Medicine, Taichung Veterans General Hospital, Taichung, Taiwan

<sup>h</sup>Department of Physical Therapy and Graduate Institute of Rehabilitation Science, China Medical University, Taichung, Taiwan

Received 27 October 2010; received in revised form 22 January 2011; accepted 29 January 2011

### Abstract

**Background:** Atherosclerosis is a chronic inflammatory disease of the vessel wall associated with oxidized low-density lipoprotein (oxLDL)-induced apoptosis of endothelial cells. Coenzyme Q10 (CoQ10), a potent antioxidant and a critical intermediate of the electron transport chain, has been reported to inhibit LDL oxidation and thus the progression of atherosclerosis. However, its molecular mechanisms on endothelial cells remain still unclarified.

**Methods:** In this study, primary human umbilical vein endothelial cell cultures treated with oxLDL were used to explore the protective effects of CoQ10.

**Results:** Our results showed that CoQ10 attenuated the oxLDL-induced generation of reactive oxygen species and improved the antioxidant capacity. CoQ10 also attenuated the oxLDL-mediated down-regulation of endothelial nitric oxide synthase (eNOS) and up-regulation of inducible nitric oxide synthase (iNOS). In addition, CoQ10 suppressed oxLDL-activated NF- $\kappa$ B and downstream inflammatory mediators, including expression of adhesion molecules, release of proinflammatory cytokines and the adherence of monocytic THP-1 cells. Moreover, CoQ10 attenuated oxLDL-altered proapoptotic responses. The inhibitor of eNOS (L-NIO 10  $\mu$ M) and iNOS (1400W 10  $\mu$ M) as well as NO enhancer (SNP 10  $\mu$ M) were used to clean up the mechanism.

**Conclusion:** These results provide new insight into the possible molecular mechanisms by which CoQ10 protects against atherogenesis by NO-related pathways.

© 2011 Published by Elsevier Inc.

**Keywords:** oxLDL; CoQ10; NO; ROS; Apoptosis; Inflammation

### 1. Introduction

Atherosclerotic cardiovascular disease is associated with oxidative stress [1]. There is an increasing body of evidence showing that oxidized low-density lipoprotein (oxLDL)-induced apoptosis of vascular endothelial cells participates in the pathogenesis of atherosclerosis. The early stages of the atherosclerotic process are initiated by accumulation of oxLDL and activation of endothelial cells with subsequent expression of adhesion molecules and increased binding of monocytes to the vascular endothelium. Proinflammatory cytokines, such as interleukin 8 (IL-8) and tumor necrosis factor  $\alpha$  (TNF- $\alpha$ ), which are released when endothelial cells are exposed to oxLDL, up-regulate the expression of cell adhesion molecules. This series of adverse changes is also associated with a decrease in the bioavailability of nitric oxide (NO); NO is produced in endothelial cells by the constitutively expressed enzyme endothelial nitric oxide synthase (eNOS). Considerable evidence indicates that oxLDL-induced

<sup>\*</sup> Funding: This study was supported by grants from The National Science Council (NSC 98-2320-B-039-020-MY3, NSC 97-3111-B-075-001-MY3 and 97-2320-B-075-003-MY3), China Medical University (CMU99-S-13), Taiwan, ROC, Taipei Veterans General Hospital (V97B1-006, E1-008 and F-001), the Joint Projects of UTVGH (VGHUST 98-p1-01), Yen-Tjing-Ling Medical Foundation (96/97/98), National Yang-Ming University (Ministry of Education, Aim for the Top University Plan) & Genomic Center Project, Institute of Biological medicine (IBMS-CRC99-p01), Academia Sinica and Center of Excellence for Cancer Research at Taipei Veterans General Hospital (DOH99-TD-C-111-007), Taiwan.

<sup>\*\*</sup> Conflict of Interest: None declared.

<sup>\*</sup> Corresponding author. Department of Medical Research and Education, Taipei Veterans General Hospital, Taipei 11217, Taiwan. Tel.: +886 2 28757394; fax: +886 2 28742375.

E-mail address: shchiou@vghtpe.gov.tw (S.-H. Chiou).

<sup>1</sup> Dr. Hsiu-Chung Ou and Dr. Shih-Hwa Chiou contributed equally to

endothelial dysfunction is associated with down-regulation of eNOS and up-regulation of inducible nitric oxide synthase (iNOS). Reactive oxygen species (ROS), especially superoxide, generated by oxLDL directly reacts with NO to form peroxynitrite, a stable molecule that is toxic to endothelial cells. As a superoxide scavenger, NO inhibits the generation of hydrogen peroxide and impedes the activation of NF- $\kappa$ B and the subsequent expression of inflammatory mediators that promote leukocyte adhesion [2] and macrophage recruitment [3].

Moreover, several pathways involved in the cytotoxicity of oxLDL are dependent on the generation of ROS [4]. At high concentrations, ROS can induce cell injury and death by oxidatively modifying proteins, carbohydrates, nucleic acids and lipids. In addition, the proapoptotic effects of oxLDL-induced ROS generation in endothelial cells involve the disturbance of mitochondrial membrane permeability followed by cytochrome *c* release and subsequently the activation of executioner caspases [5]. Therefore, therapeutic interventions involving inhibitors of oxLDL-induced endothelial apoptosis may prevent the development of atherogenic diseases as well as reduce the morbidity and increase the survival rate of patients with cardiovascular diseases.

Coenzyme Q10 (CoQ10) is a key component of mitochondrial oxidative phosphorylation and adenosine triphosphate production [6]. It has also been shown that CoQ10 located in the mitochondria, lysosomes, Golgi and plasma membranes has antioxidant activity either by directly reacting with free radicals or by regenerating tocopherol and ascorbate from their oxidized state [6]. In humans, CoQ10 content is highest at 20 years old in the lung, heart, spleen, liver and kidney and then gradually decreases upon further aging [7]. In addition, patients with cardiovascular disease show decreased CoQ10 levels in the myocardium [8]. CoQ10 has been reported to have numerous biologic effects. For example, it has been shown *in vitro* that CoQ10 promotes cellular membrane repair *via* patch formation [9], affects the expression of genes involved in G-protein-coupled receptor-mediated JAK/STAT signaling pathways and mediates inflammatory-related response by inhibiting NF- $\kappa$ B activation [10]. In addition, it has been shown *in vivo* that CoQ10 reduces exercise-induced muscular injury and leads to a reduction in plasma concentrations of the oxidative stress marker malondialdehyde [11], attenuates hypertension-mediated oxidative damage [12], increases the antioxidant capacity of glutathione reductase and superoxide dismutase (SOD) [13] and attenuates obesity-induced inflammation and oxidative stress damage [14]. Hamilton et al. [15] showed that CoQ10 improved endothelial dysfunction in statin-treated type II diabetic patients. Clinically, CoQ10 has potential for use in prevention and treatment of cardiovascular diseases such as myocardial infarction, congestive heart failure and other drug-induced/disease-induced cardiomyopathies [8,16]. Daily supplementation of CoQ10 decreases plasma LDL concentration as well as increase HDL concentration in humans with hypercholesterolemia [17]. Chapidze et al. [18] reported that treatment with CoQ10 provide roles in positive modification of oxidative stress, anti-atherogenic fraction of lipid profile, atherogenic ratio and platelet aggregability; moreover, CoQ10 revealed anti-aggregatory ability in patients with coronary atherosclerosis [19].

However, the direct effects of CoQ10 against oxidative stress-induced endothelial injury have not been well clarified. In this study, we explored whether CoQ10 attenuates oxLDL-induced damage by modulating the NO-related pathways.

## 2. Materials and methods

### 2.1. Reagents

Fetal bovine serum, M199 and trypsin-EDTA were obtained from Gibco (Grand Island, NY); low serum growth supplement was obtained from Cascade (Portland, OR); CoQ10 (purity  $\geq$ 98% by HPLC), 2',7'-bis-2-carboxyethyl-5 (and -6)-carboxyfluores-

cein-acetoxymethyl ester (BCECF-AM), 1400 W, (1-N<sup>5</sup>-(1-iminoethyl)ornithine hydrochloride; L-NIO), sodium nitroprusside (SNP), 4,6-diamidino-2-phenylindole (DAPI), EDTA, penicillin, and streptomycin were obtained from Sigma (St. Louis, MO); the terminal deoxynucleotidyl transferase-mediated dUTP nick end-labeling (TUNEL) staining kit was obtained from Boehringer Mannheim (Mannheim, Germany); the SOD activity assay kit and the catalase activity assay kit were obtained from Calbiochem (San Diego, CA); 2',7'-dichlorofluorescein acetoxymethyl ester (DCF-AM), Fura-2 AM and the EnzChek caspase 3 assay kit were purchased from Molecular Probes (Eugene, OR); 5,5,8,6,6,8-tetraethylbenzimidazolcarbocyanine iodide (JC-1) and anti-active caspase 3 were obtained from BioVision (Palo Alto, CA); anti-vascular cell adhesion molecule 1 (VCAM-1), anti-intercellular adhesion molecules (ICAM-1) and anti-E-selectin, IL-8 and endothelin 1 (ET-1) ELISA kits and the annexin V apoptosis kit were purchased from R&D Systems (Minneapolis, MN); anti-eNOS, anti-iNOS, anti-nitrotyrosine, anti-cyclooxygenase II (COX-II), anti-P53, anti-phospho-P53, anti-cytochrome *c*, anti-Bcl 2 and anti-Bax were obtained from Transduction Laboratories (San Diego, CA).

### 2.2. Cell cultures

This study was approved by the Research Ethics Committee of the China Medical University Hospital. After receiving written informed consent from the parents, fresh human umbilical cords were obtained from neonates after birth, suspended in Hanks' balanced salt solution (Gibco) and stored at 4°C. Human umbilical vein endothelial cells (HUVECs) were isolated with collagenase and used at passages 2-3 as described previously [20]. THP-1, a human monocytic leukemia cell line, was obtained from ATCC (Rockville, MD) and cultured in RPMI with 10% FBS at a density of 2 to 5  $\times$  10<sup>6</sup> cells/ml, as suggested in the product specification sheet provided by the vendor.

### 2.3. Lipoprotein separation

The protocol for LDL separation used in this study has been described previously [21]. Briefly, native LDL was isolated from fresh normolipidemic human serum by sequential ultracentrifugation ( $\rho$ =1.019-1.210 g/ml) in KBr solution containing 30 mM EDTA. Immediately before oxidation, LDL was separated from EDTA and from diffusible low molecular mass compounds by gel filtration on PD-10 Sephadex G-25 M gel (Pharmacia, St-Quentin, France) in 0.01 M phosphate-buffered saline (PBS; 136.9 mM NaCl, 2.68 mM KCl, 4 mM Na<sub>2</sub>HPO<sub>4</sub>, 1.76 mM KH<sub>2</sub>PO<sub>4</sub>, pH 7.4). Copper-modified LDL (1 mg protein/ml) was prepared by exposing LDL to 10  $\mu$ M CuSO<sub>4</sub> for 16 h at 37°C. The oxLDL we studied had a TBARS value of 16-20 nM/mg protein of LDL protein; native LDL had no detectable TBARS.

### 2.4. Determination of cytotoxicity and indices of apoptosis

To determine the effect of CoQ10 on oxLDL-induced cytotoxicity, HUVECs were incubated with indicated concentrations of CoQ10 (chloroform-dissolved) for 2 h and then exposed to 130  $\mu$ g/ml oxLDL for an additional 24 h. Cell viability was assessed by the MTT assay [22]. CoQ10 was present during oxLDL exposure. Apoptotic cells were determined by annexin V and TUNEL assays. In preparation for flow cytometry, cells were harvested and stained with both annexin V and PI for 10 min. They were then washed in PBS, dissolved in HEPES buffer. Apoptotic cells assessed by the TUNEL assay were visualized under a fluorescence microscope or analyzed by flow cytometry [23].

### 2.5. Measurement of ROS production

HUVECs (10<sup>4</sup> cells/well) in 96-well plates were preincubated with various concentrations of CoQ10 for 2 h and then incubated with 10  $\mu$ M DCF-AM for 1 h; the fluorescence intensity was measured with a fluorescence microplate reader (Labsystems, CA) calibrated for excitation at 485 nm and emission at 538 nm (before and after 2 h of stimulation with 130  $\mu$ g/ml oxLDL). CoQ10 was present during oxLDL exposure. The percentage increase in fluorescence per well was calculated by the formula [(F<sub>t2</sub> - F<sub>t0</sub>)/F<sub>t0</sub>] $\times$ 100, where F<sub>t2</sub> is the fluorescence at 2 h of oxLDL exposure and F<sub>t0</sub> is the fluorescence at 0 min of oxLDL exposure.

### 2.6. Measurement of antioxidant enzyme activity

To determine the effects of CoQ10 after oxLDL exposure, SOD and catalase activity in the homogenate was determined by an enzymatic assay method using a commercial kit according to the manufacturer's instructions. Enzyme activity was converted to units per milligram of protein.

### 2.7. Immunoblotting

To determine whether CoQ10 could attenuate the oxLDL-induced expression of apoptosis-regulating proteins, HUVECs were pretreated with CoQ10 for 2 h and then stimulated with oxLDL for 24 h; CoQ10 was present during oxLDL exposure. At the end of stimulation, cells were lysed in RIPA buffer. Proteins were then separated by electrophoresis on SDS-polyacrylamide gel. After the proteins had been transferred onto a polyvinylidene difluoride membrane (Millipore, Bedford, MA), the blot was incubated with blocking buffer (1  $\times$  PBS and 5% nonfat dry milk) for 1 h at room

- 178 temperature and then probed with primary antibodies (eNOS, iNOS, nitrotyrosine, 244  
179 COX-II, phosphorylated P53, P53, Bcl-2, Bax; 1:1000 dilutions) overnight at 4°C, 245  
180 followed by incubation with horseradish peroxidase-conjugated secondary antibody  
181 (1:5000) for 1 h. The intensities were quantified by densitometric analysis (Digital  
182 Protein DNA ImagineWare, Huntington Station, NY).
- 183 **2.8. Nitrite (NO<sub>2</sub><sup>-</sup>) accumulation** 246
- 184 NO<sub>2</sub><sup>-</sup> accumulation was used as an indicator of NO production in the medium and 247  
185 was assayed by Gries reagent [24]. Briefly, 100 μl of Gries reagent (1% sulfanilamide– 248  
186 0.1% naphthylethylene diamine dihydrochloride–2.5% H<sub>3</sub>PO<sub>4</sub>) (Sigma) was added to 249  
187 100 μl of each supernatant in triplicate wells of 96-well plates. The plates were read in a 250  
188 microplate reader (Molecular Devices, Palo Alto, CA) at 550 nm against a standard 251  
189 curve of NaNO<sub>2</sub> in culture medium. 252
- 190 **2.9. NF-κB assay** 253
- 191 To explore the effects of CoQ10 on oxLDL-induced NF-κB activation, HUVECs were 254  
192 pretreated with CoQ10 for 2 h and then stimulated with oxLDL (130 μg/ml) for 1 h. In 255  
193 some cases, HUVECs were incubated with specific eNOS inhibitor (L-NIO, 10 μM),  
194 specific iNOS inhibitor (1400W, 10 μM) or exogenous NO donor SNP (25 μM) for 1 h  
195 before the treatment with CoQ10 and oxLDL. CoQ10 was present during oxLDL  
196 exposure. To prepare nuclear extracts for the NF-κB assay, the cells were first  
197 resuspended in buffer containing 10 mM HEPES (pH 7.9), 1.5 mM MgCl<sub>2</sub>, 10 mM KCl,  
198 0.5 mM dithiothreitol and 0.2 mM phenylmethylsulfonyl fluoride (PMSF), followed by  
199 vigorous vortexing for 15 s. The samples were allowed to stand at 4°C for 10 min and  
200 then were centrifuged at 2000 rpm for 2 min. The pelleted nuclei were resuspended in  
201 30 μl buffer containing 20 mM HEPES (pH 7.9), 25% glycerol, 420 mM NaCl, 1.5 mM  
202 MgCl<sub>2</sub>, 0.2 mM EDTA, 0.5 mM dithiothreitol and 0.2 mM PMSF and incubated for 20 min  
203 on ice. The nuclear lysates were then centrifuged at 15,000 rpm for 2 min. Supernatants  
204 containing the solubilized nuclear proteins were stored at –70°C for subsequent NF-κB  
205 assay. NF-κB activity was measured by an NF-κB p65 Active ELISA kit (Imgenex, San  
206 Diego, CA) according to the manufacturer's instructions. The absorbance at 405 nm was  
207 determined using a microplate reader (spectraMAX 340).
- 208 **2.10. Assay for ET-1 and IL-8 secretion**
- 209 HUVECs were pretreated with the indicated concentrations of CoQ10 for 2 h  
210 followed by treatment with oxLDL (130 μg/ml) for 24 h; CoQ10 was present during  
211 oxLDL exposure. At the end of the oxLDL incubation period, cell supernatants were  
212 removed and assayed for ET-1 and IL-8 concentrations using an ELISA kit obtained from  
213 R&D Systems. Data are expressed in nanograms per milliliter of duplicate samples.
- 214 **2.11. Adhesion assay**
- 215 HUVECs at 1×10<sup>5</sup> cells/ml were cultured in 96-well plates. Cells were pretreated  
216 with the indicated concentrations of CoQ10 for 2 h followed by oxLDL (130 μg/ml) for  
217 24 h; CoQ10 was present during oxLDL exposure. The medium was then removed and  
218 0.1 ml/well of THP-1 cells (prelabeled with BECF-AM 4 μM for 30 min in RPMI at 1×10<sup>6</sup>  
219 cell/ml density) were added to fresh RPMI. The cells were allowed to adhere at 37°C for  
220 1 h in a 5% CO<sub>2</sub> incubator. Plates were washed three times with to remove the  
221 nonadherent cells by M199. The number of adherent cells was estimated by  
222 microscopic examination and then the cells were lysed with 0.1 ml 0.25% Triton  
223 X-100. The fluorescence intensity was measured with a fluorescence microplate reader  
224 (Lab System) calibrated for an excitation at 485 nm and for emission at 538 nm.
- 225 **2.12. Adhesion molecule expression**
- 226 To determine whether CoQ10 could attenuate the adhesion molecule expression  
227 induced by oxLDL, HUVECs were pretreated with CoQ10 for 2 h and stimulated with  
228 oxLDL (130 μg/ml) for 24 h; CoQ10 was present during oxLDL exposure. At the end of  
229 stimulation, HUVECs were harvested and incubated with fluorescein isothiocyanate  
230 (FITC)-conjugated antibody (R&D Systems) for 45 min at room temperature. Their  
231 immunofluorescence intensity was analyzed by flow cytometry using a Becton  
232 Dickinson FACScan flow cytometer (Mountain View, CA).
- 233 **2.13. Measurement of [Ca<sup>2+</sup>]<sub>i</sub>**
- 234 To determine the effect of CoQ10 on the oxLDL-induced increase in intracellular  
235 calcium concentration, HUVECs were seeded onto 24-mm glass coverslips, pretreated  
236 with CoQ10 for 2 h and then stimulated with oxLDL (130 μg/ml) for 24 h. The cells on  
237 the coverslips were loaded with 2 μM fura-2 AM (Molecular Probe) in M199 and  
238 allowed to stand for 30 min at 37°C. After loading, the cells were washed with PBS to  
239 remove excess fluorescent dye. Then, the fluorescence of the cells on each coverslip  
240 was measured and recorded using an inverted Olympus microscope IX-70. [Ca<sup>2+</sup>]<sub>i</sub> in  
241 endothelial cells was monitored at an emission wavelength of 510 nm with excitation  
242 wavelengths alternating between 340 and 380 nm with the use of a cooled charge-  
243 coupled device (CCD) camera (MicroMAX, 782YHS; Roper Scientific, Trenton, NJ),
- recorded using SimplePCI 6.0 (Compix Institute, Cranberry Township, PA) and 244  
calculated using Grynkiewicz's method [25]. 245
- 2.14. Measurement of mitochondrial membrane potential** 246
- The lipophilic cationic probe fluorochrome 5,5',6,6'-tetraethylbenzimidazolcar- 247  
boyanine iodide (JC-1) was used to explore the effects of CoQ10 on mitochondria 248  
membrane potential (ΔΨ<sub>m</sub>). JC-1 exists either as a green fluorescent monomer at 249  
depolarized membrane potentials or as a red fluorescent J-aggregate at hyperpolarized 250  
membrane potentials. After treating cells with oxLDL (130 μg/ml) for 24 h in the 251  
presence or absence of various concentrations of CoQ10, cells were rinsed with M199 252  
and JC-1 (5 μM) was loaded. After 20 min of incubation at 37°C, cells were examined 253  
under a fluorescence microscope. Determination of the ΔΨ<sub>m</sub> was carried out using a 254  
FACScan flow cytometer [26]. 255
- 2.15. Isolation of cytosolic fraction for cytochrome c analysis** 256
- After treating cells with oxLDL in the presence or absence of CoQ10, the cells were 257  
collected and lysed with lysis buffer for 20 min on ice. The samples were homogenized 258  
by 10 passages through two 2-gauge needle. The homogenate was centrifuged at 259  
12,000 rpm for 20 min at 4°C. A volume of cell lysates containing 30 μg of protein was 260  
analyzed by Western blot analysis for cytochrome c (1:1000) and β-actin (1:50,000). 261
- 2.16. Measurement of active caspase 3** 262
- To explore the effects of CoQ10 on oxLDL-induced caspase 3 activation, HUVECs 263  
were pretreated with CoQ10 for 2 h and then stimulated with oxLDL (130 μg/ml) for 24 264  
h. In some cases, HUVECs were incubated with specific eNOS inhibitor (L-NIO, 10 μM), 265  
specific iNOS inhibitor (1400W, 10 μM) or exogenous NO donor SNP (25 μM) for 1 h 266  
before the treatment with CoQ10 and oxLDL. The level of active caspase-3 was detected 267  
by flow cytometry using a commercial fluorescein active caspase kit (Mountain View, 268  
CA) under a fluorescence microscope. The activity of caspase 3 was also measured by an 269  
EnzChek caspase-3 assay kit according to the manufacturer's instructions (Molecular 270  
Probes). After being lysed by repeated freeze–thaw cycles. Equal amounts of protein 271  
(50 μg) were added to the reaction buffer containing 5 mM of caspase 3 substrate 272  
Z-DEVD-R110, and the mixture was incubated at room temperature for 30 min. The 273  
fluorescence generated from cleavage of the substrate by caspase 3 was monitored 274  
with a fluorescence microplate reader (Labsystems) calibrated for excitation at 496 nm 275  
and for emission at 520 nm. 276
- 2.17. Statistical analyses** 277
- Results are expressed as mean±S.E. Differences between groups were analyzed 278  
using one-way ANOVA followed by Bonferroni's post hoc test. A P value <0.05 was 279  
considered statistically significant. 280
- 3. Results** 281
- 3.1. CoQ10 blocked phosphatidylserine translocation and DNA damage 282**  
**and cell death induced by oxLDL in HUVECs** 283
- After exposure to oxLDL, the number of shrunken cells or cells 284  
with blebbing membranes was significantly lower in HUVECs that had 285  
been pretreated with CoQ10 than in those that had not been exposed 286  
to CoQ10 (Fig. 1A). As seen in Fig. 1B, the viability of HUVECs was 287  
reduced by almost 55% after exposure to oxLDL; however, CoQ10 288  
significantly increased the viability in a dose-dependent manner. 289  
Normal LDL (nLDL) was used as one negative control, as our finding, 290  
the viability of HUVECs was not repressed after exposure to nLDL. 291  
In addition, CoQ10 treatment was not cytotoxic up to a 292  
concentration of 50 μM. The ED50 (half-maximal effective concen- 293  
tration) of CoQ10 was 4.2 μM after exposure to 130 μg/ml oxLDL for 24 294  
h. Therefore, 130 μg/ml of oxLDL and 2.5–20 μM of CoQ10 were used 295  
in the following experiments. 296
- The annexin V and TUNEL assays were used to ascertain the 297  
apoptotic effects of oxLDL and the antiapoptotic effects of CoQ10 in 298  
HUVECs. Flow cytometric analysis revealed that the percentage of 299  
apoptotic cells in untreated HUVECs was less than 3%. In HUVECs 300  
exposed to oxLDL, however, the percentage of apoptotic cells (20.4%) 301  
and necrotic cells (10.9%) was markedly higher. The percentage of 302  
apoptotic cells in HUVECs pretreated with 20 μM CoQ10 (4.9%) was 303  
close to that in untreated HUVECs (2.9%) (Fig. 1C). 304




Fig. 1. Effects of CoQ10 on oxLDL-induced endothelial cell death. HUVECs were incubated with oxLDL (130 µg/ml) or nLDL in the absence or presence of indicated concentrations of CoQ10 for 24 h. Photomicrographs from phase-contrast microscopy (A). Viability was determined via MTT assay. Effects of CoQ10 on oxLDL-induced endothelial apoptosis (B). HUVECs were incubated with oxLDL (130 µg/ml) in the absence or presence of indicated concentrations of CoQ10 for 24 h. Apoptotic and necrotic death of oxLDL-exposed HUVECs were measured using a FITC-labeled annexin V assay and PI staining. Flow cytometry was used for confirmation (C). HUVECs were incubated with oxLDL in the absence (middle) or presence (right) of 10 µM CoQ10 for 24 h. The lower right quadrants represent the apoptotic cells and the upper right quadrants represent the necrotic cells. Late apoptotic death of oxLDL-exposed HUVECs was evaluated using the TUNEL assay (D). Data are expressed as the mean±S.E. of three independent analyses (E). <sup>#</sup>*P*<.05 vs. untreated control; <sup>\*</sup>*P*<.05 vs. oxLDL treatment.

380 The TUNEL and DAPI staining assays were then used to clarify the  
 381 protective effects of CoQ10 against oxLDL-induced DNA damage. As  
 382 shown in Fig. 1D,E, cells incubated with oxLDL for 24 h showed typical  
 383 features of apoptosis, including the formation of condensed nuclei.  
 384 Those morphologic features were not observed in HUVECs pretreated  
 with CoQ10.

### 385 3.2. CoQ10 inhibited oxLDL-induced intracellular ROS generation 386 in HUVECs

387 To clarify whether the observed antiapoptotic effect of CoQ10 can  
 388 be attributed to reduction in oxidative stress. We found that  
 389 treatment with oxLDL for 2 h produced an eightfold increase in ROS  
 390 generation. Pretreatment of HUVECs with CoQ10 (2.5–20  $\mu$ M) led to a  
 391 dose-dependent reduction in ROS (Fig. 2A, B, all  $P < .05$ ). In this  
 392 investigation, nLDL was used as one negative control, as our result,  
 393 there are no significance difference between control group and nLDL  
 394 group in intracellular ROS level.

395 To examine the mechanisms involved in the antioxidant action of  
 396 CoQ10 in HUVECs exposed to oxLDL, we analyzed the activities of  
 397 antioxidant enzyme activity in HUVECs treated with 130  $\mu$ g/ml of  
 398 oxLDL for 2 h. As shown in Fig. 2C,D, the activity of SOD and catalase  
 399 were reduced by 48% and 68%, respectively, in cells exposed to oxLDL;  
 400 in contrast, pretreatment of cells with CoQ10 (2.5–20  $\mu$ M) signifi-  
 401 cantly potentiated the activity of those antioxidant enzymes in a  
 402 dose-dependent manner (all  $P < .05$ ).

### 387 3.3. CoQ10 protects against oxLDL-induced release of ET-1 and 388 suppression of NOS protein expression

405 To validate whether NO and NO synthase were involved in CoQ10  
 406 suppress oxLDL-induced injury. We, therefore, studied the effects of  
 407 CoQ10 on protein expression of eNOS and iNOS as well as NO content  
 408 and formation of nitrotyrosine. As shown in Fig. 3A,B, exposure to  
 409 oxLDL resulted in significantly lower eNOS expression, higher iNOS  
 410 expression and higher levels of nitrotyrosin than control cells;  
 411 however, in HUVECs pretreated with 10  $\mu$ M CoQ10, the levels of  
 412 eNOS, iNOS and nitrotyrosin expression were close to those seen in  
 413 control cells. Furthermore, the oxLDL-enhanced release of NO  
 414 (Fig. 3C) was also suppressed in HUVECs pretreated with CoQ10  
 415 and the selective iNOS inhibitor 1400W but not eNOS inhibitor L-NIO.  
 416 It has been speculated that reduced NO release/bioavailability  
 417 and enhanced release of ET-1 may contribute to oxLDL-induced  
 418 endothelial dysfunction and the development of atherosclerosis [27].  
 419 Our results display that CoQ10 inhibited oxLDL-enhanced ET-1  
 420 secretion (Fig. 3D).

### 421 3.4. CoQ10-mediated protective function involving eNOS action and 422 CoQ10 mitigated the NF- $\kappa$ B and NF- $\kappa$ B-related proinflammatory 423 response in HUVECs

424 NF- $\kappa$ B is a vital player in the regulation of inflammatory response,  
 425 apoptosis and cell survival [28]. In addition, NO inhibits cleavage of




Fig. 2. The protective effects of CoQ10 on oxLDL-mediated ROS generation in HUVECs. After preincubation for 2 h with the indicated concentrations of CoQ10 (2.5–20  $\mu$ M), HUVECs were incubated with the  $H_2O_2$ -sensitive fluorescent probe DCF-AM (10  $\mu$ M) for 1 h, followed by treatment with 130  $\mu$ g/ml oxLDL or nLDL for 2 h. (A) Fluorescence images exhibited the ROS level in control cells (left) and HUVECs stimulated with oxLDL (middle) in the presence of 10  $\mu$ M CoQ10 (right). (B) Fluorescence intensity of HUVECs was measured with a fluorescence microplate reader. Fluorescence distribution of DCF-AM oxidation was expressed as a percentage of increased intensity. The activity of (C) SOD and (D) catalase in HUVECs stimulated with oxLDL in the absence or presence of indicated concentrations of CoQ10 were determined. Data are expressed as the mean  $\pm$  S.E. of three independent analyses. <sup>#</sup> $P < .05$  vs. untreated control; <sup>\*</sup> $P < .05$  compared with oxLDL treatment.




Fig. 3. CoQ10 suppressed the oxLDL-induced down-regulation of eNOS, up-regulation of iNOS protein expression (A, B) and oxLDL-enhanced NO release (C). In addition, CoQ10 attenuated the oxLDL-enhanced secretion of ET-1 (D). HUVECs were pretreated for 2 h with the indicated concentrations of CoQ10 followed by stimulation with oxLDL (130 µg/ml) for 24 h. At the end of the incubation period, level of eNOS, iNOS and nitrotyrosine protein were determined by immunoblotting; content of NO was assayed using Griess reagent; ET-1 secretion was measured by ELISA. The values represent means±S.E. from three separate experiments. <sup>#</sup>*P*<.05 vs. untreated control; <sup>\*</sup>*P*<.05 vs. oxLDL treatment.

426 IκB and NF-κB activation [29]. We hypothesized that oxLDL induces  
 427 NF-κB activation by reducing the bioavailability of NO and that oxLDL-  
 428 induced NF-κB activation could be reversed by CoQ10. As shown in  
 429 Fig. 4A, pretreatment of HUVECs with CoQ10 conspicuously inhibited  
 430 the oxLDL-induced activation of NF-κB. Moreover, cells pretreatment  
 431 with L-NIO partially antagonized the inhibitory effect of CoQ10.  
 432 Additionally, cells pretreated with 1400W or exogenous donor of  
 433 NO (SNP) showed a marked reduction in the activation of NF-κB  
 434 (all *P*<.05).

435 Proinflammatory cytokines, COX-II and adhesion molecules were  
 436 mediated by NF-κB. All of the proinflammatory events subsequently  
 437 lead to the tethering and adherence of monocytic cells to endothelial  
 438 cells. Our results showed that pretreatment with CoQ10 attenuated  
 439 the release of IL-8 (Fig. 4B), the expression of COX-II (Fig. 4C,D),  
 440 the adhesion of monocytic THP-1 cells to HUVECs exposed to oxLDL (Fig.  
 441 4E, F) and the expression of adhesion molecules (ICAM-1, VCAM-1  
 442 and E-selectin) (Fig. 4G).

### 443 3.5. CoQ10 suppressed the oxLDL-induced apoptotic responses

444 Intracellular calcium rise is involved in oxLDL-induced endothelial  
 445 apoptosis [30]. To validate the protective effect of chronic exposure of  
 446 HUVECs to a detrimental concentration of oxLDL on intracellular  
 447 calcium concentration, we incubated HUVECs with oxLDL (130 µg/ml)  
 448 in the absence or presence of different concentrations of CoQ10. Our

449 finding that the 340/380 ratio of  $[Ca^{2+}]_i$  elevated in oxLDL-exposed  
 450 cells after 24 h was reduced in cells pretreatment of CoQ10 (Fig. 5A,  
 451 all *P*<.05).

452 The increase in intracellular  $Ca^{2+}$  triggers the activation of several  
 453 calcium-dependent proapoptotic signaling transduction pathways.  
 454 In addition, calcium is the most important signal for opening of the  
 455 mitochondrial permeability transition pore (PTP), a mechanism that  
 456 triggers apoptosis. As a consequence of both the dysfunction of the  
 457 electrochemical gradient caused by pore opening and rupture of the  
 458 outer mitochondrial membrane, the mitochondrial membrane po-  
 459 tential ( $\Psi_m$ ) generally collapses. We, therefore, determined the  
 460 mitochondrial permeability to understand whether CoQ10 preserves  
 461 mitochondrial stability after exposure to oxLDL. As shown in Fig. 5B,  
 462 oxLDL depolarized the mitochondrial transmembrane potential in  
 463 HUVECs, as shown by the increase in green fluorescence (middle  
 464 panel); however, pretreatment with CoQ10 contributed to the  
 465 maintenance of mitochondrial transmembrane potential, as indicated  
 466 by repression of green fluorescence and restoration of red fluores-  
 467 cence (right panel). The results of flow cytometry supported those  
 468 findings (Fig. 5C).

469 Apoptosis regulator proteins in the Bcl-2 family govern mitochon-  
 470 drial outer membrane permeabilization and can be either proapopto-  
 471 tic or antiapoptotic. Cheng et al. [31] reported that oxLDL-induced  
 472 generation of ROS in endothelial cells leads to the activation of P53,  
 473 which subsequently induces a conformational change in Bax that 473




Fig. 4. Effects of CoQ10 on oxLDL-induced NF- $\kappa$ B activation. HUVECs were pretreated with each inhibitor 1 h before incubated for additional 1 h with oxLDL (130  $\mu$ g/ml). Nucleic proteins were extracted for nuclear translocation assay of NF- $\kappa$ Bp65 (A). Effects of CoQ10 on oxLDL-induced IL-8 release, COX-II expression, adhesion molecule expression and adhesiveness of THP-1 monocytic cells to HUVECs. HUVECs were pretreated with indicated concentrations of CoQ10 for 2 h followed by stimulation with oxLDL (130  $\mu$ g/ml) for another 24 h. (B) Media were collected for ELISA assay of IL-8 protein levels. (C, D) Protein levels of COX-II were assessed by Western blotting. (E, F) THP-1 cells preloaded with BECEF were incubated with HUVECs for 1 h. The adhesiveness of HUVECs to THP-1 was measured as described in the Materials and Methods. (G) Cell surface expression of ICAM-1, VCAM-1 and E-selectin was determined by flow cytometry. The values represent means  $\pm$  S.E. from three separate experiments. <sup>#</sup> $P$  < .05 vs. untreated control; <sup>\*</sup> $P$  < .05 vs. oxLDL treatment. <sup>&</sup> $P$  < .05 vs. oxLDL+CoQ10 treatment.




Fig. 5. Effects of CoQ10 on oxLDL-triggered intracellular calcium rise (A). HUVECs were pretreated with indicated concentrations of CoQ10 for 2 h followed by stimulation with oxLDL (130 μg/ml) for another 24 h. The measurement of intracellular calcium is described in the Materials and Methods. The influence of CoQ10 on mitochondrial transmembrane permeability transition induced by oxLDL. (B)  $\Delta\Psi_m$  was inspected with the signal from monomeric and J-aggregate JC-1 fluorescence, as described in the Materials and Methods. (left) No treatment; (middle) oxLDL; (right) oxLDL + CoQ10. (C) JC-1 fluorescence was confirmed by flow cytometry. (D–F) Immunoblotting analysis of apoptotic cells in HUVECs exposed to oxLDL and CoQ10. HUVECs were incubated with 130 μg/ml oxLDL in the absence or presence of indicated concentrations (2.5–20 μM) of CoQ10 for 24 h. Representative Western blots and summary data showing that oxLDL up-regulated proapoptotic (phosphorylated P53, Bax) and down-regulated antiapoptotic (Bcl-2) proteins and led to an increase in the concentration of cytosolic cytochrome c. Intervention with CoQ10 suppressed the above-mentioned oxLDL-induced responses. Results were confirmed by densitometric analysis; the values are presented as means  $\pm$  S.E. of three separate experiments. # $P$  < .05 vs. untreated control; \* $P$  < .05 vs. oxLDL treatment.






Fig. 6. Effects of CoQ10 on oxLDL-induced caspase 3 activation. (A) HUVECs were incubated for 1 h with indicated concentrations of CoQ10, followed by exposure to oxLDL (130 μg/ml) for another 24 h. (B) Fluorescent images show the activated caspase 3 level in control cells (left), HUVECs stimulated with oxLDL (middle), and in the presence of 10 μM CoQ10 (right). (C) Fluorescence intensity of cells was measured by flow cytometry. The activity of caspase 3 was measured by EnzCaspase-3 assay kit. Data are expressed as the mean±S.E. of three independent analyses. <sup>#</sup>*P*<.05 vs. untreated control; <sup>\*</sup>*P*<.05 vs. oxLDL treatment. <sup>&</sup>*P*<.05 vs. oxLDL+CoQ10 treatment.

474 enables the mitochondrial translocation of that proapoptotic protein.  
475 Our results showed that CoQ10 significantly reduced the activation of  
476 P53 and the expression of Bax and significantly increased the  
477 expression of the antiapoptotic protein Bcl-2 in a concentration-  
478 dependent manner (Fig. 5C-E).

479 Disintegration of mitochondrial membrane function contributes to  
480 the release of cytochrome *c* from the mitochondria. As shown in  
481 Fig. 5C,F, CoQ10 reduced the cytosolic cytochrome *c* level in oxLDL-  
482 exposed HUVECs.

### 483 3.6. CoQ10 prevented oxLDL-induced caspase 3 activation

484 To examine whether CoQ10 ultimately influences the activity of  
485 this apoptotic effector in modulating apoptosis, we studied the effects  
486 of CoQ10 on oxLDL-induced activation of caspase 3 using fluorescence  
487 microscopy and flow cytometry. As shown in Fig. 6A,B, CoQ10  
488 inhibited oxLDL-induced caspase 3 activation. We also determined  
489 the activity of caspase 3 using the EnzCaspase-3 assay kit. As seen in  
490 Fig. 6C, CoQ10 effectively inhibited the cleavage of caspase 3 activated  
491 by oxLDL. Simultaneous treatment of HUVECs with CoQ10 and eNOS  
492 inhibitor partially abolished the inhibitory effects of CoQ10 on  
493 caspase 3 activity. Furthermore, the addition of 1400W or SNP  
494 definitely inhibited oxLDL-induced activation of caspase 3.

## 495 4. Discussion

496 In the present study, we demonstrated that CoQ10 ameliorated  
497 oxLDL-induced endothelial dysfunction by inhibiting inflammatory  
498 and oxidative damage that leads to cellular apoptosis. Specifically,  
499 CoQ10 suppressed the generation of ROS, which subsequently  
500 attenuated the oxLDL-impaired expression of antioxidant enzymes,  
501 increased the bioavailability of NO, reduced ET-1 secretion, stabilized

the mitochondrial membrane and maintained the endothelial [Ca<sup>2+</sup>]<sub>i</sub> 502  
level, thereby preventing the release of cytochrome *c*, a molecule 503  
required for the activation of the proapoptotic protein caspase 3. 504

The generation of ROS and the activities of antioxidant enzymes 505  
must be kept in balance to preserve homeostasis and a stable 506  
intracellular redox state for normal cell function. Consistent with 507  
findings from previous studies that CoQ10 supplementation reversed 508  
the oxidative stress-induced suppression of antioxidant enzyme 509  
expression [13,32], our findings show that pretreatment with 510  
CoQ10 suppressed the oxLDL-induced reduction in SOD and catalase 511  
activities and resulted in decreased ROS generation in endothelial 512  
cells exposed to oxLDL (Fig. 2). 513

The relationship between NO and cellular inflammation is 514  
complex because NO is cytotoxic at high concentrations and has a 515  
protective effect at low concentrations. High levels of NO have the 516  
opportunity to react with superoxide leading to peroxynitrite 517  
formation and cell toxicity [33]. It has been shown that iNOS-derived 518  
overproduction of NO can lead to activation of NF-κB, which in turn 519  
leads to the up-regulation of several major proinflammatory 520  
mediators such as COX-2, iNOS and the adhesion molecules. In the 521  
present study, we found that pretreatment of CoQ10 suppressed the 522  
oxLDL-induced down-regulation of eNOS and up-regulation of iNOS. 523  
Furthermore, our data revealed that 1400W but not L-NIO represses 524  
NO formation (Fig. 3C), indicating that oxLDL-facilitated NO gener- 525  
ation mainly through iNOS. In addition, pretreatment with L-NIO 526  
partially antagonized the effect of CoQ10 on inhibition of NF-κB, and 527  
pretreatment with 1400W or SNP partially reversed the oxLDL- 528  
induced activation of NF-κB, suggesting that the protective effect of 529  
CoQ10 is due to its ability to modulate the NO-mediated signaling 530  
pathway (Fig. 5A). 531

Intracellular calcium signaling is associated with the development 532  
of atherosclerosis and several cardiovascular diseases. Vindis et al. 533

[34] showed that oxLDL-induced apoptosis in endothelial cells involves two distinct calcium-dependent pathways, the first mediated by calpain/mPTP/cytochrome *c*/caspase-3 and the second mediated by apoptosis-inducing factor, which is caspase-independent. Clinical studies have shown that calcium channel blockers, such as azelnidipine and dihydropyridine, protect against oxidative stress-related injuries by inhibiting intracellular ROS generation, mitigating NF- $\kappa$ B activation and inhibiting the expression of adhesion molecules and proinflammatory factors [35,36]. We found that CoQ10 protects against oxLDL-induced apoptosis by suppressing the oxLDL-induced rise in intracellular calcium.

Endothelial cell damage induced by oxLDL was mediated through the mitochondrial-dependent apoptotic pathway. There is evidence that NF- $\kappa$ B activation is accompanied by elevated P53 levels, indicating that NF- $\kappa$ B might play a critical role in regulating P53 [37], which subsequently induces a conformational change in Bax that enables the mitochondrial translocation of that proapoptotic protein [31]. Our results showed that CoQ10 significantly reduced the activation of NF- $\kappa$ B, suppressed the expression of P53 and the expression of Bax and led to a significant increase in expression of the antiapoptotic protein Bcl-2, which prevented the release of cytochrome *c* and further activation of caspase 3. It has been linked to the inhibition of caspase activation and prevention of endothelial apoptosis caused by oxLDL [38]. Our findings that CoQ10 reduced the activity of caspase 3 in oxLDL-treated HUVECs and that inhibitors of eNOS partially blocked this effect clearly identify the antiapoptotic effect of CoQ10 is due, at least in part, to its ability to modulate the NO-mediated signaling pathway.

The concentrations (2.5–20  $\mu$ M) we used to attenuate oxLDL-induced endothelial cell dysfunction are similar to those investigated to mitigate other oxidative stress-related responses. For example, 10  $\mu$ M of CoQ10 was shown to inhibit high glucose-induced endothelial cell adhesion molecule expression, restore high glucose-mitigated eNOS expression and attenuate high-glucose-induced ROS generation and endothelial cell apoptosis [39]. The dosage of 3000 mg/day, which reach a plasma concentration of 8.69  $\mu$ M, is safe and tolerable in patients with Parkinson's disease [40]. In this study, pretreatment with 2.5  $\mu$ M, a physiologically achievable concentration, was enough to mitigate oxLDL-induced ROS generation, restore the suppression of antioxidant enzyme activities induced by oxLDL and suppress the activation of ROS-mediated proapoptotic signaling pathways.

In summary, we have demonstrated that CoQ10 suppressed oxidative stress-related responses by modulating NO-related signaling. Our results provide insight into some of the mechanisms by which CoQ10 protects against endothelial damage.

## Acknowledgments

Flow cytometry was performed at the Instrument center of Chung-Shan Medical University, which is partly supported by National Science Council, Ministry of Education and Chung-Shan Medical University.

## References

[1] Lum H, Roebuck KA. Oxidant stress and endothelial cell dysfunction. *Am J Physiol Cell Physiol* 2001;280(4):C719–41.

[2] Chen XL, Zhang Q, Zhao R, Ding X, Tummala PE, Medford RM. Rac1 and superoxide are required for the expression of cell adhesion molecules induced by tumor necrosis factor- $\alpha$  in endothelial cells. *J Pharmacol Exp Ther* 2003;305(2):573–80.

[3] Chen XL, Zhang Q, Zhao R, Medford RM. Superoxide, H<sub>2</sub>O<sub>2</sub>, and iron are required for TNF- $\alpha$ -induced MCP-1 gene expression in endothelial cells: role of Rac1 and NADPH oxidase. *Am J Physiol Heart Circ Physiol* 2004;286(3):H1001–7.

[4] Cominacini L, Pasini AF, Garbin U, Davoli A, Tosetti ML, Campagnola M, et al. Oxidized low density lipoprotein (ox-LDL) binding to ox-LDL receptor-1 in endothelial cells induces the activation of NF- $\kappa$ B through an increased

production of intracellular reactive oxygen species. *J Biol Chem* 2000;275(17):12633–8.

[5] Dimmeler S, Zeiher AM. Reactive oxygen species and vascular cell apoptosis in response to angiotensin II and pro-atherosclerotic factors. *Regul Pept* 2000;90(1–3):19–25.

[6] Crane FL. Biochemical functions of coenzyme Q10. *J Am Coll Nutr* 2001;20(6):591–8.

[7] Kalen A, Appelkvist EL, Dallner G. Age-related changes in the lipid compositions of rat and human tissues. *Lipids* 1989;24(7):579–84.

[8] Sarter B. Coenzyme Q10 and cardiovascular disease: a review. *J Cardiovasc Nurs* 2002;16(4):9–20.

[9] Potgieter M, Pretorius E, Oberholzer HM. Qualitative electron microscopic analysis of cultured chick embryonic cardiac and skeletal muscle cells: the cellular effect of coenzyme q10 after exposure to triton x-100. *Ultrastruct Pathol* 2009;33(3):93–101.

[10] Schmelzer C, Lindner I, Rimbach G, Niklowitz P, Menke T, Doring F. Functions of coenzyme Q10 in inflammation and gene expression. *Biofactors* 2008;32(1–4):179–83.

[11] Kon M, Kimura F, Akimoto T, Tanabe K, Murase Y, Ikemune S, et al. Effect of coenzyme Q(10) supplementation on exercise-induced muscular injury of rats. *Exerc Immunol Rev* 2007;13:76–88.

[12] Murad LB, Guimaraes MR, Vianna LM. Effects of decylubiquinone (coenzyme Q10 analog) supplementation on SHRSP. *Biofactors* 2007;30(1):13–8.

[13] Wang H, Zhao X, Yin S. Effects of coenzyme Q10 or combined with micronutrients on antioxidant defense system in rats. *Wei Sheng Yan Jiu* 2008;37(3):311–3.

[14] Sohet FM, Neyrinck AM, Pachikian BD, de Backer FC, Bindels LB, Niklowitz P, et al. Coenzyme Q10 supplementation lowers hepatic oxidative stress and inflammation associated with diet-induced obesity in mice. *Biochem Pharmacol* 2009. 624

[15] Hamilton SJ, Chew GT, Watts GF. Coenzyme Q10 improves endothelial dysfunction in statin-treated type 2 diabetic patients. *Diabetes Care* 2009;32(5):810–2.

[16] Rosenfeldt FL, Pepe S, Linnane A, Nagley P, Rowland M, Ou R, et al. Coenzyme Q10 protects the aging heart against stress: studies in rats, human tissues, and patients. *Ann NY Acad Sci* 2002;959:355–9 [discussion 463–355]. 630

[17] Mabuchi H, Nohara A, Kobayashi J, Kawashiri MA, Katsuda S, Inazu A, et al. Effects of CoQ10 supplementation on plasma lipoprotein lipid, CoQ10 and liver and muscle enzyme levels in hypercholesterolemic patients treated with atorvastatin: a randomized double-blind study. *Atherosclerosis* 2007;195(2):e182–9. 634

[18] Chapidze G, Kapanadze S, Dolidze N, Bachutashvili Z, Latsabidze N. Prevention of coronary atherosclerosis by the use of combination therapy with antioxidant coenzyme Q10 and statins. *Georgian Med News* 2005(118):20–5. 636

[19] Chapidze GE, Kapanadze SD, Dolidze NK, Latsabidze NE, Bakhtushvili ZV. Combination treatment with coenzyme Q10 and simvastatin in patients with coronary atherosclerosis. *Kardiologia* 2006;46(8):11–3. 640

[20] Ou HC, Song TY, Yeh YC, Huang CY, Yang SF, Chiu TH, et al. EGCG protects against oxidized LDL-induced endothelial dysfunction by inhibiting LOX-1-mediated signaling. *J Appl Physiol* 2010;108(6):1745–56. 642

[21] Kuo WW, Huang CY, Chung JG, Yang SF, Tsai KL, Chiu TH, et al. Crude extracts of Solanum lyratum protect endothelial cells against oxidized low-density lipoprotein-induced injury by direct antioxidant action. *J Vasc Surg* 2009;50(4):849–60. 645

[22] Denizot F, Lang R. Rapid colorimetric assay for cell growth and survival. Modifications to the tetrazolium dye procedure giving improved sensitivity and reliability. *J Immunol Methods* 1986;89(2):271–7. 649

[23] Gavrieli Y, Sherman Y, Ben-Sasson SA. Identification of programmed cell death in situ via specific labeling of nuclear DNA fragmentation. *J Cell Biol* 1992;119(3):493–501. 652

[24] Ding AH, Nathan CF, Stuehr DJ. Release of reactive nitrogen intermediates and reactive oxygen intermediates from mouse peritoneal macrophages. Comparison of activating cytokines and evidence for independent production. *J Immunol* 1988;141(7):2407–12. 656

[25] Grynkiewicz G, Poenie M, Tsien RY. A new generation of Ca<sup>2+</sup> indicators with greatly improved fluorescence properties. *J Biol Chem* 1985;260(6):3440–50. 658

[26] Bedner E, Li X, Gorczyca W, Melamed MR, Darzynkiewicz Z. Analysis of apoptosis by laser scanning cytometry. *Cytometry* 1999;35(3):181–95. 660

[27] Lubrano V, Baldi S, Ferrannini E, L'Abbate A, Natali A. Role of thromboxane A2 receptor on the effects of oxidized LDL on microvascular endothelium nitric oxide, endothelin-1, and IL-6 production. *Microcirculation* 2008;15(6):543–53. 663

[28] Robbesyn F, Salvayre R, Negre-Salvayre A. Dual role of oxidized LDL on the NF- $\kappa$ B signaling pathway. *Free Radic Res* 2004;38(6):541–51. 665

[29] Marshall HE, Merchant K, Stamler JS. Nitrosation and oxidation in the regulation of gene expression. *FASEB J* 2000;14(13):1889–900. 667

[30] Sanson M, Ingueneau C, Vindis C, Thiers JC, Glock Y, Rousseau H, et al. Oxygen-regulated protein-150 prevents calcium homeostasis deregulation and apoptosis induced by oxidized LDL in vascular cells. *Cell Death Differ* 2008;15(8):1255–65. 670

[31] Cheng J, Cui R, Chen CH, Du J. Oxidized low-density lipoprotein stimulates p53-dependent activation of proapoptotic bax leading to apoptosis of differentiated endothelial progenitor cells. *Endocrinology* 2007;148(5):2085–94. 673

[32] Rauscher FM, Sanders RA, Watkins III JB. Effects of coenzyme Q10 treatment on antioxidant pathways in normal and streptozotocin-induced diabetic rats. *J Biochem Mol Toxicol* 2001;15(1):41–6. 676

[33] Zou MH, Leist M, Ullrich V. Selective nitration of prostacyclin synthase and defective vasorelaxation in atherosclerotic bovine coronary arteries. *Am J Pathol* 1999;154(5):1359–65. 679

[34] Vindis C, Elbaz M, Escargueil-Blanc I, Auge N, Heniquez A, Thiers JC, et al. Two distinct calcium-dependent mitochondrial pathways are involved in 681

- 682 oxidized LDL-induced apoptosis. *Arterioscler Thromb Vasc Biol* 2005;25(3):  
683 639-45.
- 684 [35] Naito Y, Shimozawa M, Manabe H, Nakabe N, Katada K, Kokura S, et al.  
685 Azelnidipine, a new calcium channel blocker, inhibits endothelial inflammatory  
686 response by reducing intracellular levels of reactive oxygen species. *Eur J*  
687 *Pharmacol* 2006;546(1-3):11-8.
- 688 [36] Takayama M, Yao K, Wada M. The dihydropyridine calcium channel blocker  
689 benidipine prevents lysophosphatidylcholine-induced endothelial dysfunction in  
690 rat aorta. *J Biomed Sci* 2009;16:57.
- 691 [37] Dentelli P, Rosso A, Zeoli A, Gambino R, Pegoraro L, Pagano G, et al. Oxidative  
692 stress-mediated mesangial cell proliferation requires RAC-1/reactive oxygen  
704 species production and beta4 integrin expression. *J Biol Chem* 2007;282(36): 693  
26101-10. 694
- [38] Hoffmann J, Haendeler J, Aicher A, Rossig L, Vasa M, Zeiher AM, et al. Aging  
695 enhances the sensitivity of endothelial cells toward apoptotic stimuli: important  
696 role of nitric oxide. *Circ Res* 2001;89(8):709-15. 697
- [39] Tsuneki H, Sekizaki N, Suzuki T, Kobayashi S, Wada T, Okamoto T, et al. Coenzyme  
698 Q10 prevents high glucose-induced oxidative stress in human umbilical vein  
699 endothelial cells. *Eur J Pharmacol* 2007;566(1-3):1-10. 700
- [40] Shults CW, Flint Beal M, Song D, Fontaine D. Pilot trial of high dosages of  
701 coenzyme Q10 in patients with Parkinson's disease. *Exp Neurol* 2004;188(2): 702  
491-4. 703

UNCORRECTED PROOF

**AUTHOR QUERY FORM**

 ELSEVIER	<b>Journal: JNB</b>  <b>Article Number: 6676</b>	<b>Please e-mail or fax your responses and any corrections to:</b> <b>Jill Shepherd</b> <b>E-mail: <a href="mailto:J.Shepherd@Elsevier.com">J.Shepherd@Elsevier.com</a></b> <b>Tel: 352-483-8113</b> <b>Fax: 352-483-3417</b>
---	--	---

Dear Author,

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list.

For correction or revision of any artwork, please consult <http://www.elsevier.com/artworkinstructions>.

No queries have arisen during the processing of your article.

Thank you for your assistance.