

RECURRENT DISTAL 16Q DUPLICATION AND TERMINAL 22Q DELETION: PRENATAL DIAGNOSIS AND GENETIC COUNSELING

Chih-Ping Chen^{1,2,3,4,5,6*}, Ming-Chao Huang¹, Yi-Ning Su⁷, Fuu-Jen Tsai^{4,8}, Pei-Chen Wu¹,
Chen-Chi Lee¹, Dai-Dyi Town¹, Chen-Wen Pan¹, Wayseen Wang^{2,9}

*Departments of*¹*Obstetrics and Gynecology and*²*Medical Research, Mackay Memorial Hospital,*⁵*Institute of Clinical and Community Health Nursing, and*⁶*Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University,*⁷*Department of Medical Genetics, National Taiwan University Hospital, and*⁹*Department of Bioengineering, Tatung University, Taipei;*³*Department of Biotechnology, Asia University,*⁴*School of Chinese Medicine, College of Chinese Medicine, China Medical University, and*⁸*Departments of Medical Genetics and Medical Research, China Medical University Hospital, Taichung, Taiwan.*

A 36-year-old woman, gravida 4, para 0, was referred for amniocentesis at 18 gestational weeks because of advanced maternal age and an autosomal reciprocal translocation in her second spouse. This was the woman's fourth pregnancy, and she had experienced two preterm deliveries with neonatal death during her previous marriage and one spontaneous abortion following a relationship with her current spouse. Nine years before, the ex-wife of the woman's current spouse gave birth to a growth-restricted malformed baby at term with a karyotype of 46,XX,der(22)t(16;22)(q12.1;q13.3) and an unbalanced reciprocal translocation between 16q and 22q [1]. The baby's chromosomal aberration led to the diagnosis of a 46,XY,t(16;22)(q12.1;q13.3) karyotype in the man. Four years later, the man's ex-wife delivered a malformed baby again with a karyotype of 46,XX,der(22)t(16;22)(q12.1;q13.3) [2]. During this pregnancy, amniocentesis at 18 gestational weeks revealed a karyotype of 46,XX,der(22)t(16;22)(q12.1;q13.3) (Figure 1). Level II ultrasound revealed a singleton with fetal biometry consistent with the gestational age, dolichocephaly, decreased fetal movements and a thickened nuchal fold. The pregnancy was subsequently terminated. At 22 gestational weeks, a 342 g malformed female fetus was delivered with a high forehead, bitemporal narrowing, frontal bossing, dolichocephaly, a prominent nose, hypertelorism, large low-set ears, micrognathia, a short neck with a thickened nuchal fold and

clinodactyly (Figure 2). Array comparative genomic hybridization (aCGH) using genomic DNA extracted from the uncultured umbilical cord confirmed a distal 16q duplication and a terminal 22q deletion (Figure 3).

We previously reported the prenatal diagnosis of an inherited unbalanced reciprocal translocation by aCGH using uncultured amniocytes [3]. Our case demonstrates that aCGH can be applied for rapid confirmation of prenatally diagnosed aneuploidy using uncultured postnatal tissues. The present case manifested some of the characteristic features associated with 22q13.3 deletion syndrome and partial trisomy 16q. The 22q13.3 deletion syndrome or Phelan-McDermid syndrome (OMIM 606232) is characterized by long eye lashes, large or unusual ears, relatively large hands, dysplastic toenails, a full brow, dolichocephaly, ptosis, full cheeks, a bulbous nose, a pointed chin, autistic behavior, neonatal hypotonia, global developmental delay, normal or accelerated growth and absent to severely delayed speech [4]. The reported abnormal findings of trisomy 16q on prenatal ultrasound include hydrocephalus, intrauterine growth restriction, micrognathia, congenital heart defects, clinodactyly and abnormal external genitalia [5,6]. The present case had haploinsufficiency of the *SHANK3* gene (49,459,936–49,518,507 bp) (NCBI Build 36). Haploinsufficiency of the *SHANK3* gene is a major cause of the neurological symptoms of the 22q13 deletion syndrome [7,8]. The abnormal gene dosage of *SHANK3* is associated with autism spectrum disorders and language and speech disorders [8].

In the present case, the reason for prenatal chromosome analysis was the paternal carrier status, which was primarily ascertained through a previous term aneuploid child. Inherited unbalanced structural chromosomal abnormalities detected at prenatal chromosome analysis



ELSEVIER

*Correspondence to: Dr Chih-Ping Chen, Department of Obstetrics and Gynecology, Mackay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei, Taiwan.

E-mail: cpc_mmh@yahoo.com

Accepted: July 28, 2010




Figure 1. 46,XX,der(22)t(16;22)(q12.1;q13.3) karyotype in the fetus. The arrows indicate the breakpoints.




Figure 2. (A) Anterior view and (B) lateral view of a fetus with dolichocephaly, bitemporal narrowing, a short neck, frontal bossing, a prominent nose, hypertelorism, large low-set ears and micrognathia, and (C) clinodactyly.




Figure 3. Whole genome view of array comparative genomic hybridization shows a distal 16q duplication (arrows at chromosome 16) and a terminal 22q deletion (arrows at chromosome 22).

are most commonly ascertained through a previous child with an unbalanced karyotype [9]. In a study of 56 inherited unbalanced structural chromosomal abnormalities detected at prenatal chromosome analysis, Franssen et al [9] found that the modes of ascertainment

were a previous child with an unbalanced karyotype (48%, 27/56), ultrasound abnormalities (20%, 11/56), advanced maternal age (9%, 5/56), abnormal serum screening (4%, 2/56), abnormal nuchal translucency (4%, 2/56), intracytoplasmic sperm injection (4%, 2/56),

and recurrent miscarriage (2%, 1/56), among others. Structural chromosomal rearrangements in couples are usually identified through recurrent miscarriages, previous aneuploid live birth or stillbirth, and prenatal diagnosis of chromosomal aberrations. Chen et al [10,11] found that 17 of 22 (77.3%) families with prenatal diagnosis of an inherited acrocentric rearrangement by amniocentesis were aware of their carrier status only after the diagnosis of a fetus with a translocation.

Structural chromosomal rearrangements in couples are estimated to be 2.2% after one miscarriage, 4.8% after two miscarriages and 5.2% after three miscarriages compared with 0.7% in the general population [12–14]. In a study of reproductive outcome after chromosome analysis in couples with two or more miscarriages, Franssen et al [15] concluded that the risk (0.4% at prenatal diagnosis and 0.4% at birth) of viable offspring with chromosomal abnormalities was low, and that the chance of having a healthy child (83%) was as high as that of non-carrier couples (84%) despite a higher risk of a subsequent miscarriage. Nonetheless, carrier couples ascertained through a previous aneuploid child are at a higher risk of having unbalanced viable offspring than those ascertained through miscarriages [15–18]. Boué and Gallano [16] reported a 20.8% risk of unbalanced fetuses at prenatal diagnosis when the translocation was ascertained through an aneuploid infant, compared with a risk of 4.9% when the ascertainment was through other means. Daniel et al [17] reported a 20–25% risk of unbalanced pregnancy in carriers of 2:2 segregating reciprocal translocations ascertained by previous term unbalanced offspring irrespective of carrier parents. Barišić et al [18] reported a 31.6% risk of an unbalanced fetal karyotype at prenatal diagnosis when the translocation was determined through an aneuploid infant, compared with a risk of 11.8% when the ascertainment was through spontaneous abortions. Therefore, for carrier couples whose carrier status is determined through a previous aneuploid child, pre-implantation genetic diagnosis by fluorescence *in situ* hybridization [19] or by whole genome amplification [20,21] may be an alternative to prenatal diagnosis in cases where there are multiple failed attempts to achieve a successful pregnancy.

Acknowledgments

This work was supported by research grants NSC-96-2314-B-195-008-MY3 and NSC-97-2314-B-195-006-MY3 from the National Science Council, and MMH-E-99004 from Mackay Memorial Hospital, Taipei, Taiwan.

References

1. Chen CP, Lin SP, Chern SR, Shih SL, Lee CC, Wang W, Liao SW. Perinatal findings and molecular cytogenetic analysis of trisomy 16q and 22q13.3 deletion. *Prenat Diagn* 2003;23:504–8.
2. Chen CP, Hsu CY, Huang JK, Lee CC, Chen WL, Wang W. Prenatal diagnosis of partial trisomy 16q and distal 22q13 deletion associated with dolichocephaly and frontal bossing on second-trimester ultrasound. *Prenat Diagn* 2005;25:964–6.
3. Chen CP, Su YN, Tsai FJ, et al. Terminal 2q deletion and distal 15q duplication: prenatal diagnosis by array comparative genomic hybridization using uncultured amniocytes. *Taiwan J Obstet Gynecol* 2009a;48:441–5.
4. Phelan MC. Deletion 22q13.3 syndrome. *Orphanet J Rare Dis* 2008;3:14.
5. Bianchi DW, Nicholls RD, Russell KA, Miller WA, Ellin M, Lage JM. Pericentric inversion of chromosome 16 in a large kindred: spectrum of morbidity and mortality in offspring. *Am J Med Genet* 1992;43:791–5.
6. Paladini D, D'Agostino A, Liguori M, Teodoro A, Tartaglione A, Colombari S, Martinelli P. Prenatal findings in trisomy 16q of paternal origin. *Prenat Diagn* 1999;19:472–5.
7. Wilson HL, Wong AC, Shaw SR, et al. Molecular characterisation of the 22q13 deletion syndrome supports the role of haploinsufficiency of SHANK3/PROSAP2 in the major neurological symptoms. *J Med Genet* 2003;40:575–84.
8. Durand CM, Betancur C, Boeckers TM, et al. Mutations in the gene encoding the synaptic scaffolding protein SHANK3 are associated with autism spectrum disorders. *Nat Genet* 2007;39:25–7.
9. Franssen MT, Korevaar JC, Tjoa WM, et al. Inherited unbalanced structural chromosome abnormalities at prenatal chromosome analysis are rarely ascertained through recurrent miscarriage. *Prenat Diagn* 2008;28:408–11.
10. Chen CP, Chern SR, Wu PC, et al. Unbalanced and balanced acrocentric rearrangements involving chromosomes other than chromosome 21 at amniocentesis. *Taiwan J Obstet Gynecol* 2009b;48:389–99.
11. Chen CP, Chern SR, Tsai FJ, et al. Unbalanced and balanced heterologous Robertsonian translocations involving chromosome 21 at amniocentesis. *Taiwan J Obstet Gynecol* 2010;49:62–8.
12. Hook EB, Healy NP, Willey AM. How much difference does chromosome banding make? Adjustments in prevalence and mutation rates of human structural cytogenetic abnormalities. *Ann Hum Genet* 1989;53:237–42.
13. De Braekeleer M, Dao TN. Cytogenetic studies in couples experiencing repeated pregnancy losses. *Hum Reprod* 1990;5:519–28.
14. Franssen MT, Korevaar JC, Leschot NJ, et al. Selective chromosome analysis in couples with two or more miscarriages: case-control study. *BMJ* 2005;331:137–41.
15. Franssen MT, Korevaar JC, van der Veen F, Leschot NJ, Bossuyt PMM, Goddijn M. Reproductive outcome after chromosome analysis in couples with two or more miscarriages: case-control study. *BMJ* 2006;332:759–63.

16. Boué A, Gallano P. A collaborative study of the segregation of inherited chromosome structural rearrangements in 1356 prenatal diagnoses. *Prenat Diagn* 1984;4:45–67.
17. Daniel A, Hook EB, Wulf G. Risks of unbalanced progeny at amniocentesis to carriers of chromosome rearrangements: data from United States and Canadian laboratories. *Am J Med Genet* 1989;33:14–53.
18. Barišić I, Zergollern L, Mu inić D, Hitrec V. Risk estimates for balanced reciprocal translocation carriers—prenatal diagnosis experience. *Clin Genet* 1996;49:145–51.
19. Holmes LB, Gargiulo AR, Nadel AS, Racowsky C. Case 11-2005. A 32-year-old pregnant woman with an abnormal fetal karyotype. *N Engl J Med* 2005;352:1579–87.
20. Delhanty JDA, Wells D. Preimplantation genetic diagnosis: an alternative to prenatal diagnosis. *Expert Rev Mol Diagn* 2002;2:395–9.
21. Peng W, Takabayashi H, Ikawa K. Whole genome amplification from single cells in preimplantation genetic diagnosis and prenatal diagnosis. *Eur J Obstet Gynecol Reprod Biol* 2007; 131:13–20.