

1 Anti-angiogenic potential of three triterpenic acids in human liver cancer cells

2
3
4 CHUN-CHE LIN^{o,∇,#}, CHUN-YIN HUANG^{‡,#}, MEI-CHIN MONG[†], CHIEN-YI CHAN[‡],
5 MEI-CHIN YIN^{†,‡,*}

6 ^oDepartment of Internal Medicine, Chung Shan Medical University Hospital, Taichung City,
7 Taiwan

8 [∇]Institute of Medicine, Chung Shan Medical University, Taichung City, Taiwan

9 [‡]Department of Nutrition, China Medical University, Taichung City, Taiwan

10 [†]Department of Health and Nutrition Biotechnology, Asia University, Taichung County,
11 Taiwan

12
13
14 running title: anti-angiogenic potential of triterpenic acids

15 [#]These two authors contributed equally to this study.

16 *To whom correspondence should be addressed: Dr. Mei-chin Yin, Professor, Department
17 of Nutrition, China Medical University, 16th Floor, 91, Hsueh-shih Rd, Taichung City,
18 Taiwan, ROC

19 TEL: 886-4-22053366 ext. 7510, FAX: 886-4-22062891

20 Email: meyin@mail.cmu.edu.tw

1 **ABSTRACT**

2 Three triterpenic acids, oleanolic acid, ursolic acid and maslinic acid, at 2 or 4 $\mu\text{mol/L}$
3 were used to study their anti-angiogenic potential in human liver cancer Hep3B, Huh7 and
4 HA22T cell lines. The effects of these compounds upon the level and/or expression of
5 hypoxia-inducible factor (HIF)-1 α , basic fibroblast growth factor (bFGF), vascular
6 endothelial growth factor (VEGF), interleukin (IL)-8, urokinase plasminogen activator
7 (uPA), reactive oxygen species (ROS), nitric oxide (NO) and cell invasion and migration
8 were examined. Results showed that these triterpenic acids at 4 $\mu\text{mol/L}$ significantly
9 suppressed HIF-1 α expression in three cell lines ($P<0.05$); and these compounds at test
10 doses failed to affect bFGF expression ($P>0.05$). Three triterpenic acids
11 dose-dependently decreased production and expression of VEGF and IL-8, retained
12 glutathione level, lowered ROS and NO levels, and declined cell invasion and migration in
13 test cell lines ($P<0.05$). These compounds also dose-dependently reduced uPA production
14 and expression in Hep3B and Huh7 cell lines ($P<0.05$); but these agents only at 4 $\mu\text{mol/L}$
15 significantly suppressed uPA production and expression in HA22T cells ($P<0.05$). These
16 findings suggest that these triterpenic acids are potent anti-angiogenic agents to retard
17 invasion and migration in liver cancer cells.

18

19 **KEYWORDS:** angiogenesis; triterpenic acid; liver cancer cell; hypoxia-inducible factor;
20 vascular endothelial growth factor

21

1 INTRODUCTION

2 Angiogenesis, also called as neovascularization, is an important process for tumor
3 growth and metastasis. Liver cancer, also called as hepatocellular carcinoma (HCC), is a
4 hypervascular tumor characterized by massive angiogenesis (1). The increased
5 circulating levels and/or expression of several angiogenetic factors such as vascular
6 endothelial growth factor (VEGF) and interleukin (IL)-8 in patients with HCC have been
7 reported (2, 3). Hypoxia-inducible factor (HIF)-1 α and basic fibroblast growth factor
8 (bFGF) are two crucial activators responsible for cancer angiogenesis, in which HIF-1 α
9 regulates the essential adaptive responses of cancer cells to hypoxia, the major
10 pathophysiological condition promoting angiogenesis; and bFGF mediates extracellular
11 matrix degradation (4, 5). In addition, VEGF through increasing vascular permeability
12 could enhance endothelial cell migration and proliferation, which are required for vascular
13 remodeling and sprouting development of new blood vessels (6). Overexpression of
14 urokinase plasminogen activator (uPA), a gene related to invasion, also contributed to HCC
15 angiogenesis (7). On the other hand, reactive oxygen species (ROS) and nitric oxide (NO)
16 play important roles in cancer angiogenesis because these free radicals enhance oxidative
17 stress and stimulate VEGF production, which in turn favors cancer progression (8, 9).
18 Obviously, these factors provide vascular support for the growth of already established
19 malignant tumor. Thus, any agents with suppressive effects on the production or
20 expression of these angiogenetic factors may be beneficial in retarding tumor progression.

21 Oleanolic acid, ursolic acid and maslinic acid are pentacyclic triterpenic acids
22 naturally occurring in many herbs and plant foods such as glossy privet fruit (*Ligustrum*
23 *lucidum* Ait.), hawthorn fruit (*Crataegi Pinnatifidae* Fructus) and olive (10, 11). The
24 inhibitory effects of oleanolic acid and ursolic acid against the growth of human bladder,
25 prostate and colorectal cancer cells have been observed (12, 13). Our previous study (14)

1 also found that these two triterpenic acids at 8 $\mu\text{mol/L}$ caused markedly apoptotic effects in
2 four human liver cancer HepG2, Hep3B, Huh7 and HA22T cell lines via increasing DNA
3 fragmentation, decreasing mitochondrial membrane potential and lowering
4 $\text{Na}^+ - \text{K}^+ - \text{ATPase}$ activity. Juan et al. (15) reported that maslinic acid, another triterpenic
5 acid, could inhibit cell proliferation in human colon cancer cell, HT-29. These studies
6 suggest that these triterpenic acids are potent anti-cancer agents to cause apoptosis or
7 retard metastasis for liver cancer. Therefore, a further study was conducted to examine
8 the anti-angiogenic potential of these triterpenic acids. The major purpose of our present
9 study was to investigate the effects of triterpenic acids on the production and/or expression
10 of HIF-1 α , bFGF, VEGF, IL-8 and uPA in liver cancer cell lines. The influence of these
11 compounds upon the production of ROS and NO in these cell lines was also evaluated.

12 On the other hand, the presence of ursolic acid and maslinic acid in apple peel has
13 been considered as major compounds responsible for the anti-cancer activity of apple (16).
14 So far, less information is available regarding the presence of these triterpenic acids in
15 fresh vegetables. In order to enrich the food component database, eleven locally available
16 fresh vegetables in Taiwan were used to analyze the content of oleanolic acid, ursolic acid
17 and maslinic acid.

18

19 **MATERIALS AND METHODS**

20 **Chemicals.** Oleanolic acid (OA, 99%), ursolic acid (UA, 98%) and maslinic acid
21 (MA, 98.5%) were purchased from Aldrich Chemical Co. (Milwaukee, WI, USA).
22 Medium, plates, antibiotics and chemicals used for cell culture were purchased from Difco
23 Laboratory (Detroit, MI, USA). All chemicals used in these measurements were of the
24 highest purity commercially available.

1 **Content of Triterpenic Acids.** Eleven locally available fresh vegetables, carrot
2 (*Daucus carota* L.), celery (*Apium graveolens* L.), okra (*Hibiscus esculentus*), eggplant
3 (*Solanum melongena* L.), spinach (*Spinacia oleracea* L.), leaf mustard (*Brassica juncea*),
4 gynura (*Gynura bicolor* DC.), Chinese chive (*Allium tuberosum* Rottler), fennel
5 (*Foeniculum vulgare*), basil (*Ocimum basilicum* L.) and daylily (*Hemerocallis fulva* L.),
6 were used to analyze the content of OA, UA and MA. These vegetables, harvested in
7 summer of 2010, were purchased from farms in central area of Taiwan. The content of
8 three triterpenic acids in these vegetables was analyzed by HPLC methods described in Liu
9 et al. (17).

10 **Cell Culture.** Human normal liver cell line (L-02 cell) and liver cancer Hep3B, Huh7
11 and HA22T cell lines were obtained from American Type Culture Collection (ATCC,
12 Rockville, MD, USA). All cells were cultured in RPMI 1640 medium, containing 10%
13 fetal bovine serum (BSA), 100 units/mL of penicillin and 100 units/mL of streptomycin
14 (pH 7.4). Hypoxia (1% O₂) was created using an anaerobic jar (AnaeroPack Series,
15 Mitsubishi Gas Chemical, Tokyo, Japan) with an AnaeroPack disposable O₂-absorbing and
16 CO₂-generating agent. The culture medium was changed every three days, and cells were
17 subcultured once a week. A phosphate buffer saline (PBS, pH 7.2) was added to adjust
18 the cell number to 10⁵/mL for various experiments and analyses.

19 **Experimental Design.** Stock solution of OA, UA or MA was prepared in dimethyl
20 sulfoxide (DMSO) and diluted with medium. An equal volume of DMSO (final
21 concentration <0.5%) was added to the controls. Our previous study (14) found that OA
22 and UA at 2 and 4 μmol/L caused 55-70% cell viability in Hep3B, Huh7 and HA22T cells
23 (8 μmol/L caused 20-30% cell viability). Thus, in order to observe the effects of these
24 triterpenic acids upon angiogenic factors in living cells, these compounds at these two
25 doses were used for present study. The cells at 10⁵/mL were treated with OA, UA or MA

1 at 2 or 4 $\mu\text{mol/L}$ under hypoxia at 37 °C for 72 h. Control group contained no OA, UA
2 or MA.

3 **Measurement of Glutathione (GSH), Reactive Oxygen Species (ROS) and Nitrite.**

4 After 72 h hypoxic treatment, cells were washed and suspended in RPMI 1640 medium.
5 GSH concentration (ng/mg protein) was determined by a commercial colorimetric GSH
6 assay kit (OxisResearch, Portland, OR, USA). The dye DCFH2-DA was used to measure
7 level of intracellular cellular ROS according to the method of Fu et al. (18). In briefly,
8 cells were incubated with 50 $\mu\text{mol/L}$ dye for 30 min and washed with PBS. After
9 centrifugation at 412 $\times g$ for 10 min, the medium was removed and cells were dissolved
10 with 1% Triton X-100. Fluorescence changes were measured at an excitation wavelength
11 of 485 nm and an emission wavelength of 530 nm using a fluorescence microplate reader.
12 The production of NO was determined by measuring the formation of nitrite. Briefly,
13 100 μL of supernatant was mixed with 100 μL Griess reagent after centrifuged at 10,000
14 $\times g$ for 15 min at 4 °C. The optical absorbance at 540 nm was measured and compared
15 with a sodium nitrite standard curve.

16 **Determination of Angiogenic Factors.** VEGF and IL-8 levels in cell homogenates
17 were quantified by commercial ELISA kits (R&D Systems, Minneapolis, MN, USA).
18 uPA level was assayed by commercial ELISA kit purchased from American Diagnostica
19 (Greenwich, CT, USA). Each measurement was made in duplicate, and the level of each
20 angiogenic factor was determined from a standard curve.

21 **Real Time Polymerase Chain Reaction (RT-PCR) for mRNA Expression.** Total
22 RNA was isolated from cells using Trizol reagent (Invitrogen, Life Technologies, Carlsbad,
23 CA, USA). Two μg of total RNA was used to generate cDNA, which was amplified
24 using Taq DNA polymerase. PCR was carried out in 50 μL of reaction mixture
25 containing Taq DNA polymerase buffer (20 mM Tris-HCl, pH 8.4, 50 mM KCl, 200 mM

1 dNTP, 2.5 mM MgCl₂, 0.5 mM of each primer) and 2.5 U Taq DNA polymerase. The
2 specific oligonucleotide primers for HIF-1 α , bFGF, VEGF, IL-8, uPA and
3 glyceraldehyde-3-phosphate dehydrogenase (GAPDH, the housekeeping gene) are as
4 follows. HIF-1 α : forward, 5'-CTG CTG TCT TAC TGG TCC TT-3', reverse, 5'-GTC
5 GCT TCT CCA ATT CTT AC-3'; bFGF: forward, 5'-GTG TGC TAA CCG TAC CTG
6 GC-3', reverse, 5'-CTG GTG ATT TCC TTG ACC GG-3'; VEGF, forward, 5'-ATG AAC
7 TTT CTG CTC TCT GG-3', reverse, 5'-TCA TCT CTC CTA TGT GCT GGC-3'; IL-8,
8 forward, 5'-CAC TCT CAA TCA CTC TCA GTT CTT TGA T-3'; reverse, 5'-CTT CCT
9 GAT TTC TGC AGC TCT GT-3'; uPA, forward, 5'-TCA CAC CAA GGA AGA GAA
10 TGG CCT-3', reverse 5'-AAT GAC AAC CAG CAA GAA AGC GGG-3'; GAPDH:
11 forward, 5'-TGA TGA CAT CAA GAA GGT GGT GAA G-3', reverse, 5'-CCT TGG
12 AGG CCA TGT AGG CCA T-3'. The cDNA was amplified under the following
13 reaction conditions: 95 °C for 1 min, 55 °C for 1 min, and 72 °C for 1 min. 28 cycles
14 were performed for GAPDH and 35 cycles were performed for HIF-1 α , bFGF, VEGF,
15 IL-8 and uPA. Generated fluorescence from each cycle was quantitatively analyzed by
16 using the Taqman system based on real-time sequence detection system (ABI Prism 7700;
17 Perkin-Elmer Inc., Foster City, CA, USA). In this study, mRNA level was calculated as
18 percentage of the control group.

19 **Invasion Assay.** Invasion was measured using 8 μ m pore BD BioCoat Matrigel
20 Invasion Chambers (BD Biosciences, San Jose, CA, USA), in which membrane was
21 coated with 500 ng/ μ L Matrigel. After 72 h hypoxic treatment, cells were resuspended in
22 RPMI 1640 medium containing 1% BSA, and at 10⁵/100 μ L were seeded into the upper
23 chamber. The chambers were incubated for 24 h at 37 °C in a humidified, 5% CO₂
24 atmosphere for cell invasion. Non-invasive cells were removed by Q-tip. Invasive cells
25 were stained with 0.2% crystal violet in 10% ethanol. Four independent fields of

1 invasive cells per well were photographed under the microscope to determine the number
2 of invasive cells. Data were calculated as percentage of the control group.

3 **Migration assay.** Migration was measured using a transwell chamber (Corning, NY,
4 USA). The underside of transwell membranes was precoated with fibronectin (10
5 $\mu\text{g}/\text{mL}$). After 72 h hypoxic treatment, cells were resuspended in RPMI 1640 medium
6 containing 1% BSA, and at $10^5/100 \mu\text{L}$ were seeded into the upper chamber of transwell
7 chambers separated by inserts with 6.5 mm polycarbonate filters of 8 μm pore size. After
8 24 h incubation, non-migratory cells on the upper surface of membranes were removed
9 and migratory cells were stained with 0.1% crystal violet. Four independent fields of
10 migratory cells per well were photographed under the microscope to determine the number
11 of migratory cells. Data were calculated as percentage of the control group.

12 **Statistical Analysis.** The effect of each treatment was analyzed from nine different
13 preparations (n=9). Data were reported as means \pm standard deviation (SD), and
14 subjected to analysis of variance (ANOVA). Differences among means were determined
15 by the Least Significance Difference Test with significance defined at $P<0.05$.

16

17 **RESULTS**

18 The content of three triterpenic acids in eleven fresh vegetables is shown in Table 1.
19 Among test vegetables, spinach had the highest content of OA and UA, and leaf mustard
20 had the highest MA content. As shown in Figure 1, HIF-1 α mRNA expression of three
21 liver cancer cells under hypoxic condition was greater than normoxic condition ($P<0.05$),
22 which indicated that hypoxia had been created.

23 OA, UA and MA at 4 $\mu\text{mol}/\text{L}$ significantly suppressed HIF-1 α mRNA expression in
24 three cell lines (Figure 2, $P<0.05$); and these compounds at test doses failed to affect bFGF

1 mRNA expression (Figure 3, $P>0.05$). Three triterpenic acids dose-dependently
2 decreased production and mRNA expression of VEGF and IL-8 in test cell lines (Figures 4
3 and 5, $P<0.05$). As shown in Table 2, three cancer cells had significantly lower GSH
4 level and higher production of ROS and NO than normal liver cells ($P<0.05$); however, test
5 compounds dose-dependently retained GSH level, and lowered production of ROS and NO
6 in test liver cancer cells ($P<0.05$). These compounds also dose-dependently diminished
7 invasion and migration in three test cell lines (Figure 6, $P<0.05$). As shown in Figure 7,
8 three test triterpenic acids dose-dependently reduced uPA production and mRNA
9 expression in Hep3B and Huh7 cell lines; but only at 4 $\mu\text{mol/L}$ significantly
10 down-regulated uPA production and mRNA expression in HA22T cells ($P<0.05$).

11

12 **DISCUSSION**

13 Our previous study already indicated that OA and UA exhibited marked apoptotic
14 effects in liver cancer cell lines (14). The results of our present study further found that
15 these two triterpenic acids and MA, another triterpenic acid, effectively down-regulated
16 mRNA expression of HIF-1 α , VEGF, IL-8, uPA, retained GSH level, decreased ROS and
17 NO production, as well as declined invasion and migration in three liver cancer cell lines
18 under hypoxic condition. These findings support that these triterpenic acids could
19 mitigate angiogenesis, which might attenuate liver cancer metastasis and deterioration.
20 Furthermore, we found that spinach, leaf mustard and daylily are rich in these triterpenic
21 acids. It has been suggested that vegetable intake could reduce cancer risk (19, 20). The
22 results of our present study further suggest that the vegetables rich in these triterpenic acids
23 may offer more protection against liver cancer.

24 Malignant tumor growth causes hypoxic condition within the tumor, which in turn
25 enhances the expression of HIF-1 α in order to mediate an adaptive response through

1 increasing angiogenesis. Subsequently, HIF-1 α , as an transcriptional activator,
2 up-regulates VEGF expression and leads to tumor neovascularization (21, 22). Thus,
3 regulation of HIF-1 α and/or VEGF has been considered as promising target for cancer
4 therapy. In our present study, three test triterpenic acids down-regulated HIF-1 α
5 expression at high dose, but decreased VEGF expression and production in dose-dependent
6 manner. Thus, the anti-angiogenic activity of these triterpenic acids could be partially
7 ascribed to these agents suppress HIF-1 α and VEGF expression in test cell lines.
8 However, it is interesting to find that these agents at lower dose (2 μ mol/L) failed to affect
9 HIF-1 α expression, but they lowered VEGF expression. Obviously, other upstream
10 factor(s) was involved for VEGF transcription in these triterpene-treated cancer cells.
11 bFGF, another angiogenic activator, is more potent than VEGF for stimulating the vascular
12 endothelial mitogenesis because bFGF enhances the release and activity of collagenases,
13 proteases and integrins on the extracellular membrane to form nascent microvascular
14 networks (23). In our present study, three triterpenic acids at test doses failed to lower
15 bFGF expression. These results suggested that the anti-angiogenic activity of test
16 triterpenic acids did not involve bFGF associated pathway.

17 IL-8 from tumor cells is an autocrine growth factor and acts on endothelial cells
18 through their receptors to promote migration, invasion and angiogenesis, which
19 consequently elevated metastatic potential (24). Thus, decreasing IL-8 expression and
20 production might also alleviate cancer progression. In our present study, test triterpenic
21 acids reduced IL-8 expression and production in dose-dependent manner, which
22 subsequently contributed to diminish angiogenesis. IL-8 expression could be evoked by
23 HIF-1 α and NO (25, 26). Since these triterpenic acids could suppress HIF-1 α expression
24 and reduce NO production, it seems reasonable to observe the declined expression of

1 downstream target gene, IL-8. On the other hand, enhanced IL-8 expression could
2 increase the activity of some matrix metalloproteins and facilitate tumor cell invasion (27).
3 Thus, the lowered cell invasion as we observed in triterpene-treated liver cancer cells could
4 be partially ascribed to these agents already down-regulated IL-8.

5 It has been indicated that oxidative stress enhanced the expression of both HIF-1 α and
6 VEGF in cancer cells (28). Intracellular ROS derived from NADPH oxidase or
7 mitochondria, acting a signaling messenger, is able to activate critical target molecules
8 such as intercellular adhesion molecule-1 and trigger cell adhesion,
9 epithelial-mesenchymal transition and migration in tumor cells (29). Nitrite is capable of
10 inducing VEGF expression and favors nitrite-driven formation of endothelial tubes in
11 hypoxic tumor microenvironment (30). Obviously, liver cancer cells released ROS and
12 NO to promote their migration and angiogenesis. In our present study, triterpenic acids
13 treatments effectively retained GSH level, and reduced ROS and NO release in test cell
14 lines, which in turn alleviated oxidative stress and diminished the expression of angiogenic
15 factors such as VEGF. Several studies have reported that OA, UA and MA could
16 attenuate oxidative stress via sparing GSH, inhibiting NO production and/or scavenging
17 ROS (31-33). Thus, the lower ROS and NO levels in triterpenic acid-treated groups as
18 we observed could be explained. These findings revealed that these triterpenic acids
19 could mitigate angiogenesis in liver cancer cells through their anti-oxidative actions.

20 uPA converts plasminogen to plasmin, which is responsible for the degradation of
21 fibrin and extracellular matrix components, and finally promotes cell migration and
22 invasion (34). In our present study, the production and expression of uPA in the liver
23 cancer cell lines could be down-regulated by triterpenic acid treatments, which
24 subsequently decreased plasminogen conversion and degradation of fibrin and matrix
25 proteins. In addition, ROS could stimulate uPA expression in tumor cells (35). Thus,

1 the reduced uPA expression from triterpenic acids could be partially due to these
2 compounds already decrease ROS production. Since uPA expression was suppressed, the
3 observed lower cell invasion and migration in these triterpenic acid-treated groups could be
4 explained. Furthermore, test triterpenic acids at 4 $\mu\text{mol/L}$ inhibited 50-70% cell invasion
5 and migration in these three cell lines. Thus, these agents might be able to retard HCC
6 progression via blocking metastatic process.

7 It is interesting to find that three test compounds at equal dose exhibited greater
8 suppressive effects upon VEGF and uPA expression in Hep3B and Huh7 cells than in
9 HA22T cells, but these agents were greater in declining IL-8 expression in HA22T cells
10 than in other two cell lines. It has been reported that Hep3B expressed more
11 mesenchymal proteins for mesenchymal transition, Huh7 was highly associated with
12 hepatitis B virus- or C virus-related HCC, and HA22T was a poorly-differentiated cell line
13 (36-38). Thus, these cells, based on different pathological features and biochemical
14 behavior, might exhibit different response to certain chemical(s). These findings implied
15 that selecting any agent for anti-hepatoma therapy should consider the responsible cancer
16 cell line, and/or target molecules. In addition, the results of our present study suggested
17 that any agent, like these triterpenic acids, with multiple anti-cancer actions toward
18 multiple HCC cell lines might be appropriate choice for anti-HCC therapy.

19 In summary, the content of oleanolic acid, ursolic acid and maslinic acid in daylily,
20 spinach and leaf mustard was in the range of 76-174 mg/100 g dry weight. These three
21 triterpenic acids at 4 $\mu\text{mol/L}$ markedly suppressed mRNA expression of angiogenic factors,
22 HIF-1 α , VEGF, IL-8 and uPA. These triterpenic acids also effectively alleviated
23 oxidative stress via retaining GSH level, lowering ROS and NO production, as well as
24 decreased cell invasion and migration in Hep3B, Huh7 and HA22T cell lines. These

1 findings suggest that these triterpenic acids are potent anti-angiogenic agents to retard
2 invasion and migration in liver cancer cells.

3

4 Acknowledgement

5 This study was supported by a grant from China Medical University, Taichung City,
6 Taiwan, ROC (Grant CMU99-asia-01).

1 LITERATURE CITED

- 2 1. Pang, R.; Poon, R.T. Angiogenesis and antiangiogenic therapy in hepatocellular
3 carcinoma. *Cancer Lett.* **2006**, *242*, 151-167.
- 4 2. Rosmorduc, O.; Housset, C. Hypoxia: a link between fibrogenesis, angiogenesis, and
5 carcinogenesis in liver disease. *Semin. Liver Dis.* **2010**, *30*, 258-270.
- 6 3. Battegay, E.J. Angiogenesis: mechanistic insights, neovascular diseases, and
7 therapeutic prospects. *J. Mol. Med.* **1995**, *73*, 333-346.
- 8 4. Pugh, C.W.; Ratcliffe, P.J. Regulation of angiogenesis by hypoxia: role of the HIF
9 system. *Nat. Med.* **2003**, *9*, 677-684.
- 10 5. Mise, M.; Arii, S.; Higashitaji, H.; Furutani, M.; Niwano, M.; Harada, T.; Ishigami, S.;
11 Toda, Y.; Nakayama, H.; Fukumoto, M.; Fujita, J.; Imamura, M. Clinical significance
12 of vascular endothelial growth factor and basic fibroblast growth factor gene
13 expression in liver tumor. *Hepatology* **1996**, *23*, 455-464.
- 14 6. Finn, R.S.; Zhu, A.X. Targeting angiogenesis in hepatocellular carcinoma: focus on
15 VEGF and bevacizumab. *Expert. Rev. Anticancer Ther.* **2009**, *9*, 503-509.
- 16 7. Andreasen, P.A.; Kjoller, L.; Christensen, L.; Duffy, M.J. The urokinase-type
17 plasminogen activator system in cancer metastasis: a review. *Int. J. Cancer* **1997**, *72*,
18 1-22.
- 19 8. Morbidelli, L.; Donnini, S.; Ziche, M. Role of nitric oxide in the modulation of
20 angiogenesis. *Curr. Pharm. Des.* **2003**, *9*, 521-530.
- 21 9. Wu, W.S. The signaling mechanism of ROS in tumor progression. *Cancer Metastasis*
22 *Rev.* **2006**, *25*, 695-705.
- 23 10. Cui, T.; Li, J.Z.; Kayahara, H.; Ma, L.; Wu, L.X.; Nakamura, K. Quantification of the
24 polyphenols and triterpene acids in Chinese hawthorn fruit by high-performance
25 liquid chromatography. *J. Agric. Food Chem.* **2006**, *54*, 4574-4581.

- 1 11. Zhou, C.; Chen, K.; Sun, C.; Chen, Q.; Zhang, W.; Li, X. Determination of oleanolic
2 acid, ursolic acid and amygdalin in the flower of *Eriobotrya japonica* Lindl. by HPLC.
3 *Biomed. Chromatogr.* **2007**, *21*, 755-761.
- 4 12. Kassi, E.; Papoutsi, Z.; Pratsinis, H.; Aligiannis, N.; Manoussakis, M.; Moutsatsou, P.
5 Ursolic acid, a naturally occurring triterpenoid, demonstrates anticancer activity on
6 human prostate cancer cells. *J. Cancer Res. Clin. Oncol.* **2007**, *133*, 493-500.
- 7 13. Janakiram, N.B.; Indranie, C.; Malisetty, S.V.; Jagan, P.; Steele, V.E.; Rao, C.V.
8 Chemoprevention of colon carcinogenesis by oleanolic acid and its analog in male
9 F344 rats and modulation of COX-2 and apoptosis in human colon HT-29 cancer cells.
10 *Pharm. Res.* **2008**, *25*, 2151-2157.
- 11 14. Yan, S.L.; Huang, C.Y.; Wu, S.T.; Yin, M.C. Oleanolic acid and ursolic acid induce
12 apoptosis in four human liver cancer cell lines. *Toxicol. Vitro.* **2010**, *24*, 842-848.
- 13 15. Juan, M.E.; Wenzel, U.; Ruiz-Gutierrez, V.; Daniel, H.; Planas, J.M. Olive fruit
14 extracts inhibit proliferation and induce apoptosis in HT-29 human colon cancer cells.
15 *J. Nutr.* **2006**, *136*, 2553-2557.
- 16 16. He, X.; Liu, R.H. Triterpenoids isolated from apple peels have potent antiproliferative
17 activity and may be partially responsible for apple's anticancer activity. *J. Agric. Food*
18 *Chem.* **2007**, *55*, 4366-4370.
- 19 17. Liu, H.; Shi, Y.; Wang, D.; Yang, G.; Yu, A.; Zhang, H. MECC determination of
20 oleanolic acid and ursolic acid isomers in *Ligustrum lucidum* Ait. *J. Pharm. Biomed.*
21 *Anal.* **2003**, *32*, 479-485.
- 22 18. Fu, W.; Luo, H.; Parthasarathy, S.; Mattson, M.P. Catecholamines potentiate amyloid
23 β -peptide neurotoxicity: involvement of oxidative stress, mitochondrial dysfunction,
24 and perturbed calcium homeostasis. *Neurobiol. Dis.* **1998**, *5*, 229-243.

- 1 19. Key, T.J.; Schatzkin, A.; Willett, W.C.; Allen, N.E.; Spencer, E.A.; Travis, R.C. Diet,
2 nutrition and the prevention of cancer. *Public Health Nutr.* **2004**, *7*, 187-200.
- 3 20. Khan, N.; Afaq, F.; Mukhtar, H. Cancer chemoprevention through dietary
4 antioxidants: progress and promise. *Antioxid. Redox Signal.* **2008**, *10*, 475-510.
- 5 21. Rey, S.; Semenza, G.L. Hypoxia-inducible factor-1-dependent mechanisms of
6 vascularization and vascular remodelling. *Cardiovasc. Res.* **2010**, *86*, 236-242.
- 7 22. Liao, D.; Johnson, R.S. Hypoxia: a key regulator of angiogenesis in cancer. *Cancer*
8 *Metastasis Rev.* **2007**, *26*, 281-290.
- 9 23. Rusnati, M.; Presta, M. Interaction of angiogenic basic fibroblast growth factor with
10 endothelial cell heparan sulfate proteoglycans. Biological implications in
11 neovascularization. *Int. J. Clin. Lab. Res.* **1996**, *26*, 15-23.
- 12 24. Waugh, D.J.; Wilson, C. The interleukin-8 pathway in cancer. *Clin. Cancer Res.* **2008**,
13 *14*, 6735-6741.
- 14 25. Xiong, Q.; Shi, Q.; Le, X.; Wang, B.; Xie, K. Regulation of interleukin-8 expression
15 by nitric oxide in human pancreatic adenocarcinoma. *J. Interferon Cytokine Res.* **2001**,
16 *21*, 529-537.
- 17 26. Mizukami, Y.; Jo, W.S.; Duerr, E.M.; Gala, M.; Li, J.; Zhang, X.; Zimmer, M.A.;
18 Iliopoulos, O.; Zukerberg, L.R.; Kohgo, Y.; Lynch, M.P.; Rueda, B.R.; Chung, D.C.
19 Induction of interleukin-8 preserves the angiogenic response in HIF-1alpha-deficient
20 colon cancer cells. *Nat. Med.* **2005**, *11*, 992-997.
- 21 27. Kubo, F.; Ueno, S.; Hiwatashi, K.; Sakoda, M.; Kawaida, K.; Nuruki, K.; Aikou, T.
22 Interleukin 8 in human hepatocellular carcinoma correlates with cancer cell invasion
23 of vessels but not with tumor angiogenesis. *Ann. Surg. Oncol.* **2005**, *12*, 800-807.

- 1 28. Shi, D.Y.; Xie, F.Z.; Zhai, C.; Stern, J.S.; Liu, Y.; Liu, S.L. The role of cellular
2 oxidative stress in regulating glycolysis energy metabolism in hepatoma cells. *Mol.*
3 *Cancer* **2009**, *8*, 32.
- 4 29. Wang, Z.; Li, Y.; Sarkar, F.H. Signaling mechanism(s) of reactive oxygen species in
5 Epithelial-Mesenchymal Transition reminiscent of cancer stem cells in tumor
6 progression. *Curr. Stem Cell Res. Ther.* **2010**, *5*, 74-80.
- 7 30. Frérart, F.; Lobysheva, I.; Gallez, B.; Dessy, C.; Feron, O. Vascular caveolin
8 deficiency supports the angiogenic effects of nitrite, a major end product of nitric
9 oxide metabolism in tumors. *Mol. Cancer Res.* **2009**, *7*, 1056-1063.
- 10 31. Hsu, C.L.; Hong, B.H.; Yu, Y.S.; Yen, G.C. Antioxidant and anti-inflammatory
11 effects of *Orthosiphon aristatus* and its bioactive compounds. *J. Agric. Food Chem.*
12 **2010**, *58*, 2150-2156.
- 13 32. Márquez Martín, A.; de la Puerta Vázquez, R.; Fernández-Arche, A.; Ruiz-Gutiérrez,
14 V. Suppressive effect of maslinic acid from pomace olive oil on oxidative stress and
15 cytokine production in stimulated murine macrophages. *Free Radic. Res.* **2006**, *40*,
16 295-302.
- 17 33. Tsai, S.J.; Yin, M.C. Antioxidative and anti-inflammatory protection of oleanolic acid
18 and ursolic acid in PC12 cells. *J. Food Sci.* **2008**, *73*, 174-178.
- 19 34. Stefansson, S.; McMahon, G.A.; Petitclerc, E.; Lawrence, D.A. Plasminogen activator
20 inhibitor-1 in tumor growth, angiogenesis and vascular remodeling. *Curr. Pharm. Des.*
21 **2003**, *9*, 1545-1564.
- 22 35. Lee, K.H.; Kim, J.R. Reactive oxygen species regulate the generation of urokinase
23 plasminogen activator in human hepatoma cells via MAPK pathways after treatment
24 with hepatocyte growth factor. *Exp. Mol. Med.* **2009**, *41*, 180-188.

- 1 36. Huang, E.J.; Wu, C.C.; Lee, S.D.; Chen, J.H.; Liu, J.Y.; Ko, J.L.; Lin, J.A.; Lu, M.C.;
2 Chen, L.M.; Huang, C.Y.; Kuo, W.W. Opposing action of estrogen receptors alpha
3 and beta on tumor necrosis factor alpha gene expression and caspase-8-mediated
4 apoptotic effects in HA22T cells. *Mol. Cell Biochem.* **2006**, *287*, 137-145.
- 5 37. Slany, A.; Haudek, V.J.; Zwickl, H.; Gundacker, N.C.; Grusch, M.; Weiss, T.S.; Seir,
6 K.; Rodgarkia-Dara, C.; Hellerbrand, C.; Gerner, C. Cell characterization by proteome
7 profiling applied to primary hepatocytes and hepatocyte cell lines Hep-G2 and
8 Hep-3B. *J. Proteome Res.* **2010**, *9*, 6-21.
- 9 38. Ura, S.; Honda, M.; Yamashita, T.; Ueda, T.; Takatori, H.; Nishino, R.; Sunakozaka,
10 H.; Sakai, Y.; Horimoto, K.; Kaneko, S. Differential microRNA expression between
11 hepatitis B and hepatitis C leading disease progression to hepatocellular carcinoma.
12 *Hepatology* **2009**, *49*, 1098-1112.

1 **Table 1.** Content (mg/100 g dry weight) of oleanolic acid (OA), ursolic acid (UA) and
 2 maslinic acid (MA) in eleven vegetables, carrot, celery, okra, eggplant, spinach, leaf
 3 mustard, gynura, Chinese chive, fennel, basil and daylily. Data are mean \pm SD, n=6.


	OA	UA	MA
carrot	25 \pm 4	31 \pm 6	-
celery	17 \pm 2	-	-
okra	61 \pm 6	23 \pm 3	40 \pm 4
eggplant	53 \pm 4	32 \pm 3	84 \pm 7
spinach	167 \pm 13	133 \pm 8	126 \pm 11
leaf mustard	135 \pm 11	92 \pm 6	174 \pm 12
gynura	- ^a	45 \pm 6	37 \pm 5
Chinese chive	-	-	38 \pm 4
fennel	54 \pm 8	17 \pm 5	-
basil	96 \pm 7	83 \pm 12	35 \pm 4
daylily	118 \pm 10	76 \pm 5	105 \pm 9

4 ^ameans too low to be detected.

5

1 **Figure 1.** HIF-1 α mRNA expression in human Hep3B, Huh7 and HA22T cell lines under
2 normoxia and hypoxia condition. Data are mean \pm SD (n = 9). ^{a,b}Means among bars
3 without a common letter differ, $P < 0.05$.


1



2


3

1 **Figure 2.** Effect of oleanolic acid (OA), ursolic acid (UA) and maslinic acid (MA) at 0
2 (control), 2 or 4 $\mu\text{mol/L}$ upon HIF-1 α mRNA expression in human Hep3B, Huh7 and
3 HA22T cell lines. Data are mean \pm SD (n = 9). ^{a-c}Means among bars without a common
4 letter differ, $P < 0.05$.



1

1 **Figure 3.** Effect of oleanolic acid (OA), ursolic acid (UA) and maslinic acid (MA) at 0
2 (control), 2 or 4 $\mu\text{mol/L}$ upon bFGF mRNA expression in human Hep3B, Huh7 and HA22T
3 cell lines. Data are mean \pm SD (n = 9). ^aMeans among bars without a common letter differ,
4 $P < 0.05$.




□ control ▨ OA, 2 ▩ OA, 4 ▪ UA, 2 ▫ UA, 4 ▬ MA, 2 ■ MA, 4

1


2

1 **Figure 4.** Effect of oleanolic acid (OA), ursolic acid (UA) and maslinic acid (MA) at 0
2 (control), 2 or 4 $\mu\text{mol/L}$ upon VEGF level (upper part) and mRNA expression (lower part) in
3 human Hep3B, Huh7 and HA22T cell lines. Data are mean \pm SD (n = 9). ^{a-d}Means among
4 bars without a common letter differ, $P < 0.05$.




1

2




1

1 **Figure 5.** Effect of oleanolic acid (OA), ursolic acid (UA) and maslinic acid (MA) at 0
2 (control), 2 or 4 $\mu\text{mol/L}$ upon IL-8 level (upper part) and mRNA expression (lower part) in
3 human Hep3B, Huh7 and HA22T cell lines. Data are mean \pm SD (n = 9). ^{a-d}Means among
4 bars without a common letter differ, $P < 0.05$.



1
2
3



1

2

1 **Table 2.** Effect of oleanolic acid (OA), ursolic acid (UA) and maslinic acid (MA) at 0
 2 (control), 2 or 4 $\mu\text{mol/L}$ upon level of GSH, ROS and NO in human normal liver cell line
 3 (L-02 cells), Hep3B, Huh7 and HA22T cell lines. Data are mean \pm SD (n = 9).


	GSH ng/mg protein			
	L-02	Hep3B	Huh7	HA22T
Control	80 \pm 4 ^a	31 \pm 2 ^a	23 \pm 3 ^a	38 \pm 4 ^a
OA, 2	84 \pm 5 ^a	42 \pm 3 ^b	36 \pm 4 ^b	47 \pm 5 ^b
OA, 4	88 \pm 3 ^a	57 \pm 5 ^c	50 \pm 5 ^c	60 \pm 5 ^c
UA, 2	82 \pm 6 ^a	40 \pm 4 ^b	40 \pm 3 ^b	45 \pm 4 ^b
UA, 4	85 \pm 3 ^a	56 \pm 4 ^c	60 \pm 4 ^d	58 \pm 2 ^c
MA, 2	79 \pm 5 ^a	47 \pm 5 ^b	33 \pm 4 ^b	50 \pm 3 ^b
MA, 4	87 \pm 4 ^a	69 \pm 3 ^d	48 \pm 2 ^c	63 \pm 5 ^c
	ROS nmol/mg protein			
	L-02	Hep3B	Huh7	HA22T
Control	0.35 \pm 0.04 ^a	2.06 \pm 0.26 ^d	2.14 \pm 0.13 ^c	1.90 \pm 0.18 ^c
OA, 2	0.33 \pm 0.06 ^a	1.67 \pm 0.20 ^c	1.75 \pm 0.17 ^b	1.58 \pm 0.10 ^b
OA, 4	0.31 \pm 0.02 ^a	1.03 \pm 0.14 ^a	1.39 \pm 0.15 ^a	1.27 \pm 0.12 ^a
UA, 2	0.29 \pm 0.05 ^a	1.72 \pm 0.21 ^c	1.65 \pm 0.09 ^b	1.65 \pm 0.15 ^b
UA, 4	0.32 \pm 0.03 ^a	1.35 \pm 0.16 ^b	1.23 \pm 0.10 ^a	1.19 \pm 0.07 ^a
MA, 2	0.30 \pm 0.06 ^a	1.70 \pm 0.18 ^c	1.72 \pm 0.15 ^b	1.52 \pm 0.14 ^b
MA, 4	0.33 \pm 0.05 ^a	1.38 \pm 0.13 ^b	1.40 \pm 0.11 ^a	1.23 \pm 0.09 ^a
	NO $\mu\text{M/mg}$ protein			
	L-02	Hep3B	Huh7	HA22T

Control	1.3±0.3 ^a	15.7±1.1 ^d	14.6±1.0 ^c	15.3±1.4 ^d
OA, 2	1.6±0.5 ^a	10.5±0.7 ^b	12.1±0.6 ^b	10.8±1.0 ^b
OA, 4	1.4±0.2 ^a	8.3±0.9 ^a	8.5±1.1 ^a	7.9±0.8 ^a
UA, 2	1.2±0.4 ^a	12.6±0.8 ^c	11.8±0.9 ^b	12.9±1.2 ^c
UA, 4	1.5±0.2 ^a	8.0±1.1 ^a	8.3±0.7 ^a	7.6±0.5 ^a
MA, 2	1.4±0.5 ^a	13.1±1.0 ^c	12.2±1.2 ^b	13.0±0.9 ^c
MA, 4	1.3±0.6 ^a	8.2±0.6 ^a	9.0±0.8 ^a	7.4±0.6 ^a


1 ^{a-d}Means in a column without a common letter differ, $P < 0.05$.

1 **Figure 6.** Effect of oleanolic acid (OA), ursolic acid (UA) and maslinic acid (MA) at 0
2 (control), 2 or 4 $\mu\text{mol/L}$ upon invasion and migration in human Hep3B, Huh7 and HA22T
3 cell lines. Data are mean \pm SD (n = 9). ^{a-d}Means among bars without a common letter
4 differ, $P < 0.05$.

invasion




1
2




1 **Figure 7.** Effect of oleanolic acid (OA), ursolic acid (UA) and maslinic acid (MA) at 0
2 (control), 2 or 4 $\mu\text{mol/L}$ upon uPA level (upper part) and mRNA expression (lower part) in
3 human Hep3B, Huh7 and HA22T cell lines. Data are mean \pm SD (n = 9). ^{a-d}Means among
4 bars without a common letter differ, $P < 0.05$.

1



2



1