

# **Oromotor variability in children with mild spastic cerebral palsy: a kinematic study of speech motor control**

Chia-ling Chen,<sup>1,2,\*</sup> MD, PhD; Hsieh-ching Chen<sup>3</sup>, PhD; Wei-hsien Hong,<sup>4</sup> PhD; Fan-pei  
Gloria Yang,<sup>5</sup> Ph.D; Liang-yi Yang,<sup>2</sup> MS; Ching-yi Wu,<sup>6</sup> ScD, OTR

<sup>1</sup>Department of Physical Medicine and Rehabilitation, Chang Gung Memorial hospital, 5  
Fuhsing St. Kweishan, Taoyuan 33302, Taiwan,

<sup>2</sup>Graduate Institute of Early Intervention, Chang Gung University, 259 Wenhwa 1 Rd.,  
Kweishan, Taoyuan 33302, Taiwan

<sup>3</sup>Department of Industrial Engineering and Management, Chaoyang University of Technology,  
168 Jifong E. Rd., Wufong, Taichung County 41349, Taiwan

<sup>4</sup>Department of Sports Medicine, China Medical University, .91 Hsueh-Shih Rd., Taichung,  
40402, Taiwan,

<sup>5</sup>Department of Radiology and Biomedical Imaging, University of California at San Francisco,  
185 Berry Street Suite 350, San Francisco, CA 94107, USA,

<sup>6</sup>Department of Occupational Therapy, Chang Gung University, 259 Wenhwa 1 Rd.,  
Kweishan, Taoyuan 33302, Taiwan

## **\*Corresponding author**

Email addresses:

CLC: clingchen@gmail.com; HCC: hcchen@mail.cyut.edu.tw

WHH: whhong@mail.cmu.edu.tw; FGY: fbyang@gmail.com

LYY: dana9909@gmail.com; CYW: cywu@mail.cgu.edu.tw

**Running head: Oromotor variability in CP**

# ABSTRACT

**Background:** Treating motor speech dysfunction in children with CP requires an understanding of the mechanism underlying speech motor control. However, there is a lack of literature in quantitative measures of motor control, which may potentially characterize the nature of the speech impairments in these children. This study investigated speech motor control in children with cerebral palsy (CP) using kinematic analysis.

**Methods:** We collected 10 children with mild spastic CP, aged 4.8 to 7.5 years, and 10 age-matched children with typical development (TD) from rehabilitation department at a tertiary hospital. All children underwent analysis of percentage of consonants correct (PCC) and kinematic analysis of speech tasks: poly-syllable (PS) and mono-syllable (MS) tasks using the Vicon Motion 370 system integrated with a digital camcorder. Kinematic parameters included spatiotemporal indexes (STIs), and average values and coefficients of variation (CVs) of utterance duration, peak oral opening displacement and velocity. An ANOVA was conducted to determine whether PCC and kinematic data significantly differed between groups.

**Results:** CP group had relatively lower PCCs (80.0-99.0%) than TD group ( $p=0.039$ ). CP group had higher STIs in PS speech tasks, but not in MS tasks, than TD group did ( $p=0.001$ ). The CVs of utterance duration for MS and PS tasks of children with CP were at least three times as large as that of TD children ( $p<0.01$ , Table 4). However, average values of utterance duration, peak oral opening displacement and velocity and CVs of other kinematic data for both tasks did not significantly differ between two groups.

**Conclusion:** High STI values and high variability on utterance durations in children with CP reflect deficits in relative spatial and/or especially temporal control for speech in the CP participants compared to the TD participants. Children with mild spastic CP may have more difficulty in processing increased articulatory demands and resulted in greater oromotor variability than normal children. The kinematic data such as STIs can be used as indices for

detection of speech motor control impairments in children with mild CP and assessment of the effectiveness in the treatment.

**Keywords:** Cerebral palsy, Oromotor variability, Kinematic analysis, Speech motor control, Motion analysis

## Background

Cerebral palsy (CP) refers to a group of developmental disorders in movement and posture, which are attributed to non-progressive disturbances that occurred in the developing fetal or infant brain [1]. Speech and language impairments are commonly associated with CP [2]. These communication disorders, especially poor speech production, may result from disturbed neuromuscular control of speech mechanism [2]. Impaired speech functions such as articulation disorders are present in 38% children with CP [3]. Reduced intelligibility in children with CP can adversely impact communication abilities and limit their vocational, educational, and social participation [4]. Such limitations may consequently diminish these children's quality of life [4].

Children with spastic CP commonly exhibit dysarthria of varying severity. Articulatory imprecision is one of the primary characteristics of dysarthria [5]. Other fairly stable features of CP dysarthria are problems in the consonantal place and manner [6]. Specifically, at the phonemic level, patients with dysarthria displayed anterior lingual place inaccuracy, reduced precision of fricative and affricate manners, and inability to achieve the extreme positions in the vowel articulatory space [6]. Other studies revealed that speakers with CP exhibited smaller vowel working space areas compared to age-matched controls and that the vowel working space area was significantly correlated with vowel and word intelligibility [7].

Quantitative measurements of speech motor control have been used to characterize language and communication deficits in diverse patient populations. These measurements include kinematic [8-11], kinetic [12], electromyographic (EMG) [12-16] and acoustic analyses [17-19]. Kinematic measures of articulatory movements include measurements of movement amplitude, velocity and duration [11], and speech movement trajectory analysis [10, 11]. The spatio-temporal index (STI) values in speech movement trajectory analysis reflect the degree to which repeated performance of a task produces movement trajectories that converge on a single pattern [10]. Therefore, the STI values indicate the degree of

oromotor stability of a speech task that produces movement trajectories [10]. At present, lip and jaw kinematic analyses in previous studies have identified the speech motor control pattern in children with normal development [9, 12, 20, 21]. However, no studies up to date have performed kinematic analysis of speech motor control in children with mild spastic CP.

It is important to conduct speech motor control analysis in children with CP for several reasons. First, quantitative measures of motor control are considerably more sensitive than conventional methods in determining the distribution and nature of orofacial motor impairments which degrade fine motor performance [22]. The speech disorder, which is often associated with athetosis, has been considered with respect to a model of motor learning [23]. Research also reported that the most frequent abnormalities of subjects with athetoid CP included large ranges of jaw movement, inappropriate positioning of the tongue for various phonetic segments, intermittency of velopharyngeal closure caused by an instability of velar elevation, prolonged transition times for articulatory movements, and retrusion of the lower lip [23]. Despite the large number of studies focusing on the description of motor disabilities, there is a lack of literature in quantitative measures of motor control, which may potentially characterize the nature of the speech impairments in children with spastic CP.

Secondly, treating motor speech dysfunction in children with CP requires an understanding of the mechanism underlying speech motor control. Previous research has demonstrated that the measures of dynamics in select structures of the oral motor system were found to be related to impairments in speech intelligibility [22]. Even in mild CP patients with intelligence levels above 70, half of the patients exhibit motor speech problems [2]. However, it remained unclear how the fine articulator movements are controlled and coordinated for speech production in children with mild spastic CP. Understanding the control and coordination mechanism for speech production is essential for developing appropriate treatment.

We hypothesize that speech motor control is impaired even in children with mild spastic

CP because these children have greater oromotor variability than TD children. We predict that these children's oromotor variability can be reflected in high variability on kinematic variables and high STI values in speech tasks. This study aims to investigate speech motor control in children with mild spastic CP using kinematic analysis. We will analyze the kinematic parameters such as utterance duration, peak oral opening displacement, peak oral opening velocity, and spatiotemporal indexes (STIs). We will also characterize kinematic parameters for the sensitive detection of speech motor control problems. The current research will offer valuable kinematic data that support neural-motor models proposed to account for speech motor control problems. More importantly, kinematic data such as STIs can be used as indices for detection of speech motor control impairments in children with mild CP and assessment of the effectiveness in the treatment. This study may potentially provide directions in linguistic and kinematic measures for future studies on patients with speech deficits.

## **METHODS**

### **Participants**

Ten children with mild spastic CP (seven male, three female), aged 4.8 to 7.5 years old (mean age:  $5.9 \pm 1.0$  years), from rehabilitation department at a tertiary hospital, Chang Gung Memorial hospital, were enrolled in the study. The inclusion criteria were as follows: (1) mild spastic CP with Gross Motor Functional Classification System (GMFCS) [24] levels I-II; (2) ability to perform speech tasks with mild articulation disorders; (3) good cooperation during examination; and (4) ability to understand the verbal commands required for analysis.

Exclusion criteria were any history of the following conditions within the previous three months: (1) significant medical problems such as active pneumonia or urinary tract infection; (2) significant hearing impairment; (3) any major surgical treatment such as orthopedic surgery or neurosurgical surgery; (4) any treatment with nerve or motor point block such as a botulinum toxin injection; and (5) history of facial palsy.

The control group consisted of ten age-matched children with typical development (TD) (six male and four female) aged 4.9 to 7.5 years (mean age:  $6.1 \pm 0.8$  years) with no history of learning disabilities, speech impairment, such as specific speech production errors, language impairment, neurological lesions, or visual or hearing impairment. The institutional review board for human studies at Chang Gung Memorial hospital approved the study protocol. All participants and their parents or guardians provided informed consent to participate in the study.

### **Instrumentation**

Kinematic analysis of head and mouth movements during speech tasks was performed using the Vicon Motion 370 system (Oxford metrics Ltd, UK) integrated with a digital camcorder. The Vicon system, which consisted of six infrared cameras, was used in conjunction with a personal computer to capture the movement of reflective markers. Kinematic data for the reflective markers were recorded at a sampling rate of 60Hz and digitally low-pass filtered using a second-order Butterworth filter with 5Hz cut-off frequency. The 5-Hz cut-off frequency was used to reduce markers' velocity error, which might be introduced by noise signal using numerical differentiation method, without significantly altering the results of marker displacements. For each speech task, a digital signal synchronized with an external LED light was collected by the Vicon system to synchronize the video images and to determine onset and offset of marker movement.

### **Assessment Procedures**

We analyzed specific speech production errors, speech intelligibility and performed kinematic analysis of speech tasks on all children. In addition to these analyses, we also analyzed motor severity of children with CP. The same speech pathologist assessed the specific speech production errors of all children with CP and the same physiatrist (CL Chen) classified the motor severity of CP using Gross Motor Functional Classification System (GMFCS) [24]. The GMFCS grades the self-initiated movement of CP patients with particular

emphasis on their functional abilities (sitting, crawling, standing and walking) and their need for assistive devices (e.g., walkers, crutches canes and wheelchairs). The GMFCS employs a 5-point scale (I-V) from “independent” (level I) to “dependent: (level V). Demographic data of all participants, including age and gender were recorded.

### ***Experimental setup for measuring speech intelligibility***

Each child was seated in a quiet room. The recording system used to measure speech intelligibility consisted of an external microphone and a laptop computer (IBM ThinkPad 570E) with 16k Hz sampling rate and 16-bit resolution. The microphone was placed on a table approximately 15 cm from the mouth of the child. The children were shown pictures or texts printed on cards and asked to read them aloud in a normal voice. Whenever the child encountered an unfamiliar word, the examiner explained the word or asked the child to read it with the assistance of phonetic transcription. The examiner did not model the correct sound production or provide other assistance. The speech recording tasks included 69 picture-cards for preschool children and 140 word-cards for school children. Before the speech task started, the examiner told the subjects that the words they read were going to be recorded. During the task, the examiner recorded a speech sample of each subject.

The percentage of consonants correct (PCC), modified from procedures outlined by Shriberg and Kwiatkowski (1982), was used to determine severity of speech intelligibility [25]. The PCC information was used as an index to quantify severity of involvement [25]. To measure PCC, a rater must make correct-incorrect judgments of individual sounds produced in the speech sample of each subject. The same rater, who was a native Mandarin speaker with normal hearing, transcribed recorded speech samples. The PCC was calculated as  $100 \times (\text{number of correct consonants} / \text{number of correct plus incorrect consonants})$  [25]. In order to test intra-rater and inter-rater reliabilities, a research assistant was recruited to rate the sound of 10 children randomly selected from the data base. The intra-class correlation coefficient (ICC) values of inter-rater and intra-rater reliability for PCC were 0.812 and 0.977,



respectively.

Additionally, the same speech pathologist identified all subjects' specific speech production errors based on the place-manner-voicing analysis with phonological process analysis: omission, substitution, and distortion [26] from the recorded speech samples. The patterns of phonological process analysis consisted of assimilation, fronting, backing, stopping, voicing, de-voicing, affrication, de-affrication, nasalization, de-nasalization, and lateralization [26].

### ***Experimental setup of Kinematic analysis***

During the Kinematic analysis task, the subjects were comfortably seated in chairs adjusted to 100% of lower leg length, measured from the lateral knee joint to the floor with the subject standing. The trunk was secured to the chair-back with a harness in order to minimize trunk flexion and rotation. Each subject wore the same facial masks (Figure 1A). Four reflective markers in diameter of 0.6 cm were attached to the facial mask at the forehead, bilateral pre-auricular areas and nose to establish a reference coordination system with the positive  $x$ ,  $y$ ,  $z$  orientation line in horizontal rightward, anterior-posterior, and vertical upward directions respectively (Fig. 1B). The direction of  $x$ -axis is defined along the line joining the bilateral markers at per-auricular areas. The  $y$ -axis is perpendicular to the frontal plane passing through markers at the forehead and bilateral pre-auricular areas. The  $z$ -axis is orthogonal to  $x$ - and  $y$ -axis. The origin of reference coordination system was located at the nose marker. For oral-movement tracking, five markers were attached to the bilateral mouth corners, the upper and lower lips at midline and jaw region (Fig. 1A). That is, nine reflective markers were used on the facial mask and oral areas (Fig. 1C).

All participants underwent mono-syllable (MS) and poly-syllable (PS) task assessments. The speech task comprised combination of two bilabial consonants ( $/p/$ ,  $/p^h/$ ) and five basic vowels ( $/a/$ ,  $/i/$ ,  $/u/$ ,  $/æ/$ , and  $/o/$ ) to form the target consonant-vowel syllable. These vowels are selected because they are the most common in human languages [27]. We chose the bilabial

consonants to elicit the lip opening-closing movement in each consonants-vowel syllable. For both tasks, the examiner pronounced the syllables themselves and asked participants to repeat after the examiner. The examiner said the target syllable(s) at a relatively slow rate for clarity purpose. During the MS tasks, participants were asked to speak /pa/, /pi/, /p<sup>h</sup>u/, /p<sup>h</sup>æ/, and /p<sup>h</sup>o/ separately. During the PS task, participants were required to speak /pa, pi, p<sup>h</sup>u, p<sup>h</sup>æ, p<sup>h</sup>o/ in a sequence.

The order of task presentation was randomized. Each task was repeated at least 10 times until we collected ten usable trials for each task in each individual. If markers' kinematic data were not correctly captured, these trials were excluded and retested. We used ten trials in each task for analysis. All participants were allowed a 5-sec rest period between each trial repetition and a 15-sec rest period between each task. All participants were allowed three practice trials to familiarize themselves with the experimental setup. A vocal cue together with an LED-light signal was provided to indicate the start of the task by the examiner.

### **Data analysis**

An analysis program for kinematic data coded by LabView (National Instruments, USA) was developed to process the kinematic data. Only the kinematic data of vertical movement (oral aperture in the z-axis) of lip markers were analyzed in this study. The utterance duration, peak oral opening displacement, peak oral opening velocity and STIs of each task were analyzed while performing speech tasks. The overall utterance period of a speech task was determined from the instance of peak closing velocity right before the initial opening of the lower lip to the instance when the lower lip was at the peak velocity of its closing movement during the final syllable (Fig. 2). The acoustic traces were used to verify the kinematically-derived onsets. For each task, lower lip displacement waveforms during individual utterance periods were used for STI analysis (Fig. 2). Each lower lip displacement waveform was first amplitude normalized by subtracting the individual mean and dividing by the standard deviation and then time normalized to 100% duration. For time normalization,

101 data points were resampled from each amplitude-normalized waveform by a linear interpolation scheme. One standard deviation was then computed every 2% normalized duration across 10 waveforms of each task. There were 50 (from 2% to 100%) standard deviations computed. These standard deviations were then summed to determine the overall STI [10]. In addition to computing STI, for each PS task, peak opening of mouth (oral aperture) was identified by the maximum vertical distance between the upper- and lower-lip markers within the entire utterance duration. Peak oral opening velocity was calculated by determining the maximum time derivatives of the vertical oral opening displacement. The mean peak oral opening velocity and displacement in a PS or a MS task were determined by averaging the maximum opening velocities and displacements, respectively, of each repeated trial.

Furthermore, the coefficient of variation (CV) for kinematic data (utterance duration, peak opening displacement, and peak opening velocity) obtained by dividing the standard deviation of kinematic data by the mean kinematic data. Larger CV indicates higher variability of kinematic data in speech tasks.

### **Statistical Analysis**

Group differences in age were compared by an independent t test. Gender differences between groups were determined by Fisher's exact test. An ANOVA was conducted to determine whether PCC and kinematic data (values and CVs of utterance duration, peak opening displacement, and peak opening velocity, and STI) significantly differed between groups. The effect size  $d$  was calculated for each PCC and kinematic data to index the magnitude of the difference in PCC and kinematic data varied between groups [28]. According to Cohen [29], an  $d$  of at least 0.50 represents a large effect; an  $d$  of at least 0.30 represents a moderate effect, and an  $d$  of at least 0.10 represents a small effect. Given that multiple comparisons are made on the speech productions of two groups of speakers, a  $p$  value of  $<0.01$  was considered statistically significant.

## RESULTS

Demographic data did not significantly differ between children with CP and children with TD. Four children with CP were at GMFCS level I, and six children with CP were at level II. Five children had specific speech production errors: stopping and voicing (2 cases), distortion (one case), backing (one case), and fronting and de-affrication (one case).

The PCC ranged from 80.0-99.0% in children with CP, and 95.5-100.0% in TD children. The ANOVA analysis showed that the CP group had relatively lower PCC scores than TD group with moderate effect, though the difference did not achieve significance ( $F_{1,18} = 4.962$ , effect size  $d = 0.465$ ,  $p = 0.039$ , Table 1).

STI for PS tasks between the CP and TD groups were significantly different ( $F_{1,18} = 14.093$ , effect size  $d = 0.663$ ,  $p = 0.001$ , Table 1). However, there were no significant differences in STI of MS tasks between the CP and TD groups (Table 1). The average STI values for PS tasks were greater in CP children than TD children (Table 1). The average STI values of children with mild CP were 19.5 in MS tasks and 30.1 in PS tasks (Table 1). Figure 3 illustrates the original waveforms, normalized waveforms and STIs in PS tasks of one child with CP and one child with TD.

The ANOVA analysis showed no significant differences in the utterance durations, peak oral opening displacement and velocity of both MS and PS tasks between the CP and TD groups (Table 2). The average utterance durations of children with mild CP were 0.95 sec/syllable in both MS and PS tasks (Table 2). The average peak oral opening displacements of children with mild CP were 1.17 cm in MS tasks and 1.84 cm in PS tasks (Table 2). The average peak oral opening velocities of children with mild CP in MS and PS tasks were 42.4 and 73.5 cm/sec, respectively (Table 2).

The CVs of utterance duration for MS and PS tasks between groups were different ( $p \leq 0.01$ , Table 3). The CVs of utterance duration for MS and PS tasks of children with CP were

at least three times as large as that of TD children ( $p \leq 0.01$ , Table 3). However, the CVs of peak oral opening displacement and velocities for MS and PS tasks did not differ between groups (Table 3).

## DISCUSSION

The present study is the first kinematic study of speech motor control in children with CP. The lack of this type of research on CP children may be due to the technical difficulty of managing movement artifacts due to head or trunk control problems in these children. To overcome this problem, we used a specially designed facial mask. More importantly, the use of multiple measures in the current research offered an alternative to understanding the underlying motor control for speech production as the different measures used here supplement each other and measured different aspects of oromotor movement and speech production. This approach enables kinematic analysis of speech motor control in children with CP and is likely to provide a corroborated account of the articulatory behaviors in this population.

Our study revealed that children with mild spastic CP had greater STIs in PS tasks than children with TD. In order to interpret this result, we need to understand the motor control system at the neural level, which is described in Smith [13]. To produce intelligible speech, the brain must generate motor commands to control activation of many different motor neuron pools that innervate the muscles for speech production [13]. Each coordinated movement requires temporal control and spatial control in the innervating muscles of the articulators, the larynx, and the chest wall [13]. Using Smith's model, high STI values in children with CP might reflect deficits in relative temporal and/or spatial control for speech, which might be caused by damage to the nervous system during development. Normally maturation of the neural systems underlying language processing and speech production follow a course of cortical, dendritic and synaptic development [30-32]. Damage to the immature brain in children with CP may cause variations in neural drive to muscles during speech production. As the developing system explores different solutions to achieving vocal tract goals, higher speech variability is produced [15]. This is supported by our finding that children with mild spastic CP had greater oromotor variability and relatively lower speech

intelligibility in speech production than children with TD.

We also found that children with mild CP had higher STIs and relatively lower PCC in speech tasks than TD children, though the PCC did not achieve significant differences. CP group's poor performance in STI and PCC can also be interpreted at the level of oromotor control, such as spasticity, weakness, and voluntary control abnormalities. Several theories have been proposed concerning the pathophysiology of dysarthria in subjects with CP [33], such as weakness of speech muscles [34], abnormalities of muscle tone due either to spasticity of speech muscles [34], primitive reflexes or pathological reactions interfering the articulatory control [35], or an imbalance of positive and negative oral reactions [36]. Children with spastic CP are quantitatively less consistent in their movement output compared to TD children. STI values are related to observed differences in severity of dysarthria [37]. Thus, children with spastic CP produce relatively lower speech intelligibility in speech tasks compared with TD children.

Notably, we observed significant between-group differences in STI values for PS, but not for MS, utterances. This indicates that the STI difference between CP and TD children becomes more distinctive as task complexity increases. Previous research also reported that children with mild spastic CP have more difficulty than normal children in processing increased articulatory demands, which is reflected in greater oromotor variability [30, 38]. Speech motor planning, execution, or both may be affected by processes remote from the motor output stage [30]. These studies have proposed that utterance length and complexity on speech motor performance are related to the effects of increased processing demands on articulatory movement stability [30]. These observations may explain why only PS tasks display group difference in the present study, as the processing demands on articulatory movement stability in the PS tasks exceeded those for MS tasks. Other clinical studies also revealed that syntactic complexity affects the speech motor stability of fluent speech in adults who stutter [38]. Their results suggest that language formulation processes may affect speech

production processes. Along with previous findings, our results suggest that the treatment strategies of high-demanding articulatory training, such as complicated speech tasks, may be especially important for children with mild CP.

It appears that the pattern of variability between mild spastic CP and TD groups is different. Children with CP had greater CVs of utterance duration for MS and PS tasks at least three times as large as those of TD children, although the CVs for other kinematic measures did not differ significantly between the two groups. Higher STI values and variability on utterance durations in children with CP in comparison with TD children might reflect deficits in relative spatial and/or especially temporal control (high variability on utterance durations) for speech. The temporal control indicates the appropriate timing control of muscle activations and de-activations and spatial control indicates the appropriate graded muscle activity control for speech production. The deficits in spatial and temporal control may arise from poor motor coordination. These findings may imply children with mild spastic CP may employ a different organizational unit or a different planning strategy in performing speech tasks from TD children.

The STI differences identified in the present study is of great value because the STIs can be used as an index for detection of speech motor problems in children with mild CP. In addition, STI index may also be used for assessing the effectiveness of treatment of such problems. Kleinow et al. reported reductions in STI in response to a speech treatment for adults with hypokinetic dysarthria associated with Parkinsonism [39]. We may potentially apply this approach to the assessment of intervention designed for children with mild CP.

The findings of this study may be limited due to its design in the aspects of sample size, measurement methods, and subject characteristics. The actual values of kinematic variables including STI, utterance durations, peak displacement and peak velocities could be influenced by multiple factors, such as instrumentation, specific tasks and signal processing. For example, the utterance durations are relatively long because the participants repeat the target syllable(s)



at a relatively slow rate for clarity purpose. The STI measures varies as a function of the speech task used, and therefore it is challenging to interpret STI differences or similarities in different tasks such as MS and PS tasks. The MS task simply requires syllable repetition whereas the PS task demands distinct phonetic composition. In prior studies, STI is typically used with actual speech utterances, while the present study uses syllable repetitions as the speech motor task for open-closing oral movement in each utterance. The utterance duration was described as a kinematic event in this study. However, it is likely that a kinematic event occurred before the actual utterance by use of other articulators, which preceded kinematic detection of the lips and jaw. Besides, the tasks are relatively simple and might not sufficiently tax the motor systems of children who have minimal speech impairment. We only enrolled children with mild spastic CP and mild speech intelligibility impairment in the study. Therefore, our results can not be generalized to all cases of CP. Despite this limitation, this study has demonstrated some heuristic value relative to the dynamic organization motor speech in children with mild forms of CP.

## **CONCLUSION**

Children with mild spastic CP showed greater STIs in PS tasks, but not in MS tasks, than children with TD do. These findings suggest that children with CP have more difficulty in processing increased articulatory demands. However, the average values of utterance duration, peak oral opening displacement and peak oral opening velocity of both speech tasks do not significantly differ between children with CP and children with TD. High STI values and high variability on utterance durations in children with CP reflect deficits in relative spatial and/or especially temporal control for speech in the CP participants compared to the TD participants. The STIs can be used as an index for sensitive detection and assessing the effectiveness in the treatment of speech motor control problems in children with mild CP.

The kinematic data for speech motor control provided in this study may help clinicians to understand the speech motor control and planning treatment strategies for children with CP.

Future studies may focus on speech tasks using a variety of linguistic structures to elicit different muscle contractions and movements to provide better diagnosis and treatment for CP children at different degrees of severities and to examine the effectiveness of different treatment strategies in these children.

## **Competing interests**

The authors declare that we have no competing interests.

## **Authors' contributions**

CLC participated in the conception, study design, analysis, and draft of this manuscript. HCC participated in the experimental setup of kinematic analysis, kinematic data collection and analysis, and revising of this manuscript. WHH carried out the kinematic data collection and analysis. FGY participated in the data interpretation and the revising of this manuscript. LYY carried out the data collection and analysis. CYW carried out the data collection and interpretation. All authors read and approved the final manuscript.

## **Acknowledgments**

The authors would like to thank the National Science Council, Taiwan for financially supporting this research under Contract No. NSC 92-2314-B-182A-050.

## References:

1. Rosenbaum P, Paneth N, Leviton A, Goldstein M, Bax M, Damiano D, Dan B, Jacobsson B: **A report: the definition and classification of cerebral palsy April 2006.** *Dev Med Child Neurol Suppl* 2007, **109**:8-14.
2. Pirila S, van der Meere J, Pentikainen T, Ruusu-Niemi P, Korpela R, Kilpinen J, Nieminen P: **Language and motor speech skills in children with cerebral palsy.** *J Commun Disord* 2007, **40**:116-128.
3. Sankar C, Mundkur N: **Cerebral palsy-definition, classification, etiology and early diagnosis.** *Indian J Pediatr* 2005, **72**:865-868.
4. Hustad KC: **The relationship between listener comprehension and intelligibility scores for speakers with dysarthria.** *J Speech Lang Hear Res* 2008, **51**:562-573.
5. Marchant J, McAuliffe MJ, Huckabee ML: **Treatment of articulatory impairment in a child with spastic dysarthria associated with cerebral palsy.** *Dev Neurorehabil* 2008, **11**:81-90.
6. Platt LJ, Andrews G, Young M, Quinn PT: **Dysarthria of adult cerebral palsy: I. Intelligibility and articulatory impairment.** *J Speech Hear Res* 1980, **23**:28-40.
7. Liu HM, Tsao FM, Kuhl PK: **The effect of reduced vowel working space on speech intelligibility in Mandarin-speaking young adults with cerebral palsy.** *J Acoust Soc Am* 2005, **117**:3879-3889.
8. Green JR, Moore CA, Higashikawa M, Steeve RW: **The physiologic development of speech motor control: lip and jaw coordination.** *J Speech Lang Hear Res* 2000, **43**:239-255.
9. Green JR, Moore CA, Reilly KJ: **The sequential development of jaw and lip control for speech.** *J Speech Lang Hear Res* 2002, **45**:66-79.
10. Smith A, Goffman L, Zelaznik HN, Ying G, McGillem C: **Spatiotemporal stability and patterning of speech movement sequences.** *Exp Brain Res* 1995, **104**:493-501.

11. Smith A, Johnson M, McGillem C, Goffman L: **On the assessment of stability and patterning of speech movements.** *J Speech Lang Hear Res* 2000, **43**:277-286.
12. Smith A: **The control of orofacial movements in speech.** *Crit Rev Oral Biol Med* 1992, **3**:233-267.
13. Smith A: **Speech motor development: Integrating muscles, movements, and linguistic units.** *J Commun Disord* 2006, **39**:331-349.
14. Green JR, Moore CA, Ruark JL, Rodda PR, Morvee WT, VanWitzenburg MJ: **Development of chewing in children from 12 to 48 months: longitudinal study of EMG patterns.** *J Neurophysiol* 1997, **77**:2704-2716.
15. Wohlert AB, Smith A: **Developmental change in variability of lip muscle activity during speech.** *J Speech Lang Hear Res* 2002, **45**:1077-1087.
16. Folkins JW, Zimmermann GN: **Jaw-muscle activity during speech with the mandible fixed.** *J Acoust Soc Am* 1981, **69**:1441-1445.
17. Ansel BM, Kent RD: **Acoustic-phonetic contrasts and intelligibility in the dysarthria associated with mixed cerebral palsy.** *J Speech Hear Res* 1992, **35**:296-308.
18. Goffman L: **Prosodic influences on speech production in children with specific language impairment and speech deficits: kinematic, acoustic, and transcription evidence.** *J Speech Lang Hear Res* 1999, **42**:1499-1517.
19. Vorperian HK, Kent RD: **Vowel acoustic space development in children: a synthesis of acoustic and anatomic data.** *J Speech Lang Hear Res* 2007, **50**:1510-1545.
20. Smith A, Goffman L: **Stability and patterning of speech movement sequences in children and adults.** *J Speech Lang Hear Res* 1998, **41**:18-30.
21. Smith BL, Kenney MK, Hussain S: **A longitudinal investigation of duration and temporal variability in children's speech production.** *J Acoust Soc Am* 1996,

- 99:2344-2349.
22. Barlow SM, Abbs JH: **Fine force and position control of select orofacial structures in the upper motor neuron syndrome.** *Exp Neurol* 1986, **94**:699-713.
  23. Kent R, Netsell R: **Articulatory abnormalities in athetoid cerebral palsy.** *J Speech Hear Disord* 1978, **43**:353-373.
  24. Palisano R, Rosenbaum P, Walter S, Russell D, Wood E, Galuppi B: **Development and reliability of a system to classify gross motor function in children with cerebral palsy.** *Dev Med Child Neurol* 1997, **39**:214-223.
  25. Shriberg LD, Kwiatkowski J: **Phonological disorders III: a procedure for assessing severity of involvement.** *J Speech Hear Disord* 1982, **47**:256-270.
  26. Bankson NW, Bernthal JE: **Phonological assessment procedures.** In *Articulation and Phonological Disorders* 5<sup>th</sup> edition. Edited by Bernthal JE, Bankson NW. USA, Boston: Pearson Education, Inc; 2004: 233-241
  27. Ladefoged P, Maddieson I: *The sounds of the world's languages.* Malden, MA: Blackwell; 1996.
  28. Rosenthal R, Rosnow RL: *Essentials of behavioral research: methods and data analysis.* 2nd edn. New York: McGraw-Hill; 1991.
  29. Cohen J: *Statistical power analysis for the behavioral sciences.* 2nd edn. Hillsdale: Lawrence Erlbaum Associates; 1988.
  30. Maner KJ, Smith A, Grayson L: **Influences of utterance length and complexity on speech motor performance in children and adults.** *J Speech Lang Hear Res* 2000, **43**:560-573.
  31. Huttenlocher PR: **Morphometric study of human cerebral cortex development.** *Neuropsychologia* 1990, **28**:517-527.
  32. Paus T, Zijdenbos A, Worsley K, Collins DL, Blumenthal J, Giedd JN, Rapoport JL, Evans AC: **Structural maturation of neural pathways in children and adolescents:**

- in vivo study.** *Science* 1999, **283**:1908-1911.
33. O'Dwyer NJ, Neilson PD: **Voluntary muscle control in normal and athetoid dysarthric speakers.** *Brain* 1988, **111 ( Pt 4)**:877-899.
  34. Ingram TT, Barn J: **A description and classification of common speech disorders associated with cerebral palsy.** *Cereb Palsy Bull* 1961, **3**:57-69.
  35. Mysak ED: **Dysarthria and Oropharyngeal Reflexology: A Review.** *J Speech Hear Disord* 1963, **28**:252-260.
  36. Clement M, Twitchell TE: **Dysarthria in cerebral palsy.** *J Speech Hear Disord* 1959, **24**:118-122.
  37. McHenry MA: **The effect of pacing strategies on the variability of speech movement sequences in dysarthria.** *J Speech Lang Hear Res* 2003, **46**:702-710.
  38. Kleinow J, Smith A: **Influences of length and syntactic complexity on the speech motor stability of the fluent speech of adults who stutter.** *J Speech Lang Hear Res* 2000, **43**:548-559.
  39. Kleinow J, Smith A, Ramig LO: **Speech motor stability in IPD: effects of rate and loudness manipulations.** *J Speech Lang Hear Res* 2001, **44**:1041-1051.

## Figures

Figure 1 - Experimental setup for kinematic analysis of speech tasks. The reflective markers were attached to the facial mask and oral areas and reference coordination system was established by marks on the mask.

Figure 2 - Illustration indicates the data used in analyses for poly-syllable speech task. Left vertical line is identified as the instance of peak closing velocity right before the initial opening of the lower lip marker. Right vertical line is defined as the instance when the lower lip was at the peak velocity of its closing movement during the final syllable of the lower lip marker. Both vertical lines mark the displacement period for spatiotemporal index (STI) analysis and the time interval between two points used to measure overall utterance duration in the poly-syllable speech task.

Figure 3 - Illustration indicates the original waveforms, normalized wave forms and spatiotemporal indexes (STIs). Comparison of original waveforms, normalized wave forms and spatiotemporal indexes (STIs) between a child with cerebral palsy and a child with typical development.



Table 1 - Speech intelligibility and spatiotemporal index in children with cerebral palsy and typical development.

Data	Children groups		ANOVA		
	Spastic CP (n =10)	TD (n =10)	$F_{1,18}$	$p$ value	Effect size $d$
Speech intelligibility					
Percentage of consonants correct (PCC)	92.6±7.2	97.7±1.5	4.962	0.039	0.465
Spatiotemporal index (STI)					
Mono-syllable task	19.5±5.1	16.5±4.3	1.937	0.181	0.312
Poly-syllable task	30.1±6.9	21.5±2.3	14.093	0.001*	0.663

\*  $p < 0.01$ .

Values are expressed as mean ± SD.

CP: cerebral palsy; TD: typical development.

PCC: percentage of consonants correct.

Table 2 - Average values of kinematic data in children with cerebral palsy and typical development.

Kinematic parameters	Children groups		ANOVA		
	Spastic CP (n =10)	TD (n =10)	$F_{1,18}$	$p$ value	Effect size $d$
Utterance duration (sec/syllable)					
Mono-syllable task	0.95±0.21	1.01±0.06	0.858	0.367	0.213
Poly-syllable task	0.95±0.30	0.97±0.09	0.049	0.828	0.052
Peak vertical oral opening displacement (cm)					
Mono-syllable task	1.17±0.34	1.23±0.55	0.096	0.760	0.073
Poly-syllable task	1.84±0.46	1.83±0.68	0.001	0.970	0.009
Peak vertical oral opening velocity (mm/sec)					
Mono-syllable task	42.4±10.0	42.5±12.5	0.001	0.980	0.006
Poly-syllable task	73.5±17.5	71.4±16.9	0.079	0.782	0.066

Values are expressed as mean ±SD.

CP: cerebral palsy; TD: typical development.

Table 3 - Coefficients of variation (CVs) of kinematic data in children with cerebral palsy and typical development.

Coefficients of variation (CVs)	Children groups		ANOVA		
	Spastic CP (n =10)	TD (n =10)	$F_{1,18}$	$p$ value	Effect size $d$
Utterance duration					
Mono-syllable task	17.87±11.87	5.23±2.33	10.923	0.004*	0.615
Poly-syllable task	24.30±17.90	7.57±4.11	8.299	0.010	0.562
Peak vertical oral opening displacement					
Mono-syllable task	24.34±13.53	34.14±26.16	1.108	0.307	0.241
Poly-syllable task	18.30±15.82	29.42±20.20	1.880	0.187	0.308
Peak vertical oral opening velocity					
Mono-syllable task	17.21±14.97	24.69±13.50	1.376	0.256	0.266
Poly-syllable task	15.63±17.09	15.16±18.78	0.003	0.954	0.014

\*  $p < 0.01$ .

Values are expressed as mean  $\pm$  SD.

CP: cerebral palsy; TD: typical development.




Fig. 1. Experimental setup for kinematic analysis of speech tasks.




Fig. 2. Illustration indicates the data used in analyses for poly-syllable speech task.




Fig 3. Illustration indicates the original waveforms, normalized wave forms and spatiotemporal indexes (STIs).