

FORUM REVIEW ARTICLE

Special Topic Issue: Novel Risk Factors for Cardiovascular Diseases and Its Therapy

Guest Editor: Prof. Nilanjana Maulik, ARS

A Central Role of Heme Oxygenase-1 in Cardiovascular Protection

MENG-LING WU,^{1,2} YEN-CHUN HO,^{1,3} and SHAW-FANG YET^{1,4}

¹Institute of Cellular and System Medicine, National Health Research Institutes, Zhunan, Taiwan.

²College of Life Sciences, National Tsing Hua University, Hsinchu, Taiwan.

³Graduate Institute of Life Sciences, National Defense Medical Center, Taipei, Taiwan.

⁴Graduate Institute of Basic Medical Science, China Medical University, Taichung, Taiwan.

Running head: HO-1 in cardiovascular protection

Address correspondence to:

Shaw-Fang Yet, Ph.D.

Institute of Cellular and System Medicine

National Health Research Institutes

35 Keyan Road, Zhunan, Miaoli County 35053, Taiwan

E-mail: syet@nhri.org.tw

Abstract

The intrinsic defense mechanisms of the body are critical in protecting tissues from injury in response to pathological stress. Heme oxygenase-1 (HO-1), a stress response protein, is induced in response to various pathological stimuli to serve a cytoprotective function. By degrading the oxidant heme and generating the antioxidant bilirubin and anti-inflammatory molecule carbon monoxide, HO-1 may protect cell from injury due to oxidative and pathological stress. Oxidative stress in the heart caused by ischemia and reperfusion leads to cardiomyocyte death and subsequent myocardial infarction. Vascular diseases including atherosclerosis, graft failure, and restenosis are all associated with reactive oxygen species-induced injury and inflammation. Given that cardiovascular disease is the leading cause of death worldwide, there is considerable interest in developing new strategies for preventing and treating cardiovascular disease. Because HO-1 is induced in the heart and blood vessels in response to various stresses a role of HO-1 has been implicated in cardiovascular homeostasis. Numerous studies using pharmacological method or genetic approach have since demonstrated the cardiovascular protective function of HO-1. Importantly, a number of studies have associated human HO-1 gene promoter polymorphisms with risk for vascular diseases. Taken together, HO-1 has a great therapeutic potential for cardiovascular disease.

Introduction

In response to pathophysiological stress, the body elicits an endogenous adaptive mechanism to protect cells and tissues from injury. Heme oxygenase-1 (HO-1) upregulation is one such intrinsic defense system. HO-1 is a member of the heme oxygenase family of proteins, which include 3 isoforms, HO-1, HO-2, and HO-3. HO-1, also known as heat shock protein 32, is the inducible isoform, HO-2 is constitutively expressed, whereas HO-3 has lower enzymatic activity and is less well characterized. Heme oxygenase degrades the prooxidant heme to carbon monoxide (CO), biliverdin (subsequently reduced to bilirubin), and ferrous iron (56). By degrading the prooxidant heme and generates antioxidant bilirubin, HO-1 may protect cells against oxidative stress. CO is a gas molecule that increases intracellular levels of cGMP, which regulates vascular tone and smooth muscle cell (SMC) development. CO has recently been shown to have anti-proliferative and anti-inflammatory properties. Ferrous iron can induce ferritin expression for iron sequestration. By generating these biologically active molecules and its upregulation in response to stress conditions, HO-1 may thus protect cells and tissues from oxidative stress-induced injury.

Cardiovascular disease is a leading cause of death worldwide. Vascular diseases including atherosclerosis, graft failure, and restenosis are all associated with reactive oxygen species-induced injury. In addition, oxidative stress in the heart caused by ischemia and reperfusion leads to cardiomyocyte death and subsequent myocardial infarction. Therefore, increased expression of antioxidant enzymes and stress proteins might protect against the development of cardiovascular diseases. Because HO-1 is induced in the heart and blood vessels in response to various stresses a role of HO-1

has been implicated in cardiovascular homeostasis. Numerous studies using pharmacological method or genetic approach have since demonstrated the cardiovascular protective function of HO-1. A number of studies have associated human HO-1 gene promoter polymorphisms with risk for vascular diseases (11,22,81,83). Importantly, the clinical significance of HO-1 in human cardiovascular disease has been highlighted in a child with HO-1 deficiency (45,92).

HO-1 in the development of atherosclerosis

Atherosclerosis is the leading cause of mortality and morbidity in the United States and many other countries. When low density lipoprotein (LDL) particles in individuals with hypercholesterolemia become trapped in an artery, they undergo oxidation and are then internalized by macrophages, resulting in foam cell formation (75). Removal and sequestration of oxidized LDL (oxLDL) are the initial, protective role of macrophages in the inflammatory response and minimize the damaging effects of modified LDL on endothelial and smooth muscle cells (SMCs). However, the inflammatory response may be further amplified and atherosclerotic lesions develop by accumulating macrophage-derived foam cells in the intima (32,75). Subsequent migration and proliferation of SMCs from media into intima lead to formation of advanced complicated lesions (76). Recognizing the important contribution of inflammation in atherogenesis, Ross has proposed that atherosclerosis is an inflammatory disease (76).

The detection of HO-1 expression in endothelium, foam cells, and macrophages of advanced lesions from both humans and experimental apolipoprotein E-deficient

(apoE^{-/-}) mice indicates a role of HO-1 in atherogenesis (89). Supporting a function of HO-1 in atherosclerosis, induction of HO-1 inhibits the monocyte transmigration induced by mildly oxidized LDL (38). Pharmacological inhibition of HO-1 activity by Sn-protoporphyrin IX (SnPP) in Watanabe heritable hyperlipidemic rabbits significantly increases atherosclerotic lesions, suggesting HO-1 has anti-atherogenic properties in vivo (39). Similarly, HO-1 inhibition by SnPP in LDL-receptor knockout mice fed high-fat diets results in increased lesion size (40). Evaluation of plasma lipid levels shows that HO-1 inhibition increases plasma and tissue lipid peroxide levels without affecting plasma lipid composition, suggesting the anti-atherogenic properties of HO-1 might be conducted through the prevention of lipid peroxidation (39,40).

Since the effect of metalloporphyrins might have nonselective effects on HO-1 (31), to demonstrate unequivocally the role of HO-1 in atherosclerosis, we used a loss of function approach by first generating mice deficient in both HO-1 and apoE (HO-1^{-/-}apoE^{-/-}) and then subjected mice to Western diet (93). In response to hypercholesterolemia, although total plasma cholesterol levels are similar between HO-1^{-/-}apoE^{-/-} and apoE^{-/-} mice, HO-1^{-/-}apoE^{-/-} mice develop larger and more advanced lesions than mice deficient in apoE alone (93) (Fig. 1A and B). Furthermore, Sudan IV staining revealed greater lipid accumulation in mice deficient in both HO-1 and apoE (93) (Fig. 1C and D). Our genetic study clearly demonstrates that an absence of HO-1 exacerbates atherosclerotic lesion formation (93).

Orozco et al explored whether HO-1 expression in macrophages contributes to the anti-atherogenic effect of HO-1 (66). Using peritoneal macrophages from wild type, HO-1 heterozygous, or HO-1^{-/-} mice they found that HO-1 expression in macrophages

reduces inflammatory responses against oxLDL (66). Furthermore, HO-1^{-/-} macrophages exhibit increased foam cell formation. Bone marrow transplantation experiments reveal that lack of HO-1 in the reconstituted bone marrow results in an increase in the inflammatory component of atherosclerotic lesions in LDL-receptor knockout mice. These results indicate that HO-1 expression in macrophages contributes to anti-atherogenic effect by decreasing the inflammatory component of atherosclerotic lesions (66).

Given that lack of HO-1 exacerbates atherosclerosis, the next question is whether overexpression of HO-1 would attenuate atherosclerotic lesion formation. HO-1 induction by chemical inducers (hemin or hemin and desferrioxamine) significantly attenuates atherosclerotic lesions in LDL-receptor knockout mice fed high-fat diets (40). Similarly, adenovirus-mediated HO-1 gene transfer attenuates the development of atherosclerosis in apoE^{-/-} mice (42). Furthermore, the iron deposition as well as tissue iron content was much less in aortic tissue of mice treated with adenovirus expressing HO-1, indicating overexpression of HO-1 in vascular cells facilitates iron metabolism and attenuates development of atherosclerosis in apoE^{-/-} mice (42). These gain of function experiments demonstrate a critical protective role of HO-1 in atherosclerosis.

HO-1 and plaque stability

As the atherosclerotic lesions progress, the lesions may become unstable and vulnerable with thinning of the fibrous cap and a large necrotic core with lipid accumulation, leading to plaque rupture, acute thrombus formation, and subsequent myocardial infarction. Accumulating evidence indicates that matrix metalloproteinases

(MMPs) and their inhibitors tissue inhibitor of metalloproteinase (TIMP) are important mediators in intimal thickening and atherosclerotic plaque rupture (19,63,64). Early studies show that increased expression and matrix degrading activity of MMPs in vulnerable regions of human atherosclerotic plaques might contribute to plaque rupture (27,35). On the other hand, expression of TIMPs, particularly TIMP-3, might counteract MMP activity in atheroma and increase plaque stability (23). Further, analysis of human carotid endarterectomy tissues revealed a differential distribution of MMPs and TIMPs over atherosclerotic plaques (15).

Although it is well accepted that MMPs are critically involved in atherosclerotic plaque instability; however, the molecular mechanisms regulating plaque stability remain unclear. Recently, Cheng et al (13) explored the role of HO-1 in the progression of atherosclerotic lesions from stable toward vulnerable plaque. Analysis of atherectomy biopsy from patients with clinical carotid artery disease revealed that HO-1 expression levels correlate closely with features of vulnerable human atheromatous plaque, including increased thrombogenicity and increased expression levels of MMP-9, but inversely correlate with intraplaque vascular SMCs and collagen deposition (13). The increased MMP-9 expression is consistent with previous findings that MMP-9 is very abundant in segments of human carotid endarterectomy tissues with intraplaque hemorrhage (15). The correlation of HO-1 and MMP-9 levels suggested a role of HO-1 in regulating MMP-9 expression. This idea is corroborated by recent findings that HO-1 inhibits human breast carcinoma cell invasion and migration through suppressing MMP-9 expression (49). Further supporting this notion, transplantation of HO-1-overexpressing mesenchymal stem cells (MSCs) into rat hearts after myocardial

infarction significantly reduced MMP-9 expression in the infarcted myocardium when compared with transplantation of MSCs expressing vector only (80).

The upregulation of HO-1 in advanced atherosclerotic plaques with a vulnerable phenotype might be an adaptive endogenous defense mechanism. Thus, further induction of HO-1 expression could aid in the stabilization of the atheromatous plaque. This hypothesis was tested in an apoE^{-/-} mouse model for vulnerable plaque formation. HO-1 inhibition by ZnPP augments lipid accumulation and increases necrotic core size. In contrast, HO-1 induction by CoPP or adenovirus-mediated gene transfer prevents plaque progression into vulnerable lesions by increasing fibrous cap thickness and intimal VSMC accumulation, whereas the necrotic core area and intraplaque lipid deposition is reduced (13). Supporting these findings, HO-1 has been shown to protect VSMCs from oxidative stress-induced apoptosis/cell death (6,93). Taken together, it is conceivable that by suppressing MMP-9 expression and by preventing VSMC death in the lesion HO-1 may thus promote plaque stability.

HO-1 in arterial injury

The proliferation and migration of VSMCs from the media into the intima contribute to arterial intimal thickening (24,41). Interestingly, Morita et al (59) show that VSMC-derived CO, a product of HO-1, inhibits mitogen endothelin-1 (ET-1) and platelet-derived growth factor-B (PDGF-B) expression of endothelial cells, which in turn inhibits VSMC proliferation. On the other hand, increasing CO production or exposing cells to exogenous CO leads to a markedly attenuated growth response of VSMCs (60), implicating HO-1 and its reaction product CO in the response to vascular injury. Indeed,

the in vivo protective role of HO-1 in vascular remodeling is demonstrated in a pig model of arterial injury (21). HO-1 gene transfer significantly reduced cellular proliferation in the intima and media. The decrease in cell proliferation in HO-1 transduced arteries is associated with a significant reduction in intimal lesion formation, indicating HO-1 confers protection against vascular injury through its effects on proliferation (21). Further supporting this notion, in response to femoral artery wire injury HO-1^{-/-} mice develop larger intimal lesions than wild type mice (21). Moreover, VSMCs from HO-1^{-/-} mice display enhanced DNA synthesis and proliferation compared with wild type VSMCs (21).

Following carotid artery balloon injury, wild-type Wistar rats (HsdBlu: GUNNJ/J) develop maximum neointimal hyperplasia with proliferating VSMC in the neointima (65). In contrast, the artery of hyperbilirubinemic Gunn rats (HsdBlu: GUNNj/j) shows very few proliferating cells and minimal neointimal hyperplasia compared with the control (65), suggesting bilirubin and its precursor biliverdin might have anti-proliferative effect during vascular remodeling. Supporting this notion, biliverdin is found to ameliorate neointimal hyperplasia associated with balloon injury and that bilirubin/biliverdin inhibits VSMC proliferation through cell cycle arrest (65). Thus, in addition to CO, bilirubin/biliverdin—another reaction product of HO-1—contributes to the reduction of neointima formation following arterial injury by its anti-proliferative effect on VSMCs.

In addition to proliferation, SMC migration contributes to the development of neointima after injury (24,41,67). PDGF, by inducing VSMC chemotaxis, is one of the growth factors involved in this process (41). Interestingly, generation of H₂O₂ is required for PDGF signal transduction (82). Therefore, reactive oxygen species (ROS) and HO-1

might be involved in regulating VSMC migration following arterial injury. In a recent study, Rodriguez et al showed that overexpression of HO-1 by adenovirus or chemical inducer CoPP results in decreased VSMC migration (74). They further show that CO is responsible for the anti-migratory effects seen with increased HO-1 expression. Since nicotinamide adenine dinucleotide phosphate (NADPH) oxidase is an important source of ROS generation (84), it is not surprising that the effect of HO-1/CO on inhibiting PDGF-induced VSMC migration is mediated by inhibition of NADPH oxidase (74).

HO-1 in in-stent restenosis and vein graft stenosis

Percutaneous coronary angioplasty to restore blood flow is a routine procedure to treat coronary artery disease, however, recurrent stenosis remains a major drawback in percutaneous transluminal angioplasty. Although stenting provides additional benefit by decreasing reocclusion (25), an acute coronary syndrome presentation in patients with in-stent restenosis is associated with a higher incidence of recurrent adverse cardiovascular events and angiographic restenosis (3). Drug-eluting stents with antiproliferative drugs reduce the rates of restenosis and neointimal hyperplasia and associated clinical events (61,69). However, questions regarding the long-term safety of drug-eluting stents have been raised (33,44). To reduce in-stent restenosis without compromising re-endothelialization, Hyvelin et al recently investigated the potential of HO-1 induction in this respect (37). In rat and rabbit models of in-stent stenosis, treatment with hemin, a potent HO-1 inducer, reduces neointima growth without compromising re-endothelialization of the stented arteries (37). Hemin increases HO-1 expression and limits the early inflammatory, apoptotic, and proliferative cellular events that are

associated with in-stent stenosis (37). SnPP, an HO-1 inhibitor, abolished the beneficial effect afforded by hemin. Furthermore, a CO donor also reduces in-stent stenosis in rat aorta, suggesting that CO might contribute to the protection. Their results suggest that HO-1 plays an important role in limiting in-stent stenosis and might serve as a novel target after endovascular therapies (37). This conclusion is supported by a previous report that patients with long (GT) repeat length polymorphism in the HO-1 promoter (see the session: HO-1 and microsatellite polymorphism in humans) have lower transcriptional activity and have a greater risk for angiographic restenosis as well as adverse cardiac events after coronary stenting (10).

Although coronary artery stenting is increasingly being used to treat patients with obstructive atherosclerotic lesions, stenting is associated with a greater need for repeated procedures in comparison with coronary artery bypass graft surgery (CABG) (79). A recent study showed that for patients with multivessel disease, CABG continues to be associated with lower mortality rates than does treatment with drug-eluting stents and is also associated with lower rates of death or myocardial infarction and repeat revascularization (33). Therefore, CABG remains an important treatment for multivessel disease and autologous vein grafts provide important and convenient conduits for bypass graft surgery (1,70). However, vein graft occlusion does develop due to intimal thickening (73). Given that HO-1 is induced in the neointima of the vein grafts in a mouse jugular vein/carotid artery autograft model (93), which mimics the vein graft used in bypass surgery, we hypothesized that HO-1 may play a role in the adaptation of VSMCs to hemodynamic stress in vascular wall remodeling. Consistent with the notion that HO-1 and its product CO inhibit VSMC proliferation (60), in response to similar

hemodynamic stress 10 days after grafting, HO-1^{-/-} mice have much larger neointima than wild type mice (93). Surprisingly, instead of developing even larger lesions 14 days after surgery, there is massive cell death in the neointima of HO-1^{-/-} mice (93), suggesting that lack of HO-1 might render VSMCs more prone to oxidant (generated by hemodynamic stress)-induced cell death. Indeed, HO-1^{-/-} VSMCs are more sensitive to H₂O₂-induced cell death, indicating that susceptibility to oxidative stress is one of the mechanisms leading to cell death in the absence of HO-1 (93).

HO-1 in transplant arteriosclerosis

Chronic allograft rejection is mainly manifested by progressive arteriosclerosis; thus reducing transplant arteriosclerosis would improve graft survival. Interestingly, in an aortic allotransplant model, Allotrap peptide RDP58 therapy markedly inhibits vascular intimal thickening, media necrosis, and adventitial cellular inflammation. The attenuation of arteriosclerosis is associated with the induction of HO-1 expression (48). Using a rat aorta chronic rejection model, Bouche et al show that specific HO-1 overexpression following gene transfer of HO-1 inhibits chronic rejection by reducing leukocyte and VSMC infiltration of the aorta intima (5). In another study using rat aortic transplant model, adenoviral gene transfer of HO-1 results in a significant reduction in leukocyte infiltration and a decreased number of VSMCs in the intima, and significantly lower levels of NF-κB and fewer apoptotic cells in the aortas (20). The protective role of HO-1 in inhibiting graft arteriosclerosis seems to correlate with reduced levels of NF-κB and inhibition of apoptosis in the grafts (20). A study showed that donor HO-1 expression has a direct influence on the recipient immune response. Endothelial cells

overexpressing HO-1 significantly inhibits proliferation and interferon- γ production in allogeneic CD8(+) T cells, both are important in chronic rejection (16). Their data suggest that donor HO-1 expression may be useful to augment other immunosuppressive therapies to prolong graft survival and inhibit intimal hyperplasia (16). Cheng et al (14) further demonstrated the important role of HO-1 in the regulation of vascular alloimmune response elicited by dendritic cells. In a murine model for transplantation arteriosclerosis, adoptive transfer of HO1^{-/-} dendritic cells before allograft transplantation is associated with pronounced intragraft CD4(+) T cell infiltration and increased IgG deposition, suggestive of an accelerated development of vasculopathy toward the chronic phase (14).

HO-1 in thrombosis

Thrombosis is induced after vascular injury and leads to several cardiovascular diseases associated with stroke, myocardial infarction, and venous thromboembolic disorders (26). Thrombus formation involves the interaction of injured vessel and platelets. Tissue factors, von willebrand factor, and fibrinogen trigger platelets activation and aggregation. Several lines of evidence suggest a role of HO-1 in thrombosis. Using ex vivo blood clotting perfusion chamber, induction of HO-1 expression has been found to inhibit platelet-dependent thrombosis (71). In a mouse-to-rat cardiac transplant model, inhibition of HO-1 activity by SnPP causes graft rejection, which is associated with platelet aggregation and thrombosis of coronary arterioles (77). Exogenous CO reverses the SnPP effect, including inhibition of platelet aggregation and thrombosis, suggesting CO generated by HO-1 suppresses graft rejection by inhibiting platelet

aggregation and vascular thrombosis (77). HO-1 gene transfer into injured vessels of apoE^{-/-} mice immediately after angioplasty results in earlier thrombolysis and restoration of blood flow (12). Furthermore, CO mediates this thrombolytic effect through inhibiting plasminogen activator inhibitor-1 expression and fibrin deposition (12). Bilirubin, another byproduct of HO-1, also contributes to the anti-thrombotic property of HO-1, because bilirubin was found as effective as HO-1 induction in delaying ferric chloride-induced thrombus formation (52). To investigate the direct function of HO-1 on thrombosis, True et al used photochemical-induced vascular injury in wild type and HO-1^{-/-} mice. HO-1 deficiency leads to accelerated, occlusive arterial thrombus formation compared with wild type mice (87). In addition, inhaled CO and biliverdin administration rescue the prothrombotic phenotype in HO-1^{-/-} mice (87). Confirming the role of HO-1/CO in protection against vascular arterial thrombosis, adoptive transfer of HO-1-expressing platelets rescues HO-1^{-/-} mice from arterial thrombosis in allogeneic aortic transplantation (8). In addition to arterial thrombosis, HO-1 also has a role in venous thrombosis. In a model of stasis-induced thrombosis created by ligation of the inferior vena cava, HO-1 deficiency impairs thrombus resolution and exaggerates the inflammatory response to thrombus formation (86). The clinical relevance is demonstrated in a prospective cohort study that patients with long (GT) repeat alleles in HO-1 gene promoter have an increased risk of recurrent venous thromboembolism (62).

HO-1 and microsatellite polymorphism in humans

A number of studies associate human HO-1 gene promoter polymorphisms with risk for vascular diseases (11,22). A microsatellite DNA with dinucleotide (GT) repeat in

the promoter region of the HO-1 gene shows a length polymorphism that modulates the level of HO-1 gene transcription (11,22). Patients with short (<25 GT) dinucleotide repeats in the HO-1 gene promoter on either allele have higher expression levels of HO-1 and significantly less often restenosis than patients with longer (\geq 25 GT) dinucleotide repeats (22) and coronary atherosclerosis (7). In a Hisayama cohort study, HO-1 expression is found to be intimately associated with atherogenesis and may play an important role as an adaptive molecule in the inflammatory/repair process (81). The significance of (GT)_n polymorphism in the HO-1 promoter was further demonstrated in a recent study that (GT)_n allelic variants of the promoter directly modulate HO-1 expression levels and the proangiogenic and antiinflammatory functions of HO-1 in human endothelium (83). Thus, the capacity to upregulate HO-1 expression may be genetically regulated and reduced ability to induce HO-1 may be involved in the mechanism of coronary atherosclerosis (7,11,22,83).

HO-1 and hypoxia in the heart

Reduced blood flow due to lumen narrowing from arterial thrombosis or atherosclerotic lesions leads to myocardial hypoxia. A potential function of HO-1 in cardiomyocytes was implicated when primary rat neonatal cardiomyocytes respond to hypoxia with increased expression levels of HO-1 mRNA and protein while under normal physiological conditions, HO-1 is expressed at low levels in cardiomyocytes (4). Hypoxia causes pulmonary hypertension and induces right ventricular hypertrophy. To gain insight into the role of HO-1 in cardiac adaptation to hypoxic stress, we generated HO-1^{-/-} mice and subjected the mice to chronic hypoxia in an early study (94). Although

the myocardium in HO-1^{-/-} mice appears to be normal under normoxic conditions, under hypoxic conditions, severe right ventricular dilatation and infarcts with mural thrombi develop in HO-1^{-/-}, but not wild type mice, despite similar degree of pulmonary hypertension after hypoxia exposure (94) (Fig. 2). In addition to lipid peroxidation and oxidative damage in the right ventricular cardiomyocytes of HO-1^{-/-} mice, we also detected apoptotic cardiomyocytes surrounding areas of infarcted myocardium. These data suggest that the absence of HO-1 in cardiomyocytes leads to an accumulation of ROS that cause cardiomyocyte death and the high percentage of right ventricular mural thrombi in HO-1^{-/-} mice maybe due to increased platelet aggregation (94).

HO-1 in cardiac ischemia and reperfusion injury

The results from hypoxia studies led us to test the hypothesis that HO-1 may play a central role in cardiac homeostasis (95). For gain of function experiments, we generated cardiac-specific transgenic mice overexpressing different levels of HO-1 (95). To determine whether overexpression of HO-1 protects against postischemic injury and to exclude the involvement of inflammatory components on reperfusion, we used an isolated perfused heart preparation by subjecting hearts to 30 minutes of global ischemia, followed by 40 minutes of reperfusion (95). Compared with wild type hearts, postischemic recovery of cardiac function in all three lines of HO-1 transgenic hearts improved in an HO-1 dose-dependent manner (95) (Fig. 3). In a myocardial infarction model, in contrast to the large infarcts in wild type mouse hearts, HO-1 transgenic mouse hearts show small infarcts (95) (Fig. 4A and B). Despite a similar percentage of left ventricle at risk between wild type and transgenic mouse hearts, the infarct size is

significantly reduced in transgenic mice compared with wild type mice (95) (Fig. 4C).

Our study clearly demonstrate that overexpression of HO-1 in the cardiomyocyte protects against ischemia and reperfusion injury, thus improving the recovery of cardiac function (95).

To explore the therapeutic potential of HO-1 gene transfer in long-term myocardial protection, Luis et al took advantage of an adeno-associated virus (AAV)-mediated delivery method (58). AAV-mediated transfer of the HO-1 gene into normal rat hearts 8 weeks before ischemia/reperfusion injury leads to a dramatic reduction in left ventricular myocardial infarction (58), supporting the findings of our transgenic studies (95). The HO-1-mediated protection from myocardial ischemia/reperfusion injury is associated with a decrease in oxidative stress and proapoptotic and proinflammatory protein levels (58). These findings demonstrate the therapeutic potential of AAV-mediated HO-1 gene transfer for sustained cardiac protection from ischemic injury and introduce the concept of pre-event gene therapy in protection against future injury (58).

After showing that predelivery of HO-1 by AAV to the heart can markedly reduce injury after acute myocardial infarction (58), Liu et al investigated the effect of HO-1 gene delivery on postinfarction recovery in a follow-up study (53). Interestingly, AAV-mediated HO-1 gene transfer normalizes postinfarction echocardiographic left ventricular function and chamber dimensions 3 months after myocardial infarction by preventing ventricular remodeling (53). When the effect of HO-1 gene delivery was assessed one year after acute myocardial infarction, mortality was markedly reduced in the HO-1-treated animals compared with the LacZ-treated animals (54). The therapeutic efficacy of preemptive AAV-HO-1 delivery is further demonstrated by the findings that

chronic recurrent myocardial ischemic injury is significantly attenuated by AAV-HO-1 delivery (68). These results suggest that preemptive HO-1 gene delivery may be useful as a therapeutic strategy to reduce post-myocardial infarction left ventricular remodeling and heart failure.

HO-1 in diabetic heart

Myocardial infarction and heart disease is one of the chronic complications of diabetes. Given the cardioprotective role of HO-1 against ischemia/reperfusion injury, we tested the role of HO-1 in the settings of diabetes (55). Although an absence of HO-1 significantly increases infarct size in normoglycemic mice, diabetes exacerbates myocardial infarction in the setting of HO-1 deficiency (55). In addition to exaggerated infarct size, mortality is 2-fold higher in diabetic HO-1^{-/-} than wild type mice after ischemia/reperfusion injury. Interestingly, the mortality rate of diabetic patients suffering from myocardial infarction due to ischemia/reperfusion injury is 2-fold higher than that of nondiabetic patients. Importantly, we observed 55% of diabetic HO-1^{-/-} mice that survived ischemia/reperfusion developed left ventricular thrombi (55) (Fig. 5). Intriguingly, It has been suggested that increased mortality rate in diabetic patients is in part due to the tendency toward thrombosis (2). Corroborating our findings, HO-1 inhibition in streptozotocin-induced diabetic rat further enhances infarct size during ischemia/reperfusion (18). It was found that hyperglycemia predisposed the heart to produce high levels of both the cytokines IL-1 β and CXCL8 and subsequent ischemia/reperfusion further increases the cytokine production (18).

In humans, type 2 diabetic patients with longer (GT) repeats of the HO-1 gene promoter (with lower HO-1 inducibility) are shown to have higher oxidative stress and increased susceptibility to coronary artery disease (11). In a subsequent study, by measuring serum bilirubin and ferritin levels, Chen et al links HO-1 gene promoter polymorphism and the susceptibility to coronary artery disease in diabetic patients (9). Hisayama study examining HO-1 expression in coronary atherosclerotic lesions of Japanese autopsies reveals that HO-1 is intimately associated with intraplaque in patients with diabetes (81).

HO-1 in re-endothelialization, angiogenesis, and the heart

Endothelial cell damage is an important pathophysiological step of atherosclerosis and restenosis after angioplasty and thus endothelial regeneration is critical in repairing injured vessels and protecting from the development of an intimal lesion (34). Increasing evidence suggest that endothelial progenitor cells (EPCs) derived from bone marrow augment neovascularization of tissue after ischemia and contribute to re-endothelialization after endothelial injury, thereby providing a novel therapeutic option (88).

Since human HO-1 deficiency had elevated oxidative stress with severe endothelial cell damage (45,92), it is conceivable that HO-1 may affect mobilization and homing of circulating EPCs. As demonstrated by Lin et al (51), systemic overexpression of HO-1 with adenovirus in mice leads to an acceleration of re-endothelialization of denuded vessels due to enhanced EPC mobilization; conversely, lack of HO-1 in mice results in significant attenuation of EPC mobilization and re-endothelialization.

Additionally, exposing mice to CO prior to carotid injury is also able to increase EPC mobilization and enhance re-endothelialization (51). Using a rabbit model of aortic balloon injury and pharmacological induction of HO-1, another study came to the same conclusion that HO-1 contributes to vascular repair by increasing the number and maturation of circulating EPCs derived from the bone marrow (91). Consistent with these findings, bone marrow cells from HO-1^{-/-} mice, compared with wild type mice, generate fewer endothelial colony-forming cells (91).

In addition to a role in promoting re-endothelialization, overexpression of HO-1 in coronary microvessel endothelial cells enhances endothelial cell proliferation and capillary formation, linking HO-1 to angiogenesis (17). Given the role of HO-1 in cardioprotection and its angiogenic potential, HO-1 may enhance myocardial angiogenesis in ischemic myocardium. The role of HO-1 in improving myocardial angiogenesis was supported by a resveratrol study. Resveratrol, a polyphenolic compound, enhances angiogenesis in the infarcted rat myocardium by induction of vascular endothelial growth factor (VEGF), and the induction of VEGF is mediated by thioredoxin-1 and HO-1 (43). In a hypercholesterolemic myocardial infarction model, secoisolariciresinol diglucoside treatment, a compound isolated from omega-3 fatty acids-rich flaxseed, increases capillary and arteriolar density and improves left ventricular function (72). The cardiac functional improvement might be due to increased HO-1, VEGF and p-eNOS expression (72). These studies emphasize that HO-1 mediates ischemic myocardial angiogenesis after myocardial infarction. Using heterozygous Flt-1 mice with reduced expression of Flt-1, Thirunavukkarasu et al found that reduced expression of Flt-1 results in less ventricular functional recovery compared

to the wild type mice. The reduced functional recovery paralleled with increased myocardial infarction and apoptosis (85). Interestingly, heterozygous Flt-1 mice have pronounced inhibition of the expression of HO-1, iNOS, p-AKT and p-eNOS following ischemia/reperfusion injury (85).

HO-1 gene transfer has been shown to promote neovascularization and ventricular function in the ischemic heart of mice receiving HO-1, due to higher cardiac levels of VEGF and stromal cell-derived factor-1 (SDF-1) (50). Concomitant treatment with both VEGF and SDF-1 neutralizing antibodies attenuates the protective effect of HO-1 to a greater extent than treatment with either neutralizing antibody alone (50). This emphasizes the cooperative roles of these two factors in HO-1-mediated protection. Furthermore, increased recruitment of circulating c-kit⁺ stem/progenitor cells to the infarct area in mice receiving HO-1 gene transfer suggests that EPCs participate in the angiogenic effect (50). A follow up study showed that EPC mobilization and re-endothelialization are significantly attenuated in HO-1^{-/-} mice after vascular injury, which is rescued by exposing mice to CO prior to carotid injury (51). The role of CO in myocardial angiogenesis is further supported by that pretreatment of CO-donor increases accumulation of c-kit⁺ stem/progenitor cells in the infarct areas following myocardial infarction (46). These c-kit⁺ cells are able to differentiate into VSMCs and contribute to the formation of new coronary arteries in infarct areas. The induction of proangiogenic factors hypoxia inducible factor (HIF)-1 α , SDF-1, and VEGF-B was evident in CO-donor pretreated hearts. Thus, HO-1 and CO promote neovascularization after myocardial infarction by modulating the expression of HIF-1 α , SDF-1 and VEGF-B (46). Therapeutic angiogenesis is a novel strategy for treatment of patients with

ischemic heart disease. Since HO-1-mediated angiogenesis protects ischemic myocardium, HO-1 and its product, CO, could serve as targets for the treatment of heart disease.

Therapeutic potential of HO-1

Given the cardiovascular protective function of HO-1, HO-1 offers a great therapeutic potential for cardiovascular disease. Overexpression via gene delivery methods has proven the beneficial effects of HO-1 in many animal disease models; however, therapeutic gene delivery methods in humans are not readily available. Therefore, targeted induction of HO-1 by pharmacological means might be a more promising alternative. For example, in addition to the cholesterol-lowering effects, various statins, hydroxymethylglutaryl coenzyme A (HMG-CoA) reductase inhibitors, also have antiinflammatory and antiproliferative effects. Intriguingly, these effects of statins are largely mediated through induced HO-1 expression in vascular cells (29,30,47). Furthermore, atorvastatin strongly induces angiogenesis likely via HO-1 induction, indicating a potential for therapeutic angiogenesis in ischemic diseases (57). Other than statins, another cholesterol-lowering drug probucol reduces cardiovascular disease incidence in hypercholesterolemic patients (78) and interestingly, HO-1 is the molecular target of probucol (90). Moreover, induction of HO-1 by aspirin may be a novel mechanism by which aspirin prevents cellular injury under inflammatory conditions and in cardiovascular disease (28). An alternative for HO-1 induction is to inhibit its degradation by increasing mRNA stability. Indeed, simvastatin-dependent upregulation of HO-1 was found to be mediated by stabilization of HO-1 mRNA via a

PI3K/Akt-dependent signaling pathway (36). Taken together, pharmacological induction of HO-1 holds a promise as a therapeutic intervention for treating cardiovascular disease. For a more refined therapy, time-dependent and tissue-specific induction of HO-1 is needed and thus challenges remain for the development of time- and tissue-specific pharmacological agents for HO-1 induction.

Conclusion

Collective evidence from the past two decades has demonstrated the central role of HO-1 in protection against disease. We have highlighted the protective function of HO-1 against cardiovascular diseases. Various stresses, including hypoxia, ischemia/reperfusion injury, diabetes, hypercholesterolemia, etc, lead to the production of ROS in the cardiovascular system (Fig. 6). Increased ROS in turn induces inflammation and apoptosis, among others. In the absence of HO-1, perhaps through the excessive accumulation of ROS, cardiovascular cells have a maladaptive response, resulting in cell death. In the presence of the buffering effect of HO-1, normal adaptive response provides protection to cardiovascular system (Fig. 6). As such, overexpression of HO-1 may protect against cardiovascular diseases caused by ROS. The clinical significance of HO-1 in human cardiovascular disease was highlighted in a child with HO-1 deficiency (45,92). Although the therapeutic potential of HO-1 is clear, developing HO-1 therapeutics would greatly benefit patients suffering from cardiovascular disease.

Acknowledgments

This work was supported in part by National Health Research Institutes (Taiwan) grant CS-099-PP-04 (to S.-F.Y.) and National Science Council (Taiwan) grant 98-2320-B-400-004-MY3 (to S.-F.Y.).

List of Abbreviations

AAV = adeno-associated virus

apoE^{-/-} = apolipoprotein E-deficient

CABG = coronary artery bypass graft surgery

CO = carbon monoxide

CoPP = Cobalt protoporphyrin

EPCs = endothelial progenitor cells

ET-1 = endothelin-1

GT = guanosine thymidine dinucleotide

HIF = hypoxia-inducible factor

HMG-CoA = hydroxymethylglutaryl coenzyme A

HO-1 = heme oxygenase-1

LDL = low density lipoprotein

MMPs = matrix metalloproteinases

NADPH oxidase = nicotinamide adenine dinucleotide phosphate oxidase

oxLDL = oxidized LDL

PDGF-B = platelet-derived growth factor-B

ROS = reactive oxygen species

SDF-1 = stromal cell-derived factor-1

SMC = smooth muscle cell

SnPP = Sn-protoporphyrin IX

TIMP = tissue inhibitor of metalloproteinase

VEGF = vascular endothelial growth factor

VSMC = vascular smooth muscle cell

ZnPP = Zinc protoporphyrin

References

1. Angelini GD, Jeremy JY. Towards the treatment of saphenous vein bypass graft failure--a perspective of the Bristol Heart Institute. *Biorheology* 39: 491-9, 2002.
2. Aronson D, Rayfield EJ, Chesebro JH. Mechanisms determining course and outcome of diabetic patients who have had acute myocardial infarction. *Ann Intern Med* 126: 296-306, 1997.
3. Assali AR, Moustapha A, Sdringola S, Denktas AE, Willerson JT, Holmes DR, Jr., Smalling RW. Acute coronary syndrome may occur with in-stent restenosis and is associated with adverse outcomes (the PRESTO trial). *Am J Cardiol* 98: 729-33, 2006.
4. Borger DR, Essig DA. Induction of HSP 32 gene in hypoxic cardiomyocytes is attenuated by treatment with N-acetyl-L-cysteine. *Am J Physiol* 274: H965-73, 1998.
5. Bouche D, Chauveau C, Roussel JC, Mathieu P, Braudeau C, Tesson L, Soullillou JP, Iyer S, Buelow R, Anegon I. Inhibition of graft arteriosclerosis development in

- rat aortas following heme oxygenase-1 gene transfer. *Transpl Immunol* 9: 235-8, 2002.
6. Brunt KR, Fenrich KK, Kiani G, Tse MY, Pang SC, Ward CA, Melo LG. Protection of human vascular smooth muscle cells from H₂O₂-induced apoptosis through functional codependence between HO-1 and AKT. *Arterioscler Thromb Vasc Biol* 26: 2027-34, 2006.
 7. Brydun A, Watari Y, Yamamoto Y, Okuhara K, Teragawa H, Kono F, Chayama K, Oshima T, Ozono R. Reduced expression of heme oxygenase-1 in patients with coronary atherosclerosis. *Hypertens Res* 30: 341-8, 2007.
 8. Chen B, Guo L, Fan C, Bolisetty S, Joseph R, Wright MM, Agarwal A, George JF. Carbon monoxide rescues heme oxygenase-1-deficient mice from arterial thrombosis in allogeneic aortic transplantation. *Am J Pathol* 175: 422-9, 2009.
 9. Chen YH, Chau LY, Chen JW, Lin SJ. Serum bilirubin and ferritin levels link heme oxygenase-1 gene promoter polymorphism and susceptibility to coronary artery disease in diabetic patients. *Diabetes Care* 31: 1615-20, 2008.
 10. Chen YH, Chau LY, Lin MW, Chen LC, Yo MH, Chen JW, Lin SJ. Heme oxygenase-1 gene promoter microsatellite polymorphism is associated with angiographic restenosis after coronary stenting. *Eur Heart J* 25: 39-47, 2004.
 11. Chen YH, Lin SJ, Lin MW, Tsai HL, Kuo SS, Chen JW, Charng MJ, Wu TC, Chen LC, Ding YA, Pan WH, Jou YS, Chau LY. Microsatellite polymorphism in promoter of heme oxygenase-1 gene is associated with susceptibility to coronary artery disease in type 2 diabetic patients. *Hum Genet* 111: 1-8, 2002.

12. Chen YH, Tsai HL, Chiang MT, Chau LY. Carbon monoxide-induced early thrombolysis contributes to heme oxygenase-1-mediated inhibition of neointimal growth after vascular injury in hypercholesterolemic mice. *J Biomed Sci* 13: 721-30, 2006.
13. Cheng C, Noordeloos AM, Jeney V, Soares MP, Moll F, Pasterkamp G, Serruys PW, Duckers HJ. Heme oxygenase 1 determines atherosclerotic lesion progression into a vulnerable plaque. *Circulation* 119: 3017-27, 2009.
14. Cheng C, Noorderloos M, van Deel ED, Tempel D, den Dekker W, Wagtmans K, Duncker DJ, Soares MP, Laman JD, Duckers HJ. Dendritic cell function in transplantation arteriosclerosis is regulated by heme oxygenase 1. *Circ Res* 106: 1656-66, 2010.
15. Choudhary S, Higgins CL, Chen IY, Reardon M, Lawrie G, Vick GW, 3rd, Karmonik C, Via DP, Morrisett JD. Quantitation and localization of matrix metalloproteinases and their inhibitors in human carotid endarterectomy tissues. *Arterioscler Thromb Vasc Biol* 26: 2351-8, 2006.
16. Clarke HM, Shrivastava S, Motterlini R, Sawle P, Chen D, Dorling A. Donor HO-1 expression inhibits intimal hyperplasia in unmanipulated graft recipients: a potential role for CD8+ T-cell modulation by carbon monoxide. *Transplantation* 88: 653-61, 2009.
17. Deramaudt BM, Braunstein S, Remy P, Abraham NG. Gene transfer of human heme oxygenase into coronary endothelial cells potentially promotes angiogenesis. *J Cell Biochem* 68: 121-7, 1998.

18. Di Filippo C, Marfella R, Cuzzocrea S, Piegari E, Petronella P, Giugliano D, Rossi F, D'Amico M. Hyperglycemia in streptozotocin-induced diabetic rat increases infarct size associated with low levels of myocardial HO-1 during ischemia/reperfusion. *Diabetes* 54: 803-10, 2005.
19. Dollery CM, Libby P. Atherosclerosis and proteinase activation. *Cardiovasc Res* 69: 625-35, 2006.
20. Du D, Chang S, Chen B, Zhou H, Chen ZK. Adenovirus-mediated heme oxygenase transfer inhibits graft arteriosclerosis in rat aortic transplants. *Transplant Proc* 39: 3446-8, 2007.
21. Duckers HJ, Boehm M, True AL, Yet SF, San H, Park JL, Clinton Webb R, Lee ME, Nabel GJ, Nabel EG. Heme oxygenase-1 protects against vascular constriction and proliferation. *Nat Med* 7: 693-8, 2001.
22. Exner M, Schillinger M, Minar E, Mlekusch W, Schlerka G, Haumer M, Mannhalter C, Wagner O. Heme oxygenase-1 gene promoter microsatellite polymorphism is associated with restenosis after percutaneous transluminal angioplasty. *J Endovasc Ther* 8: 433-40, 2001.
23. Fabunmi RP, Sukhova GK, Sugiyama S, Libby P. Expression of tissue inhibitor of metalloproteinases-3 in human atheroma and regulation in lesion-associated cells: a potential protective mechanism in plaque stability. *Circ Res* 83: 270-8, 1998.
24. Ferns GA, Raines EW, Sprugel KH, Motani AS, Reidy MA, Ross R. Inhibition of neointimal smooth muscle accumulation after angioplasty by an antibody to PDGF. *Science* 253: 1129-32, 1991.

25. Fischman DL, Leon MB, Baim DS, Schatz RA, Savage MP, Penn I, Detre K, Veltri L, Ricci D, Nobuyoshi M, et al. A randomized comparison of coronary-stent placement and balloon angioplasty in the treatment of coronary artery disease. Stent Restenosis Study Investigators. *N Engl J Med* 331: 496-501, 1994.
26. Fuster V, Lewis A. Conner Memorial Lecture. Mechanisms leading to myocardial infarction: insights from studies of vascular biology. *Circulation* 90: 2126-46, 1994.
27. Galis ZS, Sukhova GK, Lark MW, Libby P. Increased expression of matrix metalloproteinases and matrix degrading activity in vulnerable regions of human atherosclerotic plaques. *J Clin Invest* 94: 2493-503, 1994.
28. Grosser N, Abate A, Oberle S, Vreman HJ, Dennery PA, Becker JC, Pohle T, Seidman DS, Schroder H. Heme oxygenase-1 induction may explain the antioxidant profile of aspirin. *Biochem Biophys Res Commun* 308: 956-60, 2003.
29. Grosser N, Erdmann K, Hemmerle A, Berndt G, Hinkelmann U, Smith G, Schroder H. Rosuvastatin upregulates the antioxidant defense protein heme oxygenase-1. *Biochem Biophys Res Commun* 325: 871-6, 2004.
30. Grosser N, Hemmerle A, Berndt G, Erdmann K, Hinkelmann U, Schurger S, Wijayanti N, Immenschuh S, Schroder H. The antioxidant defense protein heme oxygenase 1 is a novel target for statins in endothelial cells. *Free Radic Biol Med* 37: 2064-71, 2004.
31. Grundemar L, Ny L. Pitfalls using metalloporphyrins in carbon monoxide research. *Trends Pharmacol Sci* 18: 193-5, 1997.

32. Han J, Hajjar DP, Febbraio M, Nicholson AC. Native and modified low density lipoproteins increase the functional expression of the macrophage class B scavenger receptor, CD36. *J Biol Chem* 272: 21654-9, 1997.
33. Hannan EL, Wu C, Walford G, Culliford AT, Gold JP, Smith CR, Higgins RS, Carlson RE, Jones RH. Drug-eluting stents vs. coronary-artery bypass grafting in multivessel coronary disease. *N Engl J Med* 358: 331-41, 2008.
34. Haudenschild CC, Schwartz SM. Endothelial regeneration. II. Restitution of endothelial continuity. *Lab Invest* 41: 407-18, 1979.
35. Henney AM, Wakeley PR, Davies MJ, Foster K, Hembry R, Murphy G, Humphries S. Localization of stromelysin gene expression in atherosclerotic plaques by in situ hybridization. *Proc Natl Acad Sci U S A* 88: 8154-8, 1991.
36. Hinkelmann U, Grosser N, Erdmann K, Schroder H, Immenschuh S. Simvastatin-dependent up-regulation of heme oxygenase-1 via mRNA stabilization in human endothelial cells. *Eur J Pharm Sci* 41: 118-24, 2010.
37. Hyvelin JM, Maurel B, Uzbekov R, Motterlini R, Lermusiaux P. Hemin prevents in-stent stenosis in rat and rabbit models by inducing heme-oxygenase-1. *J Vasc Surg* 51: 417-28, 2010.
38. Ishikawa K, Navab M, Leitinger N, Fogelman AM, Lusis AJ. Induction of heme oxygenase-1 inhibits the monocyte transmigration induced by mildly oxidized LDL. *J Clin Invest* 100: 1209-16, 1997.
39. Ishikawa K, Sugawara D, Goto J, Watanabe Y, Kawamura K, Shiomi M, Itabe H, Maruyama Y. Heme oxygenase-1 inhibits atherogenesis in Watanabe heritable hyperlipidemic rabbits. *Circulation* 104: 1831-6, 2001.

40. Ishikawa K, Sugawara D, Wang X, Suzuki K, Itabe H, Maruyama Y, Luscis AJ. Heme oxygenase-1 inhibits atherosclerotic lesion formation in LDL-receptor knockout mice. *Circ Res* 88: 506-12, 2001.
41. Jawien A, Bowen-Pope DF, Lindner V, Schwartz SM, Clowes AW. Platelet-derived growth factor promotes smooth muscle migration and intimal thickening in a rat model of balloon angioplasty. *J Clin Invest* 89: 507-11, 1992.
42. Juan SH, Lee TS, Tseng KW, Liou JY, Shyue SK, Wu KK, Chau LY. Adenovirus-mediated heme oxygenase-1 gene transfer inhibits the development of atherosclerosis in apolipoprotein E-deficient mice. *Circulation* 104: 1519-25, 2001.
43. Kaga S, Zhan L, Matsumoto M, Maulik N. Resveratrol enhances neovascularization in the infarcted rat myocardium through the induction of thioredoxin-1, heme oxygenase-1 and vascular endothelial growth factor. *J Mol Cell Cardiol* 39: 813-22, 2005.
44. Kastrati A, Mehilli J, Pache J, Kaiser C, Valgimigli M, Kelbaek H, Menichelli M, Sabate M, Suttorp MJ, Baumgart D, Seyfarth M, Pfisterer ME, Schomig A. Analysis of 14 trials comparing sirolimus-eluting stents with bare-metal stents. *N Engl J Med* 356: 1030-9, 2007.
45. Kawashima A, Oda Y, Yachie A, Koizumi S, Nakanishi I. Heme oxygenase-1 deficiency: the first autopsy case. *Hum Pathol* 33: 125-30, 2002.
46. Lakkisto P, Kytö V, Forsten H, Siren JM, Segersvard H, Voipio-Pulkki LM, Laine M, Pulkki K, Tikkanen I. Heme oxygenase-1 and carbon monoxide promote neovascularization after myocardial infarction by modulating the expression of HIF-1 α , SDF-1 α and VEGF-B. *Eur J Pharmacol* 635: 156-64, 2010.

47. Lee TS, Chang CC, Zhu Y, Shyy JY. Simvastatin induces heme oxygenase-1: a novel mechanism of vessel protection. *Circulation* 110: 1296-302, 2004.
48. Li W, Iyer S, Lu L, Buelow R, Fung JJ, Rao AS, Woo J, Qian S. Attenuation of aortic graft arteriosclerosis by systemic administration of Allotrap peptide RDP58. *Transpl Int* 16: 849-56, 2003.
49. Lin CW, Shen SC, Hou WC, Yang LY, Chen YC. Heme oxygenase-1 inhibits breast cancer invasion via suppressing the expression of matrix metalloproteinase-9. *Mol Cancer Ther* 7: 1195-206, 2008.
50. Lin HH, Chen YH, Chang PF, Lee YT, Yet SF, Chau LY. Heme oxygenase-1 promotes neovascularization in ischemic heart by coinduction of VEGF and SDF-1. *J Mol Cell Cardiol* 45: 44-55, 2008.
51. Lin HH, Chen YH, Yet SF, Chau LY. After vascular injury, heme oxygenase-1/carbon monoxide enhances re-endothelialization via promoting mobilization of circulating endothelial progenitor cells. *J Thromb Haemost* 7: 1401-8, 2009.
52. Lindenblatt N, Bordel R, Schareck W, Menger MD, Vollmar B. Vascular heme oxygenase-1 induction suppresses microvascular thrombus formation in vivo. *Arterioscler Thromb Vasc Biol* 24: 601-6, 2004.
53. Liu X, Pachori AS, Ward CA, Davis JP, Gnechi M, Kong D, Zhang L, Murduck J, Yet SF, Perrella MA, Pratt RE, Dzau VJ, Melo LG. Heme oxygenase-1 (HO-1) inhibits postmyocardial infarct remodeling and restores ventricular function. *FASEB J* 20: 207-16, 2006.
54. Liu X, Simpson JA, Brunt KR, Ward CA, Hall SR, Kinobe RT, Barrette V, Tse MY, Pang SC, Pachori AS, Dzau VJ, Ogunyankin KO, Melo LG. Preemptive heme

- oxygenase-1 gene delivery reveals reduced mortality and preservation of left ventricular function 1 yr after acute myocardial infarction. *Am J Physiol Heart Circ Physiol* 293: H48-59, 2007.
55. Liu X, Wei J, Peng DH, Layne MD, Yet SF. Absence of heme oxygenase-1 exacerbates myocardial ischemia/reperfusion injury in diabetic mice. *Diabetes* 54: 778-84, 2005.
56. Maines MD. The heme oxygenase system: a regulator of second messenger gases. *Annu Rev Pharmacol Toxicol* 37: 517-54, 1997.
57. Matsumura M, Fukuda N, Kobayashi N, Umezawa H, Takasaka A, Matsumoto T, Yao EH, Ueno T, Negishi N. Effects of atorvastatin on angiogenesis in hindlimb ischemia and endothelial progenitor cell formation in rats. *J Atheroscler Thromb* 16: 319-26, 2009.
58. Melo LG, Agrawal R, Zhang L, Rezvani M, Mangi AA, Ehsan A, Griese DP, Dell'Acqua G, Mann MJ, Oyama J, Yet SF, Layne MD, Perrella MA, Dzau VJ. Gene therapy strategy for long-term myocardial protection using adeno-associated virus-mediated delivery of heme oxygenase gene. *Circulation* 105: 602-7, 2002.
59. Morita T, Kourembanas S. Endothelial cell expression of vasoconstrictors and growth factors is regulated by smooth muscle cell-derived carbon monoxide. *J Clin Invest* 96: 2676-82, 1995.
60. Morita T, Mitsialis SA, Koike H, Liu Y, Kourembanas S. Carbon monoxide controls the proliferation of hypoxic vascular smooth muscle cells. *J Biol Chem* 272: 32804-9, 1997.

61. Moses JW, Leon MB, Popma JJ, Fitzgerald PJ, Holmes DR, O'Shaughnessy C, Caputo RP, Kereiakes DJ, Williams DO, Teirstein PS, Jaeger JL, Kuntz RE. Sirolimus-eluting stents versus standard stents in patients with stenosis in a native coronary artery. *N Engl J Med* 349: 1315-23, 2003.
62. Mustafa S, Weltermann A, Fritsche R, Marsik C, Wagner O, Kyrle PA, Eichinger S. Genetic variation in heme oxygenase 1 (HMOX1) and the risk of recurrent venous thromboembolism. *J Vasc Surg* 47: 566-70, 2008.
63. Newby AC. Dual role of matrix metalloproteinases (matrixins) in intimal thickening and atherosclerotic plaque rupture. *Physiol Rev* 85: 1-31, 2005.
64. Newby AC. Metalloproteinase expression in monocytes and macrophages and its relationship to atherosclerotic plaque instability. *Arterioscler Thromb Vasc Biol* 28: 2108-14, 2008.
65. Ollinger R, Bilban M, Erat A, Froio A, McDaid J, Tyagi S, Csizmadia E, Graca-Souza AV, Liloia A, Soares MP, Otterbein LE, Usheva A, Yamashita K, Bach FH. Bilirubin: a natural inhibitor of vascular smooth muscle cell proliferation. *Circulation* 112: 1030-9, 2005.
66. Orozco LD, Kapturczak MH, Barajas B, Wang X, Weinstein MM, Wong J, Deshane J, Bolisetty S, Shaposhnik Z, Shih DM, Agarwal A, Lusis AJ, Araujo JA. Heme oxygenase-1 expression in macrophages plays a beneficial role in atherosclerosis. *Circ Res* 100: 1703-11, 2007.
67. Owens GK, Kumar MS, Wamhoff BR. Molecular regulation of vascular smooth muscle cell differentiation in development and disease. *Physiol Rev* 84: 767-801, 2004.

68. Pachori AS, Melo LG, Zhang L, Solomon SD, Dzau VJ. Chronic recurrent myocardial ischemic injury is significantly attenuated by pre-emptive adeno-associated virus heme oxygenase-1 gene delivery. *J Am Coll Cardiol* 47: 635-43, 2006.
69. Park SJ, Shim WH, Ho DS, Raizner AE, Park SW, Hong MK, Lee CW, Choi D, Jang Y, Lam R, Weissman NJ, Mintz GS. A paclitaxel-eluting stent for the prevention of coronary restenosis. *N Engl J Med* 348: 1537-45, 2003.
70. Patil CV, Nikolsky E, Boulos M, Grenadier E, Beyar R. Multivessel coronary artery disease: current revascularization strategies. *Eur Heart J* 22: 1183-97, 2001.
71. Peng L, Mundada L, Stomel JM, Liu JJ, Sun J, Yet SF, Fay WP. Induction of heme oxygenase-1 expression inhibits platelet-dependent thrombosis. *Antioxid Redox Signal* 6: 729-35, 2004.
72. Penumathsa SV, Koneru S, Zhan L, John S, Menon VP, Prasad K, Maulik N. Secoisolariciresinol diglucoside induces neovascularization-mediated cardioprotection against ischemia-reperfusion injury in hypercholesterolemic myocardium. *J Mol Cell Cardiol* 44: 170-9, 2008.
73. Rodes-Cabau J, Facta A, Larose E, DeLarochelliere R, Dery JP, Nguyen CM, Roy L, Proulx G, Gleeton O, Barbeau G, Noel B, Rouleau J, Boudreault JR, Bertrand OF. Predictors of aorto-saphenous vein bypass narrowing late after coronary artery bypass grafting. *Am J Cardiol* 100: 640-5, 2007.
74. Rodriguez AI, Gangopadhyay A, Kelley EE, Pagano PJ, Zuckerbraun BS, Bauer PM. HO-1 and CO decrease platelet-derived growth factor-induced vascular smooth

- muscle cell migration via inhibition of Nox1. *Arterioscler Thromb Vasc Biol* 30: 98-104, 2010.
75. Ross R. Cell biology of atherosclerosis. *Annu Rev Physiol* 57: 791-804, 1995.
76. Ross R. Atherosclerosis—an inflammatory disease. *N Engl J Med* 340: 115-26, 1999.
77. Sato K, Balla J, Otterbein L, Smith RN, Brouard S, Lin Y, Csizmadia E, Sevigny J, Robson SC, Vercellotti G, Choi AM, Bach FH, Soares MP. Carbon monoxide generated by heme oxygenase-1 suppresses the rejection of mouse-to-rat cardiac transplants. *J Immunol* 166: 4185-94, 2001.
78. Sawayama Y, Shimizu C, Maeda N, Tatsukawa M, Kinukawa N, Koyanagi S, Kashiwagi S, Hayashi J. Effects of probucol and pravastatin on common carotid atherosclerosis in patients with asymptomatic hypercholesterolemia. Fukuoka Atherosclerosis Trial (FAST). *J Am Coll Cardiol* 39: 610-6, 2002.
79. Serruys PW, Unger F, Sousa JE, Jatene A, Bonnier HJ, Schonberger JP, Buller N, Bonser R, van den Brand MJ, van Herwerden LA, Morel MA, van Hout BA. Comparison of coronary-artery bypass surgery and stenting for the treatment of multivessel disease. *N Engl J Med* 344: 1117-24, 2001.
80. Shu T, Zeng B, Ren X, Li Y. HO-1 modified mesenchymal stem cells modulate MMPs/TIMPs system and adverse remodeling in infarcted myocardium. *Tissue Cell* 42: 217-22, 2010.
81. Song J, Sumiyoshi S, Nakashima Y, Doi Y, Iida M, Kiyohara Y, Sueishi K. Overexpression of heme oxygenase-1 in coronary atherosclerosis of Japanese autopsies with diabetes mellitus: Hisayama study. *Atherosclerosis* 202: 573-81, 2009.

82. Sundaresan M, Yu ZX, Ferrans VJ, Irani K, Finkel T. Requirement for generation of H₂O₂ for platelet-derived growth factor signal transduction. *Science* 270: 296-9, 1995.
83. Taha H, Skrzypek K, Guevara I, Nigisch A, Mustafa S, Grochot-Przeczek A, Ferdek P, Was H, Kotlinowski J, Kozakowska M, Balcerczyk A, Muchova L, Vitek L, Weigel G, Dulak J, Jozkowicz A. Role of Heme Oxygenase-1 in Human Endothelial Cells. Lesson From the Promoter Allelic Variants. *Arterioscler Thromb Vasc Biol*: [Epub ahead of print], 2010.
84. Taille C, El-Benna J, Lanone S, Boczkowski J, Motterlini R. Mitochondrial respiratory chain and NAD(P)H oxidase are targets for the antiproliferative effect of carbon monoxide in human airway smooth muscle. *J Biol Chem* 280: 25350-60, 2005.
85. Thirunavukkarasu M, Juhasz B, Zhan L, Menon VP, Tosaki A, Otani H, Maulik N. VEGFR1 (Flt-1+/-) gene knockout leads to the disruption of VEGF-mediated signaling through the nitric oxide/heme oxygenase pathway in ischemic preconditioned myocardium. *Free Radic Biol Med* 42: 1487-95, 2007.
86. Tracz MJ, Juncos JP, Grande JP, Croatt AJ, Ackerman AW, Katusic ZS, Nath KA. Induction of heme oxygenase-1 is a beneficial response in a murine model of venous thrombosis. *Am J Pathol* 173: 1882-90, 2008.
87. True AL, Olive M, Boehm M, San H, Westrick RJ, Raghavachari N, Xu X, Lynn EG, Sack MN, Munson PJ, Gladwin MT, Nabel EG. Heme oxygenase-1 deficiency accelerates formation of arterial thrombosis through oxidative damage to the

- endothelium, which is rescued by inhaled carbon monoxide. *Circ Res* 101: 893-901, 2007.
88. Urbich C, Dimmeler S. Endothelial progenitor cells: characterization and role in vascular biology. *Circ Res* 95: 343-53, 2004.
89. Wang LJ, Lee TS, Lee FY, Pai RC, Chau LY. Expression of heme oxygenase-1 in atherosclerotic lesions. *Am J Pathol* 152: 711-20, 1998.
90. Wu BJ, Kathir K, Witting PK, Beck K, Choy K, Li C, Croft KD, Mori TA, Tanous D, Adams MR, Lau AK, Stocker R. Antioxidants protect from atherosclerosis by a heme oxygenase-1 pathway that is independent of free radical scavenging. *J Exp Med* 203: 1117-27, 2006.
91. Wu BJ, Midwinter RG, Cassano C, Beck K, Wang Y, Changsiri D, Gamble JR, Stocker R. Heme oxygenase-1 increases endothelial progenitor cells. *Arterioscler Thromb Vasc Biol* 29: 1537-42, 2009.
92. Yachie A, Niida Y, Wada T, Igarashi N, Kaneda H, Toma T, Ohta K, Kasahara Y, Koizumi S. Oxidative stress causes enhanced endothelial cell injury in human heme oxygenase-1 deficiency. *J Clin Invest* 103: 129-35, 1999.
93. Yet SF, Layne MD, Liu X, Chen YH, Ith B, Sibinga NE, Perrella MA. Absence of heme oxygenase-1 exacerbates atherosclerotic lesion formation and vascular remodeling. *FASEB J* 17: 1759-61, 2003.
94. Yet SF, Perrella MA, Layne MD, Hsieh CM, Maemura K, Kobzik L, Wiesel P, Christou H, Kourembanas S, Lee ME. Hypoxia induces severe right ventricular dilatation and infarction in heme oxygenase-1 null mice. *J Clin Invest* 103: R23-9, 1999.

95. Yet SF, Tian R, Layne MD, Wang ZY, Maemura K, Solovyeva M, Ith B, Melo LG, Zhang L, Ingwall JS, Dzau VJ, Lee ME, Perrella MA. Cardiac-specific expression of heme oxygenase-1 protects against ischemia and reperfusion injury in transgenic mice. *Circ Res* 89: 168-73, 2001.

Figure legends

FIG. 1. Absence of HO-1 exacerbates lipid accumulation and lesion formation in arteries.

HO-1^{+/+}apoE^{-/-} and HO-1^{-/-}apoE^{-/-} mice were fed Western diet starting at 4 weeks for a total of 8 weeks. The aortic arch and its branches were then harvested and analyzed. Photographs of representative arteries from HO-1^{+/+}apoE^{-/-} (A) or HO-1^{-/-}apoE^{-/-} (B) mice. Lesions are shown as white areas and indicated by arrows. C and D, Sudan IV staining (red) of lipid in the arteries of HO-1^{+/+}apoE^{-/-} (C) or HO-1^{-/-}apoE^{-/-} (D) mice. Br, brachiocephalic artery; LCC, left common carotid artery; LSC, left subclavian artery; RCC, right common carotid artery; RSC, right subclavian artery. Original magnification x15 (A through D). Reproduced from Yet et al. (93) with permission.

FIG. 2. Right ventricular dilatation and thrombus formation in HO-1^{-/-} mice in response to hypoxia.

Heart cross-sections (at the papillary muscle level) from wild-type (+/+) and HO-1 null (-/-) mice were stained with hematoxylin and eosin (n = 5–6 in each group). Mice were exposed to normoxia (a, +/+; b, -/-), 5 wk of hypoxia (c, +/+; d, -/-), or 7 wk of hypoxia (e, +/+; f, -/-). RV, right ventricle; LV, left ventricle; T, thrombus. Original magnification, 15×. Reproduced from Yet et al. (94) with permission.

FIG. 3. HO-1 dose-dependently improves postischemic cardiac performance in isolated perfused hearts.

Hearts from mice were stabilized for 30 min, then subjected

to 30 min of global ischemia followed by 40 min of reperfusion. Cardiac contractile performance is shown during ischemia and reperfusion. Wild-type (WT): open triangles, $n = 12$; transgenic TG.L: filled diamonds, $n = 12$; TG.M: open circles, $n = 5$; TG.H: filled squares, $n = 9$. **A**, Left ventricular end diastolic pressure (LVEDP). LVEDP at the end of ischemia was 65 ± 2 , 69 ± 2 , 66 ± 3 , and 63 ± 5 mmHg for WT, TG.L, TG.M, and TG.H, respectively. **B**, Left ventricular developed pressure (LVDevP). **C**, Rate pressure product (RPP). Error bars indicate standard errors. Reproduced from Yet et al. (95) with permission.


FIG. 4. HO-1 protects against myocardial infarction in transgenic mice. Myocardial infarcts from **(A)** wild-type (WT, $n = 6$) and **(B)** TG.H transgenic (TG, $n = 8$) mice were assessed by Evans blue and TTC staining after 1 h ischemia and 24 h reperfusion. The Evans blue perfused area, which is not at risk, stained blue; viable myocardium stained red and infarcted myocardium appeared pale. Representative wild-type and transgenic heart sections were shown and oriented anterior side up in **A** and **B**, respectively. Original magnification: x15. **C**, Myocardial infarcts are reduced in HO-1 transgenic mice. Wild-type (open bars, $n = 6$) and TG.H transgenic (filled bars, $n = 8$) mice were subjected to 1 h ischemia and 24 h reperfusion as in A. Risk Area/LV, percentage of left ventricle (LV) at risk; Infarct/Risk Area, infarcted area as percentage of risk area. Error bars indicate standard errors. * $P = 0.001$ vs. Infarct/Risk Area of WT animals.

Reproduced from Yet et al. (95) with permission.

FIG. 5. Ischemia/reperfusion injury induces left ventricular mural thrombi formation in diabetic HO-1^{-/-} mice. Heart cross sections at the papillary muscle level


after ischemia/reperfusion were stained with H&E. Sections from control (**A**) and diabetic (**B**) HO-1^{+/+} mouse hearts. (**C**) Ventricular section from control HO-1^{-/-} mice. (**D**) Thrombus formation in the left ventricle of HO-1^{-/-}-diabetic mouse heart. RV, right ventricle; LV, left ventricle; T, thrombus. Original magnification: x15.

FIG. 6. HO-1 in cardiovascular system. Various stresses lead to the production of reactive oxygen species in the cardiovascular system. In the absence of HO-1, perhaps through the excessive accumulation of reactive oxygen species, cardiovascular cells have a maladaptive response, resulting in cell death. In the presence of the buffering effect of HO-1, normal adaptive response provides protection to cardiovascular system. This implies that overexpression of HO-1 may play a protective role in cardiovascular diseases caused by reactive oxygen species.




Wu et al. Figure 1

Wu et al. Figure 2




Wu et al. Figure 3



Wu et al. Figure 4

Antioxidants & Redox Signaling
 A Central Role of Heme Oxygenase-1 in Cardiovascular Protection (doi: 10.1089/ars.2010.3726)
 This article has been peer-reviewed and accepted for publication, but has yet to undergo copyediting and proof correction. The final published version may differ from this proof.



Wu et al. Figure 5

Wu et al. Figure 6