

Novel Vilsmeier-type Methylenation for Synthesis of Dipyrzolylmethane Derivatives using Formamide or *N*-methylformamide

Fung Fuh Wong,^{1,*} Yu-Ying Huang¹

¹*Graduate Institute of Pharmaceutical Chemistry, China Medical University, No. 91 Hsueh-Shih Rd. Taichung, Taiwan 40402, R.O.C.*

*Corresponding author. Tel.: +886 4 2205 3366 ext. 5603; Fax: +886 4 2207 8083.

E-mail address: wongfungfuh@yahoo.com.tw, ffwong@mail.cmu.edu.tw (F. F. Wong).


Key words: Vilsmeier-type reaction, methylenation, dipyrzolylmethane, pyrazolone, formylation

Abstract

A novel Vilsmeier-type methylenation for synthesis of dipyrzolylmethanes was developed by reacting pyrazolones with formamide or *N*-methylformamide in the presence of phosphorous oxychloride POCl₃ coupling agent. This newly developed method can be efficiently provided a series of dipyrzolylmethane derivatives as the main products in excellent yields without the formylated products. Our experimental result was different with the classical Vilsmeier-type reaction.

Introduction

In 1927, the Vilsmeier or Vilsmeier–Haack reaction was first published and applied to an immense variety of substrates, from substituted benzenes to complex heterocycles.^{1,2} The classical Vilsmeier reagent, involves electrophilic substitution of an activated aromatic ring with a halomethyleniminium salt, which results from the reaction between an acid chloride (e.g., POCl₃, SOCl₂, (COCl)₂, and COCl₂) and an amide, usually DMF (see Scheme 1). When DMF was used in the reaction, the product of Vilsmeier reaction is an aldehyde. Thus, Vilsmeier reaction is often called as Vilsmeier formylation.³




Scheme 1

Vilsmeier reagent serves not only as a formylating agent,³ but also as an activating reagent for carboxylic acids to give esters,⁴ amides⁵ and acid chlorides,⁶ and for alcohols to give alkyl chlorides,⁷ esters,⁸ alkyl aryl sulfides⁹ and imides.¹⁰ Since, it is now used as a powerful organic synthetic tool for the formylation¹⁻³ or construction^{11,12} of many aromatic and heterocyclic compounds. Herein, we are a lot of interests in the study of reactive ability of various amides.¹³ Thus, we attempted to investigate the reactivity pyrazolones with various amides including formamide, *N*-methylformamide, *N,N*-dimethylformamide (DMF), and *N,N*-diethylformamide (DEF). Eventually, we have successfully provided the new Vilsmeier type methylenation to prepare dipyrazolylmethane derivatives by using formamide or *N*-methylformamide as reaction solvent, which are important key functional¹⁴ or bioactive materials.¹⁵

Result and Discussion

5-Pyrazolone derivatives were prepared as the starting materials by following the previous literature reported procedure via tandem condensation and thermal cyclization.¹⁶ In this newly developed method, we chose 1,3-diphenyl-1*H*-pyrazol-5-one (**2**) as the model for the selection of best conditions for the new Vilsmeier-type methylenation reaction (see Scheme 1). To search for optimum conditions and establish a reproducible procedure, compound **2** was allowed to react with formamide (3.0 equivalent) and phosphorous oxychloride (POCl₃, 5.0 equivalent) in CH₂Cl₂ solution at reflux (~ 50–60 °C) within 0.5–1 h. After aqueous worked-up and purification by column chromatography on silica gel, we isolated the corresponding methylenated product **3** in 96% yield without the formylated product **4** (see Scheme 2).



Scheme 2

To examine the solvent effects, we applied the same condition to 1,3-diphenyl-1*H*-pyrazol-5-one **2a** and various of amides containing *N*-methylformamide *N,N*-dimethylformamide (DMF), and *N,N*-diethylformamide (DEF) in the presence of POCl₃. Based on the experimental result, up to 86% yields of the formylated product **4** have been achieved when the pyrazolone **2a** was reacted with DMF and DEF (see Entry 3–4 of Table 1). When pyrazolone **2a** was used toward *N*-methylformamide within 0.5–1.0 h, we also found the methylenated product **3a** was obtained in 71% yield without the formylated product. Otherwise, the starting material **2a** was recovery in same condition (see entry 2 in Table 1).

Table 1. The amide agents study of new Vilsmeier methylenation via


1,3-diphenyl-1*H*-pyrazol-5-one (**2a**)

Entry	HCONR ³ R ⁴			Products		
	R ³	R ⁴	Dipyrazoylmethanes	Yields (%)	4-Formylpyrazolones	Yields (%)
1	H	H	3a	96	-	- ^a
2	Me	H	3a	71	-	- ^a
3	Me	Me	-	- ^b	4	86
4	Et	Et	-	- ^b	4	89

^aThe formulated product was not detectable.

^bThe methylenated product was not detectable.

Furthermore, we propose the plausible mechanism shown in Scheme 1 and Scheme 3, and accounts for the difference between the classical Vilsmeier formylation and the newly developed Vilsmeier methylenation. Firstly, amide agents were fast reacted with phosphorous oxychloride POCl₃ to form the reactive halomethyleniminium salts **1** in situ and side product HCl (see Scheme 1).¹⁷ In the presence of acidic species including excess amount of POCl₃ and HCl, pyrazolones **5** were performed isomerization to convert to pyrazoles **6**. Further, pyrazoles **6** were reacted with the reactive iminium species **1** to generate the imidination intermediate **8** and **9** through the reactive species **7** (see Scheme 3). For *N,N*-dimethylformamide (DMF) and *N,N*-diethylformamide (DEF) as the reactants, the Vilsmeier formylation was very smooth to generate the corresponding formylated product **10**.




Scheme 3

When formamide and *N*-methylformamide, were used in this procedure, the imidination intermediate **9** was effectively transferred to the more stable aminomethylidene derivatives **11** according to the formation of the intramolecule hydrogen-bonding. Consequently, the Michel addition of intermediate **11** with another equivalent of pyrazole **6** leads to dipyrazolyl-aminomethane **11**. After the protonation and hydrolysis two reactions were completed, the final product **13** was usefully obtained in excellent yield (see Scheme 3).

For the further investigation of the reliable procedure of the new developed Vilsmeier type methylenation, we also extended the pyrazolone scopes to 3-phenyl-pyrazolones **2b–2i** bearing *N*1-substituted aryl group with various *mono*-substituents, including *o*-, *m*-, *p*-ClPh, *o*-, *m*-, *p*-MePh, *p*-BrPh and *p*-OMePh with formamide. When employing the condition was shown in Scheme 4, all of substrates can be successfully converted to the corresponding methylenation dipyrazolylmethane products **3b–3i** in $\geq 84\%$ yields (see Table 2). More specifically,

1-(2,4-dinitrophenyl)-3-phenyl-1*H*-pyrazol-5-one (**2j**) own two strong withdrawing NO₂ groups in N1-aromatic ring, the corresponding dipyrazolymethanes **3j** did not detected in the same condition. 1-(2,4,6-Trichlorophenyl)-3-phenyl-1*H*-pyrazol-5-one (**2k**) having three chloro-substituents was also reacted with HCONH₂ and produced the corresponding dipyrazolymethanes **3k** in 87% yield (see Table 2). The dipyrazolymethanes **3a–3k** were reported and fully characterized by spectroscopic methods.^{14a,b,d} For example, compound **2a** presented a peak at δ 8.51 ppm for –C¹H=C in ¹H NMR. In ¹³C NMR spectrum, compound **2a** possessed characterization absorptions at δ 109.5 ppm for methylene carbon –¹³CH=C. The IR absorptions of **2a** showed peaks at 2109 cm⁻¹ for stretching of the –CH=C group and at 3291 cm⁻¹ for stretching of the –OH group.



Scheme 4

Table 2. The result of synthesis of dipyrazolymethanes **3a–3o**

Substrates	Pyrazolones		Dipyrazolymethanes	
	N1-R ¹	C3-R ²	Products	Yields (%)
2a	Ph	Ph	3a	96
2b	<i>o</i> -MePh	Ph	3b	90
2c	<i>o</i> -ClPh	Ph	3c	92
2d	<i>m</i> -MePh	Ph	3d	94
2e	<i>m</i> -ClPh	Ph	3e	94
2f	<i>p</i> -MePh	Ph	3f	96

2g	<i>p</i> -ClPh	Ph	3g	95
2h	<i>p</i> -BrPh	Ph	3h	93
2i	<i>p</i> -OMePh	Ph	3i	84
2j	2,4-di-NO ₂ Ph	Ph	3j	not detectable
2k	2,4,6-tri-ClPh	Ph	3k	87
2l	Ph	Me	3l	93
2m	Ph	CF ₃	3m	90
2n	Ph	<i>n</i> -Pr	3n	92
2o	Ph	<i>i</i> -Pr	3o	91

For the further investigation of the substituent effects, we extended this newly developed method toward 1-phenyl-3-substituted pyrazolones **2l–2o**, which contained Me, CF₃, *n*-Pr, or *i*-Pr groups at the C-3 position on pyrazolic ring. The corresponding dipyrazolymethane products **3l–3o** were obtained in 90–93% yields (see Table 2). Therefore, this efficient Vilsmeier-type methylenation method can also be successfully applied to synthesize a series of dipyrazolymethane derivatives **3a–3o**.

To search for the optimum condition for *N*-methylformamide as reaction solvent, we used 1,3-diphenyl-1*H*-pyrazol-5-one **2a** as mold and tried to prolonged the reaction time from 1.0 h to 2.0 h. Fortunately, the isolated yield of desired product **3a** was promoted from 71% to 95% (see Scheme 4 and Table 3). As a result, the optimum condition was extended 1,3-disubstituted-1*H*-pyrazol-5-ones **2b–2c**, **2h**, **2l**, and **2o** with *N*-methylformamide. The desired dipyrazolymethane products **3b–3c**, **3h**, **3l**, and **3o** were also obtained in good to excellent yields (82–94%, see Table 3). However, the formylated compounds were also not detectable using *N*-methylformamide as the reaction solvent.

Table 3. The results of the pyrazolones reacted with *N*-methylformamide in the

presence of POCl₃

Pyrazolones (2a–2c , 2h , 2l , and 2o)			Reaction Time (h)	Dipyrazolymethanes	
S.M.	N1-R ¹	C3-R ²		No.	Yields (%)
2a	Ph	Ph	~1.0	3a	71
2a	Ph	Ph	2.0	3a	95
2b	<i>o</i> -MePh	Ph	2.0	3b	90
2c	<i>p</i> -OMePh	Ph	2.0	3c	82
2h	<i>p</i> -BrPh	Ph	2.0	3h	91
2l	Ph	Me	2.0	3l	94
2o	Ph	<i>i</i> -Pr	2.0	3o	91

In conclusion, we have successfully developed the newly Vilsmeier-type methylenation method by treating various 1,3-disubstituted pyrazolones with formamide or *N*-methylformamide and POCl₃ coupling agent. Based on our experimental data, the result was different with the classical Vilsmeier-type reaction. And the yielding dipyrzolylmethane products without formylated compounds seemed determinate to the solvents involved formamide and *N*-methylformamide .

Acknowledgment

We are grateful to the China Medical University (CMU99-S-18) and the National Science Council of Republic of China for financial support (NSC-99-2320-B-039-014-MY3).

Experimental Section

General Procedure: All chemicals were reagent grade and used as purchased. All reactions were carried out under nitrogen atmosphere and monitored by TLC analysis. Flash column chromatography was carried out on silica gel (230–400 mesh). Commercially available reagents were used without further purification unless

otherwise noted. Dichloromethane, ethyl acetate, hexanes, and methanol were purchased from Mallinckrodt Chemical Co. The following compounds were purchased from Acoros Chemical Co: ethyl Butyrylacetate, ethyl acetoacetate, 2,4-dinitrophenyl hydrazine, formamide, phenylhydrazine, o-tolylhydrazine hydrochloride, m-tolylhydrazine hydrochloride, p-tolylhydrazine hydrochloride. 2,4,6-Trichlorophenyl hydrazine were purchased from TCI Chemical Co. 4-Bromophenylhydrazine hydrochloride, 2-chlorophenylhydrazine hydrochloride, 3-chlorophenylhydrazine hydrochloride, 4-chlorophenylhydrazine hydrochloride, ethyl benzoylacetate, ethyl isobutyrylacetate, ethyl trifluoroacetoacetate, 4-methoxyphenylhydrazine hydrochloride were purchased from Alfa Chemical Co. Phosphorylchloride were purchased from FERAK Chemical Co. Purification by gravity column chromatography was carried out by use of Merck Reagents Silica Gel 60 (particle size 0.063–0.200 mm, 70–230 mesh ASTM). Infrared (IR) spectra were measured on a Bomem Michelson Series FT-IR spectrometer. The wavenumbers reported are referenced to the polystyrene 1601 cm^{-1} absorption. Absorption intensities are recorded by the following abbreviations: s, strong; m, medium; w, weak. Proton NMR spectra were obtained on a Bruker (200 or 400 MHz) spectrometer by use of CDCl_3 as solvent. Carbon-13 NMR spectra were obtained on a Bruker (50 or 100 MHz) spectrometer by used of CDCl_3 as solvent. Carbon-13 chemical shifts are referenced to the center of the CDCl_3 triplet (δ 77.0 ppm). Multiplicities are recorded by the following abbreviations: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; *J*, coupling constant (Hz). Elemental analyses were carried out on a Heraeus CHN–O RAPID element analyzer.

Standard Procedure for the Synthesis of Dipyrzolylmethane Derivatives (3a-3i and 3k-3o)

The reliable procedure involved the treatment of pyrazolones (**3a-3i** and **3k-3o**, 1.0 equivalent) with catalytic amount of POCl₃ (~3 equivalent) in formamide solution (2 mL) at 50–60 °C within 0.5–1 h. When the reaction was completed, the reaction mixture was concentrated, added to saturate sodium bicarbonate (15 mL) and extracted with dichloromethane (15 mL). The organic extracts were dried over MgSO₄, filtered, and concentrated under reduced pressure. The residue solution was purified by column chromatography on silica gel to give the corresponding dipyrazolylmethane products (**3a-3i** and **3k-3o**) in 82–96% yields.

4,4'-Methylidenebis(1,3-diphenyl-2-pyrazolone) (3a): mp (recrystallized from ethanol) 261.2–262.2 °C; ¹H NMR (CDCl₃, 200 MHz) δ 7.24 (t, 1H, J=7.6, ArH), 7.44–7.53 (m, 5H, ArH), 7.82–7.86 (m, 2H, ArH), 8.01 (d, 2H, J=7.6 Hz, ArH); ¹³C NMR (50 MHz, CDCl₃) δ 39.7, 108, 119.1 (2 × CH), 109.5, 125.3, 126.0 (2 × CH), 128.9 (2 × CH), 130.7, 130.9 (2 × CH), 134.8, 138.1, 154.6, 170.2; IR (KBr) 3067 (m), 1715 (s, C=O), 1601 (s, C–N), 1496 (m), 1383 (m), 1326 (m), 1172 (m), 1116 (m), 899 (m) cm⁻¹ Anal. Calcd for C₃₁H₂₂N₄O₂; C: 77.16; H: 4.60; N: 11.61, Found: C: 77.13; H:4.62; N: 11.59.

4,4'-Methylidenebis[1-(2'-methylphenyl)-3-phenyl-2-pyrazolone] (3b): mp (recrystallized from ethanol) 205.3–206.3 °C; ¹H NMR (CD₃OD, 200 MHz) δ 2.27 (s, 3H, CH₃), 7.51 (d, 2H, J=7.2, ArH), 7.54–7.59 (m, 5H, ArH), 7.82 (d, 2H, J=7.2, ArH); ¹³C NMR (50 MHz, CD₃OD) δ 15.9, 126.3 (2 × CH), 126.6, 127.2, 128.0, 128.5, 129.2 (2 × CH), 131.1, 131.2, 131.5, 131.7, 136.7, 149.6, 157.4; IR (KBr) 3293 (m), 3124 (s), 1880 (m), 1537 (s, C–N), 1472 (m), 1305 (m), 1147 (m), 982 (m), 807 (m) cm⁻¹ Anal. Calcd for C₃₃H₂₆N₄O₂; C: 77.63; H: 5.13; N: 10.97, Found: C: 77.67; H:5.15; N: 10.96.

4,4'-Methylidenebis[1-(2'-chlorophenyl)-3-phenyl-2-pyrazolone] (3c): mp (recrystallized from ethanol) 198–199 °C; ¹H NMR (CD₃OD, 200 MHz) δ 7.51 (d, 2H,

$J=7.2$, ArH), 7.54–7.59 (m, 5H, ArH), 7.82 (d, 2H, $J=7.2$, ArH); ^{13}C NMR (50 MHz, CD_3OD) δ 126.3 (2 \times CH), 126.6, 127.2, 128.0, 128.5, 129.2 (2 \times CH), 131.1, 131.2, 131.5, 131.7, 136.7, 149.6, 157.4; IR (KBr) 3293 (m), 3124 (s), 1880 (m), 1537 (s, C–N), 1472 (m), 1305 (m), 1147 (m), 982 (m), 807 (m) cm^{-1} ; Anal. Calcd for $\text{C}_{31}\text{H}_{20}\text{Cl}_2\text{N}_4\text{O}_2$; C: 67.52; H: 3.66; N: 10.16, Found: C: 67.49; H:3.68; N: 10.13.

4,4'-Methylidenebis[1-(3'-methylphenyl)-3-phenyl-2-pyrazolone] (3d): mp (recrystallized from ethanol) 190–192 °C; ^1H NMR (CD_3OD , 200 MHz) δ 2.27 (s, 3H, CH_3), 7.51 (d, 2H, $J=7.2$, ArH), 7.54–7.59 (m, 5H, ArH), 7.82 (d, 2H, $J=7.2$, ArH); ^{13}C NMR (50 MHz, CD_3OD) δ 15.9, 126.3 (2 \times CH), 126.6, 127.2, 128.0, 128.5, 129.2 (2 \times CH), 131.1, 131.2, 131.5, 131.7, 136.7, 149.6, 157.4; IR (KBr) 3293 (m), 3124 (s), 1880 (m), 1537 (s, C–N), 1472 (m), 1305 (m), 1147 (m), 982 (m), 807 (m) cm^{-1} ; Anal. Calcd for $\text{C}_{33}\text{H}_{26}\text{N}_4\text{O}_2$; C: 77.63; H: 5.13; N: 10.97, Found: C: 77.61; H:5.09; N: 10.95.

4,4'-Methylidenebis[1-(3'-chlorophenyl)-3-phenyl-2-pyrazolone] (3e): mp (recrystallized from ethanol) 190–192 °C; ^1H NMR (CD_3OD , 200 MHz) δ 7.51 (d, 2H, $J=7.2$, ArH), 7.54–7.59 (m, 5H, ArH), 7.82 (d, 2H, $J=7.2$, ArH); ^{13}C NMR (50 MHz, CD_3OD) δ 26.3 (2 \times CH), 126.6, 127.2, 128.0, 128.5, 129.2 (2 \times CH), 131.1, 131.2, 131.5, 131.7, 136.7, 149.6, 157.4; IR (KBr) 3293 (m), 3124 (s), 1880 (m), 1537 (s, C–N), 1472 (m), 1305 (m), 1147 (m), 982 (m), 807 (m) cm^{-1} ; Anal. Calcd for $\text{C}_{31}\text{H}_{20}\text{Cl}_2\text{N}_4\text{O}_2$; C: 67.52; H: 3.66; N: 10.16, Found: C: 67.54; H:3.65; N: 10.18.

4,4'-Methylidenebis[1-(4'-methylphenyl)-3-phenyl-2-pyrazolone] (3f): mp (recrystallized from ethanol) 151–152 °C; ^1H NMR (CD_3OD , 200 MHz) δ 2.27 (s, 3H, CH_3), 7.40 (m, 2H, ArH); 7.48 (m, 2H, ArH); 7.78 (m, 2H, ArH); 7.99 (m, 2H, ArH).; ^{13}C NMR (50 MHz, CD_3OD) δ 39.6; 120.0; 126.0; 128.9; 129.0; 130.4; 130.7; 130.9; 154.9; 170.1.; IR (KBr) 3293 (m), 3124 (s), 1880 (m), 1537 (s, C–N), 1472 (m), 1305 (m), 1147 (m), 982 (m), 807 (m) cm^{-1} ; Anal. Calcd for $\text{C}_{33}\text{H}_{26}\text{N}_4\text{O}_2$; C: 77.63; H: 5.13;

N: 10.97, Found: C: 77.62; H:5.16; N: 10.94.

4,4'-Methylidenebis[1-(4'-chlorophenyl)-3-phenyl-2-pyrazolone] (3g): mp (recrystallized from ethanol) 163–164 °C; ¹H NMR (CD₃OD, 200 MHz) δ 7.40 (m, 2H, ArH); 7.48 (m, 2H, ArH); 7.78 (m, 2H, ArH); 7.99 (m, 2H, ArH).; ¹³C NMR (50 MHz, CD₃OD) δ 120.0; 126.0; 128.9; 129.0; 130.4; 130.7; 130.9; 154.9; 170.1.; IR (KBr) 3293 (m), 3124 (s), 1880 (m), 1537 (s, C–N), 1472 (m), 1305 (m), 1147 (m), 982 (m), 807 (m) cm⁻¹; Anal. Calcd for C₃₁H₂₀Cl₂N₄O₂; C: 67.52; H: 3.66; N: 10.16, Found: C: 67.56; H:3.62; N: 10.15

4,4'-Methylidenebis[1-(4'-bromophenyl)-3-phenyl-2-pyrazolone] (3h): mp (recrystallized from ethanol) 183–184 °C; ¹H NMR (CD₃OD, 200 MHz) δ 7.38 (m, 2H, ArH); 7.50 (m, 2H, ArH); 7.81 (m, 2H, ArH); 7.99 (m, 2H, ArH).; ¹³C NMR (50 MHz, CD₃OD) δ 120.0; 126.0; 128.9; 129.0; 130.4; 130.7; 130.9; 154.9; 170.1.; IR (KBr) 3293 (m), 3124 (s), 1880 (m), 1537 (s, C–N), 1472 (m), 1305 (m), 1147 (m), 982 (m), 807 (m) cm⁻¹; Anal. Calcd for C₃₁H₂₀Br₂N₄O₂; C: 55.15; H: 3.15; N: 8.75, Found: C: 55.17; H:3.19; N: 8.72.

4,4'-Methylidenebis[1-(4'-methoxyphenyl)-3-phenyl-2-pyrazolone] (3i): mp (recrystallized from ethanol) 128–130 °C; ¹H NMR (CDCl₃, 200 MHz) δ 3.79 (s, 3H, CH₃), 6.90 (d, 2H, J=6.8 Hz, ArH), 7.40–7.44 (m, 3H, ArH), 7.73 (d, 2H, J=7.2 Hz, ArH), 7.81 (d, 2H, J=6.8 Hz, ArH); ¹³C NMR (50 MHz, CDCl₃) d 39.5, 55.5, 114.0 (2 × CH), 121.0 (2 × CH), 125.9 (2 × CH), 128.9 (2 × CH), 130.6, 130.9, 131.5, 154.5, 157.2, 170.0; IR (KBr) 2927 (m), 1709 (s, C=O), 1512 (s, C–N), 1500 (m), 1248 (m), 1030 (m), 832 (m) cm⁻¹; Anal. Calcd for C₃₃H₂₆N₄O₄: C: 73.05; H: 4.83; N: 10.33. Found: C: 73.07; H: 4.85; N: 10.31.

4,4'-Methylidenebis[1-(2'-4'-6'-trichlorophenyl)-3-phenyl-2-pyrazolone] (3k): mp (recrystallized from ethanol) 144–146 °C; ¹H NMR (CD₃OD, 200 MHz) δ 7.34–7.43 (m, 3H, ArH), 7.68–7.72 (m, 4H, ArH); ¹³C NMR (50 MHz, CD₃OD) δ 47.1, 125.4 (2

× CH), 128.4 (2 × CH), 128.6 (2 × CH), 129.1 (2 × CH), 130.6, 132.1, 136.4, 152.8, 156.9, 171.0; IR (KBr) 3075 (m), 1732 (s, C=O), 1559 (s, C–N), 1461 (s), 1380 (m), 1157 (m), 1097 (m), 983 (m), 829 (m) cm⁻¹; Anal. Calcd for C₃₁H₁₆Cl₆N₄O₂: C: 54.02; H: 2.34; N: 8.13. Found: C: 54.03; H: 2.36; N: 8.11.

4,4'-Methylidenebis(3-methyl-1-phenyl-2-pyrazolone) (3l): mp (recrystallized from ethanol) 127–129 °C; ¹H NMR (CDCl₃, 200 MHz) δ 2.15 (s, 3H, CH₃), 7.14 (d, 1H, J = 2.2 Hz, ArH), 7.36 (dd, 2H, J = 7.8, 2.2 Hz, ArH), 7.81 (d, 2H, J = 7.8 Hz, ArH); ¹³C NMR (50 MHz, CDCl₃) δ 17.0, 43.1, 118.9 (2 × CH), 125.1, 128.8 (2 × CH), 137.9, 156.4, 170.6; IR (KBr) 3101 (s), 2802 (m), 1622 (s, C=O), 1584 (s, C–N), 1494 (m), 1412 (m), 1301 (s), 1220 (m), 1035 (m), 967 (m) cm⁻¹; Anal. Calcd for C₂₁H₁₈N₄O₂: C: 70.38; H: 5.06; N: 8.93. Found: C: 70.41; H: 5.09; N: 8.96.

4,4'-Methylidenebis(1-phenyl-3-trifluoromethyl-2-pyrazolone) (3m): mp (recrystallized from ethanol) 194–196 °C; ¹H NMR (CDCl₃, 200 MHz) δ 6.61 (s, 1H, ArH), 7.66 (d, 2H, J = 7.8 Hz, ArH), 7.80 (d, 2H, J = 7.6 Hz, ArH); ¹³C NMR (50 MHz, DMSO-d₆) δ 85.6, 122.3, 127.2, 129.1, 137.7, 140.4 (q, 3J_{C–F} = 37.6 Hz), 153.7. Anal. Calcd for C₂₁H₁₂F₆N₄O₂: C: 54.09; H: 2.59; N: 12.01. Found: C: 54.12; H: 2.57; N: 12.59.

4,4'-Methylidenebis(3-*n*-propyl-1-phenyl-2-pyrazolone) (3n): mp (recrystallized from ethanol) 201–202 °C; ¹H NMR (CDCl₃, 200 MHz) δ 1.28 (d, J = 7 Hz, 6H), 2.79 (m, 1H), 7.21 (t, J = 7.4 Hz, 1H), 7.41 (dd, J = 7.4, 8 Hz, 2H), 7.92 (d, J = 8 Hz, 2H); ¹³C NMR (50 MHz, CDCl₃) δ 20.5, 31.2, 40.3, 119.3, 125.4, 129.2, 138.7, 164.7, 171.1; IR (KBr) 3101 (s), 2802 (m), 1622 (s, C=O), 1584 (s, C–N), 1494 (m), 1412 (m), 1301 (s), 1220 (m), 1035 (m), 967 (m) cm⁻¹. Anal. Calcd for C₂₅H₂₆N₄O₂: C: 72.44; H: 6.32; N: 13.52. Found: C: 72.47; H: 6.36; N: 13.49.

4,4'-Methylidenebis(3-isopropyl-1-phenyl-2-pyrazolone) (3o): mp (recrystallized from ethanol) 101–103 °C; ¹H NMR (CDCl₃, 200 MHz) δ 1.21 (d, 6H, J = 7.4,

CH(CH₃)₂), 2.68–2.82 (m, 1H, CHMe₂), 7.14 (t, 1H, J=7.4, ArH), 7.35 (dd, 2H, J = 8.1, 7.4 Hz, ArH), 7.84 (d, 2H, J = 8.1 Hz, ArH); ¹³C NMR (50 MHz, CDCl₃) δ 20.1 (2 × CH₃), 30.7, 39.8, 118.8 (2 × CH), 124.9, 128.8 (2 × CH), 138.2, 164.3, 170.6; IR (KBr) 3101 (s), 2802 (m), 1622 (s, C=O), 1584 (s, C–N), 1494 (m), 1412 (m), 1301 (s), 1220 (m), 1035 (m), 967 (m) cm⁻¹. Anal. Calcd for C₂₅H₂₆N₄O₂: C: 72.44; H: 6.32; N: 13.52. Found: C: 72.41; H: 6.35; N: 13.53.

Reference

- (1) (a) Vilsmeier, A.; Haack, A. *Ber. Dtsch. Chem. Ges.* **1927**, *60*, 119–125; (b) Jones, G.; Stanforth, S. P. *Organic Reactions*; Paquette, L. A., Ed.; Wiley: New York, NY, 1997; Vol. 49, pp 1–330; (c) Downie, I. M.; Earle, M. J.; Heaney, H.; Shuhaibar, K. F. *Tetrahedron* **1993**, *49*, 4015–4021.
- (2) (a) Lilienkamp, A.; Johansson, M. P.; Wahala, K. *Org. Lett.* **2003**, *5*, 3387–3391; (b) Tang, X.-Y.; Shi, M. *J. Org. Chem.* **2008**, *73*, 8317–8322; (c) Perumal, P. T. *Indian J. Heterocycl. Chem.* **2001**, *11*, 1–8; (d) Pan, W.; Dong, D.-W.; Liu, Q. *Org. Lett.* **2007**, *9*, 2421–2426; (e) Wang, K.-W.; Xiang, D.-X.; Dong, D.-W. *Org. Lett.* **2008**, *10*, 1691–1697; (f) Jones, G.; Stanforth, S. P. *Org. React.* **2000**, *56*, 355–330; (g) Marson, C. M. *Tetrahedron* **1992**, *48*, 3659–3667; (h) Chen, L.; Zhao, Y.-L.; Liu, Q.; Cheng, C.; Piao, C.-R. *J. Org. Chem.* **2007**, *72*, 9259–9263; (i) Zhang, R.; Zhang, D.; Guo, Y.; Zhou, G.; Jiang, Z.; Dong, D. *J. Org. Chem.* **2008**, *73*, 9504–9511; (j) Wuts, P. G. M.; Northuis, J. M.; Kwan, T. A. *J. Org. Chem.* **2000**, *65*, 9223–9227; (k) Xiang, D.; Yang, Y.; Zhang, R.; Liang, Y.; Pan, W.; Huang, J.; Dong, D. D. *J. Org. Chem.* **2007**, *72*, 8593–8601; (l) Mahata, P. K.; Venkatesh, C.; Syam Kumar, U. K.; Ila, H.; Junjappa, H. *J. Org. Chem.* **2003**, *68*, 3966–3973.
- (3) (a) Campaigne, E.; Archer, W. L. *J. Am. Chem. Soc.* **1953**, *75*, 989–996; (b)

- Treihls, W.; Neupert, H. J.; Hiebsch *J. Chem. Ber.* **1959**, *92*, 141–145; (c) Damodiran, M.; Panneer, N.; Paramasivan, S.; Perumal, T. *Tetrahedron Lett.* **2009**, *50*, 5474–5478.
- (4) (a) Procopiou, P. A.; Brodie, A. C.; Deal, M. J.; Hayman, D. F. *Tetrahedron Lett.* **1993**, *34*, 7483–7486; (b) Procopiou, P. A.; Brodie, A. C.; Deal, M. J.; Hayman, D. F. *J. Chem. Soc., Perkin Trans. 1* **1996**, 2249–2253.
- (5) Zaoral, M.; Arnold, Z. *Tetrahedron Lett.* **1960**, *1*, 9–13.
- (6) Eilingsfeld, H.; Seefelder, M.; Weidinger, H. *Angew. Chem.* **1960**, *72*, 836–841.
- (7) Hepburn, D. R.; Hudson, H. R. *J. Chem. Soc., Perkin Trans. 1* **1976**, 754–762.
- (8) (a) Barrett, A. G. M.; Braddock, D. C.; James, R. A.; Koike, N.; Procopiou, P. A. *J. Org. Chem.* **1998**, *63*, 6273–6279; (b) Barrett, A. G. M.; Koike, N.; Procopiou, P. A. *Chem. Commun.* **1995**, 1403–1412.
- (9) Kawano, Y.; Kaneko, N.; Mukaiyama, T. *Chem. Lett.* **2005**, *34*, 1612–1618.
- (10) Barrett, A. G. M.; Braddock, D. C.; James, R. A.; Procopiou, A. *Chem. Commun.* **1997**, 433–438.
- (11) (a) Savitha, G.; Perumal, P. T. *Tetrahedron Lett.* **2007**, *48*, 2943–2947; (b) Damodiran, M.; Kumar, R. S.; Sivakumar, P. M.; Doble, M.; Perumal, P. T. *J. Chem. Sci.* **2009**, *121*, 65–72; (c) Karthikeyan, K.; Perumal, P. T.; Etti, S.; Shanmugam, G. *Tetrahedron* **2007**, *63*, 10581–10586; (d) Selvam, N. P.; Perumal, P. T. *Tetrahedron* **2008**, *64*, 2972–2978; (e) Selvam, N. P.; Saranya, S.; Perumal, P. T. *Can. J. Chem.* **2008**, *86*, 32–36; (f) Shanthi, G.; Perumal, P. T. *Tetrahedron Lett.* **2007**, *48*, 6785–6791; (g) Selvam, N. P.; Shanthi, G.; Perumal, P. T. *Can. J. Chem.* **2007**, *84*, 989–995; (h) Thirumurugan, P.; Muralidharan, D.; Perumal, P. T. *Dyes Pigments* **2009**, *81*, 245–251; (i) Thirumurugan, P.; Perumal, P. T. *Tetrahedron Lett.* **2009**, *50*, 4145–4152; (j) Damodiran, M.; Muralidharan, D.; Perumal, P. T. *Bioorg. Med. Chem. Lett.* **2009**, *19*,

3611–3617.

- (12) (a) Sivaprasad, S.; Sridhar, R.; Perumal, P. T. *J. Heterocycl. Chem.* **2006**, *43*, 389–330; (b) Kumar, K. H.; Perumal, P. T. *Chem. Lett.* **2005**, *34*, 1346–1352; (c) Kumar, H. K.; Selvi, S.; Perumal, P. T. *J. Chem. Res.* **2004**, 218–226; (d) Sridhar, R.; Perumal, P. T. *Synth. Commun.* **2003**, *33*, 607. 1483–1496; (e) Selvi, S.; Perumal, P. T. *Indian J. Chem., Sect. B* **2002**, *41B*, 1887–1893; (f) Selvi, S.; Perumal, P. T. *Synth. Commun.* **2001**, *31*, 2199–2208; (g) Amaresh, R. R.; Perumal, P. T. *Synth. Commun.* **2000**, *30*, 2269; (h) Selvi, S.; Perumal, P. T. *Synth. Commun.* **2000**, *30*, 3925–3936; (i) Selvi, S.; Perumal, P. T. *Indian J. Chem., Sect. B* **2000**, *39B*, 163–169; (j) Amaresh, R. R.; Perumal, P. T. *Tetrahedron* **1999**, *55*, 8083–8090; (k) Perumal, P. T.; Majo, V. J.; Anand, R. V. *Indian J. Chem., Sect. B* **1999**, *38*, 763–772; (l) Amaresh, R. R.; Perumal, P. T. *Tetrahedron* **1998**, *54*, 14327–14333; (m) Amaresh, R. R.; Perumal, P. T. *Tetrahedron Lett.* **1998**, *39*, 3837–3842; (n) Majo, V. J.; Perumal, P. T. *J. Org. Chem.* **1998**, *63*, 7136–330; (o) Selvi, S.; Perumal, P. T. *Tetrahedron Lett.* **1997**, *38*, 6263–6267; (p) Majo, V. J.; Perumal, P. T. *Tetrahedron Lett.* **1997**, *38*, 6889–6992; (q) Majo, V. J.; Perumal, P. T. *Tetrahedron Lett.* **1996**, *37*, 5015–5019; (r) Balasundaram, B.; Venugopal, M.; Perumal, P. T. *Tetrahedron Lett.* **1993**, *34*, 4249–4256.
- (13) Yilmaz, V. T.; Topcu, Y. *Thermochimica Acta.* **1997**, *307*, 143–149.
- (14) (a) Azev, Y. A.; Shorshnev, S. V.; Golomolzin, B. V. *Tetrahedron Lett.* **2009**, *50*, 2899–2903; (b) Azev, Y. A.; Shorshnev, S. V. *Chem. Heterocycl. Comp.* **2009**, *45*, 115–116; (c) Wang, J.; Cao, W.-F.; Sua, J.-H.; Tian, H.; Huang, Y.-H.; Sun, Z.-R. *Dyes Pigments* **2003**, *57*, 171–179; (d) Shermolovich, Y. G.; Yemets, S. V. *J. Fluor. Chem.* **2000**, *101*, 111–116; (e) Azev, Y.A.; Neunhoeffler, H.; Foro, S.; Lindner, H.J.; Shorshnev, S.V. *Mendeleev Commun.*

- 1995**, 5, 229–231.
- (15) Radi, M.; Bernardo, V.; Bechi, B.; Castagnolo, D.; Pagano, M.; Botta, M. *Tetrahedron Lett.* **2009**, 50, 6572–6575.
- (16) (a) Wang, X.-j.; Tan, J.; Grozinger, K. *Tetrahedron Lett.* **2000**, 41, 4713–4716.
(b) DeRuiter, J.; Carter, D. A.; Arledge, W. S.; Sullivan, P. J. *J. Heterocycl. Chem.* **1987**, 24, 149–153. (c) Braña, M. F.; Gradillas, A.; Ovalles, A. G.; López, B.; Acero, N.; Llinares, F. Mingarro, D. M. *Bioorg. Med. Chem.* **2006**, 14, 9–16.
- (17) (a) Srivastava, V.; Negi, A. S.; Kumar, J. K.; Gupta, M. M. *Steroids* **2006**, 71, 632–638; (b) El-Shishtawy, R.M.; Almeida, P. *Tetrahedron* **2006**, 62, 7793–7798.

Fung Fuh Wong,* Yu-Ying Huang

Novel Vilsmeier-type Reaction for Synthesis of Dipyrzolylmethane

Derivatives using Formamide or *N*-methylformamide

