

Case Report

Pediatric Suprarenal Paraganglioma Presented as Blurred Vision — A Case Report

Chia-Hung Liu¹, Chia-Ying Lee², Chang-Ching Wei², Chao-Hsiang Chang³,
and Ching-Yuang Lin²

¹Department of Pediatrics, Taichung Armed Forces General Hospital

²Department of Pediatrics

³Department of Urology, Chinese Medical University Hospital
Taichung, Taiwan, Republic of China

Abstract

Ocular manifestations of hypertensive retinopathy such as vision impairment or even loss associated with pheochromocytoma or paraganglioma were rarely reported. In this paper, we report a case of a fifteen-year-old girl with bilateral blurred vision for two months. Initial blood pressure (BP) was 218/159 mm Hg. Ophthalmoscopy showed bilateral hypertensive retinopathy with optic disc edema. Two-dimensional echocardiography revealed hypertrophy of the interventricular septum and left ventricular wall. Initial emergent hypertension was stabilized with phenoxybenzamine and a calcium channel blocker. Renal ultrasound and abdominal computed tomography revealed a huge right suprarenal tumor compressing renal hilar vessels. Urinary vanillylmandelic acid (VMA) level was 56.5 mg/24 hr (normal, 2-7 mg/24 hr). Retroperitoneal suprarenal paraganglioma was highly suspected. Tumor resection was performed after blood pressure was stabilized for two weeks. Pathological study revealed a paraganglioma of 10 × 6.7 × 4 cm in size with suspicion of metastasis. Postoperative 18-fluorodeoxyglucose positron emission tomography scan revealed no metastasis. Her visual acuity, BP and VMA level returned to normal after tumor resection. While pediatric pheochromocytoma and paraganglioma are rare, blurred vision with hypertensive retinopathy may be induced by pheochromocytoma/paraganglioma in children. It is worthwhile reporting our experience with a patient who presented with initial blurred vision caused by paraganglioma. (Acta Nephrologica 2011; 25: 196-200)

KEY WORDS: paraganglioma, hypertensive retinopathy, vanillylmandelic acid

Introduction

Pheochromocytomas in children, arising from chromaffin cells of adrenal medullary or extra-adrenal paraganglionic tissue, are rare. Tumors from extra-adrenal chromaffin tissues are referred to as paragangliomas or extra-adrenal pheochromocytomas. Incidence of pheochromocytoma and paraganglioma is 0.40-2.06 per million per year (1, 2). Most cases are found in adults with incidence in children lower than 5% (3). Most symptoms are due to elevated levels of

catecholamines. Sustained hypertension in pediatric pheochromocytoma cases is more common than in adult cases.

Pheochromocytoma is estimated to be prevalent in approximately 1% of hypertensive pediatric patients and should be considered after exclusion of more common causes (4, 5). Paroxysmal hypertension or development of orthostatic hypotension against the background of sustained hypertension or amelioration or even inversion of circadian blood pressure (BP) rhythm may be crucial to diagnosing pheochromocytoma.

Corresponding author: Ching-Yuang Lin, MD, PhD, College of Medicine, China Medical University, Division of Pediatric Nephrology, China Medical University Hospital, Clinical Immunological Center, China Medical University Hospital, No. 2, Yuh-Der Rd., Taichung 40402, Taiwan, R.O.C. Tel: +886-4-2207-1501, Fax: +886-4-2207 ext. 1352, E-mail: cylin@mail.cmuh.org.tw
Received: September 17, 2010; Revised: October 22, 2010; Accepted: December 3, 2010.

toma (6). Palpitation, headache, excessive sweating and pallor are well-known symptoms of continuous or paroxysmal catecholamine excess and should arouse suspicion of pheochromocytoma (7). Sweating, nausea, vomiting, weight loss, polyuria and visual disturbances are reported more often in children than in adults (8, 9). While pediatric pheochromocytoma and paraganglioma are rare, they provide an important differential diagnosis for childhood hypertension. Ocular manifestations of hypertensive retinopathy such as vision impairment or loss associated with pheochromocytoma were reported.

Case Report

A fifteen-year-old girl visited an ophthalmologic clinic because of a two-month history of blurred vision. Excessive sweating had also been noted for three years. Her medical history was unremarkable with no family history of hypertension. Hypertensive retinopathy was suspected at ophthalmologic clinic and she was admitted for further observation. Her body weight was 51.8 kg (50th percentile) and body height was 158 cm (50-75th percentile). She had best corrected visual acuity of 6/60 in both eyes, which was worse than visual acuity of 6/6 in both eyes before this hypertensive episode. Results of slit lamp examination and intraocular pressure were normal. Ophthalmoscopic examination showed papilledema over the optic disc with hemorrhage and cotton-wool spots at the retina in each eye. Heart rate was 122 beats/minute and respiration rate was 20/minute. Wide fluctuations in BP were noted: maximal of 218/159 mmHg and minimal of 150/110 mmHg. Auscultation of heart sounds revealed grade 2 short systolic murmurs over left middle sternal border and no palpable abdominal mass. Urine analysis was unremarkable. Complete cell count, blood urea nitrogen, creatinine, and electrolyte levels of blood were all within normal limits. Urinary vanillylmandelic acid (VMA) level was 56.5 mg/24 hr (normal, 2-7 mg/24 hr) and plasma renin activity was 9.62 ng/mL/hr during supine rest (normal, 0.15-2.33 ng/mL/hr). Chest roentgenography and electrocardiography also showed normal results. Two-dimensional echocardiography revealed hypertrophy of interventricular septum and left ventricular free wall. Ultrasound revealed a right, huge suprarenal mass with compression of right kidney. Computed tomography of the abdomen demonstrated a right retroperitoneal, suprarenal tumor of 7.2 × 5.5 cm in size with tortuous veins and displacement of right renal artery and inferior vena cava (Fig. 1). Captopril renoscintigraphy revealed no probability of renovascular hypertension. Before operation, right retroperitoneal, suprarenal paraganglioma was suspected. Her thyroid function, calcitonin and intact PTH level were within normal



Fig. 1. Computed tomography of the abdomen demonstrating a 7.2 × 5.5-cm hypervascular mass lesion (arrow) over right retroperitoneum, suprarenal region with tortuous veins and displacement of right renal artery and inferior vena cava.

limits. The patient was treated with phenoxybenzamine, an α -adrenergic blocking agent, and a calcium channel blocker. Initial phenoxybenzamine dosage was 30 mg/day but increased to 120 mg/day after 4 weeks of therapy to control systolic BP within the range of 130 to 100 mm Hg. Surgery was arranged after hypertension was controlled. Tumor resection was performed after BP was stabilized for two weeks. A 10 × 6.7 × 4-cm tumor located between the aorta and the inferior vena cava with abundant collateral circulation and dense adhesion was noted intraoperatively. Microscopically, the tumor was encapsulated and tumor cells within the mass (tumor = mass) were characterized by round and focal pleomorphic nuclei, indistinct to small nucleoli, and pink cytoplasm (Fig. 2a). Tumor cells were arranged in nest or Zellballen pattern (Fig. 2b) and surrounded by thin fibrovascular channels. Both vascular and capsular invasion with patchy hemorrhage and confluent necrosis were noted. Nerve bundles both within and at the periphery of the tumor were also noted. Several big vessels within the tumor and vascular wall were focally invaded by tumor cells. There was no adrenal component in this specimen. Immunohistochemical study showed tumor cells with positive stain for synaptophysin (Fig. 2c) and chromogranin (Fig. 2d). Sustentacular frameworks were highlighted by S-100 (Fig. 2e). All resection margins were free, but tumor nests were very close to resection margins focally (less than 1 mm). Possibility of malignancy and metastasis was highly suspected,




Fig. 2. Hematoxylin and eosin stain demonstrating tumor cells characterized by round and focal pleomorphic nuclei (2a) and arranged in nest or Zellballen pattern (2b). Immunohistochemical stain showed tumor cells with positive stain for synaptophysin (2c) and chromogranin (2d). Sustentacular frameworks highlighted by S-100 (2e).

with tumor necrosis, capsular invasion, vascular invasion and large vessels within the tumor having invaded vascular wall observed. Ki-67 positive cells

were about 4-5%. The diagnosis of paraganglioma with possibility of metastasis was made (A person not a disease is diagnosed). Postoperative 18-fluorodeox-

glucose positron emission tomography scan revealed no metastasis. She had complete resolution of the symptoms with normalized blood pressure 2 months postoperatively without anti-hypertensive medication. Visual acuity and VMA level returned normal after tumor resection and were regularly monitored at our outpatient clinic for 6 months.

Discussion

Pheochromocytomas, originating from chromaffin cells of adrenal medullary or extra-adrenal paraganglionic tissue, are rare in children. Those that arise from the autonomic nervous system of abdomen, thorax, neck or head are more specifically termed as paragangliomas and often referred to as extra-adrenal pheochromocytomas. While the incidence of pheochromocytoma and paraganglioma in childhood is low, they represent the most common pediatric endocrine tumor and constitute a key cause of severe hypertension in children. Clinical presentations of childhood pheochromocytoma or paraganglioma are highly variable, with tumors found in some patients who are completely asymptomatic. The average age at onset is 11 years, with a male/female ratio of 2:1 (5, 10). Hypertension caused by secretion of one or more catecholamine hormones is the most prominent feature and may be paroxysmal or persistent. Compared with adults, children more often present with symptoms related to hypertension: headache, tachycardia, diaphoresis, postural hypotension, nausea, vomiting, abdominal or chest pain, weight loss, shortness of breath, polydipsia, polyuria, acrocyanosis, convulsions, and visual disturbances (11). Yet symptoms and signs of pheochromocytoma or paraganglioma are not so specific in children, thus making early diagnosis difficult. Clinical alertness and BP measurement are vital for detecting child hypertension. Sustained hypertension is found in 60-90% of pheochromocytoma in pediatric cases, versus 50% in adult cases. Pheochromocytoma or paraganglioma should be considered after excluding common causes of pediatric hypertension such as cardiovascular or renovascular hypertension, renal parenchymal disease and renal congenital anomaly. Other endocrine causes of pediatric hypertension include thyroid disease, Cushing syndrome and congenital adrenal hyperplasia. Evaluation of suspected pheochromocytoma or paraganglioma should include 24-hour norepinephrine, epinephrine, total metanephrines, and VMA concentrations in urine. In more than 95% of patients of pheochromocytoma or paraganglioma, diagnosis can be established by a finding of an urinary VMA concentration (12). Plasma catecholamine determination is generally less useful than urinary assessment, except in hypertension crisis. Upon diagnosis, surgical

excision is the only curative treatment. Surgical complications may emanate from excessive and abrupt release of catecholamines (12). Adequate preoperative preparation and treatment with non-competitive α -adrenoreceptor blocker like phenoxybenzamine can treat adrenergic manifestations and hypertensive crisis during anesthetic induction or surgical manipulation of tumors and reduce morbidity and mortality effectively. Patients with adequate phenoxybenzamine are prone to orthostatic hypotension (13). This preoperative treatment should be given in the hospital; patients should be instructed on the need for and assisted in maintaining upright position to prevent complications. Hypertensive retinopathy is uncommon but a critical condition for children. Delay diagnosis may result in permanent visual damage owing to optic atrophy. In its early stages, fundus picture shows retinal arteriolar constriction. As the disease progresses, superficial hemorrhages and small white superficial foci of retinal ischemia in nerve fiber layer (cotton-wool spots) develop. In cases of severe hypertension, optic discs become congested and edematous. If elevated BP is controlled promptly with medication or by removal of the tumor, retinal blood vessels may recover without permanent pathologic changes (14). In our case, fundus ophthalmoscopy revealed intraretinal hemorrhage, cotton-wool spots and optic disc edema; visual acuity returned dramatically to normal after tumor resection, indicating that early correction may reverse hypertensive retinopathy in a child.

Though pediatric pheochromocytoma and paraganglioma are rare, blurred vision with hypertensive retinopathy may arise from either of them. Permanent visual damage can be severe if not treated early. Quantification of 24-hour urine VMA, plasma catecholamine and localized image study can aid diagnosis. Preoperative stabilization of BP and surgical intervention can result in uneventful recovery.

In summary, pheochromocytomas or paraganglioma have varied clinical ocular manifestations of blurred vision and hypertensive retinopathy. While pediatric pheochromocytoma and paraganglioma are rare, they constitute a key diagnosis for child hypertension.

References

1. Fernandez-Calvet L, Garcia-Mayor RV. Incidence of pheochromocytoma in South Galicia, Spain. *J Intern Med* 236: 675-677, 1994.
2. DeGraeff J, Horak BJ. The incidence of pheochromocytoma in the Netherlands. *Acta Med Scand* 176: 583-593, 1964.
3. Abramson SJ. Adrenal neoplasms in children. *Radiol Clin North Am* 35: 1415-1453, 1997.
4. Ein SH, Shandling B, Wesson D, Filler R. Recurrent pheochromocytomas in children. *J Pediatr Surg* 25: 1063-1065, 1990.
5. Ross JH. Pheochromocytoma. Special considerations in children. *Urol Clin North Am* 27: 393-402, 2000.

6. Zelinka T, Pacak K, Widimsky J Jr. Characteristics of blood pressure in pheochromocytoma. *Ann NY Acad Sci* 1073: 86-93, 2006.
7. Lenders JW, Eisenhofer G, Mannelli M, Pacak K. Pheochromocytoma. *Lancet* 366: 665-675, 2005.
8. Ciftci AO, Tanyel FC, Senocak ME, Buyukpamukcu N. Pheochromocytoma in children. *J Pediatr Surg* 36: 447-452, 2001.
9. Hack HA. The perioperative management of children with pheochromocytoma. *Paediatr Anaesth* 10: 463-476, 2000.
10. Beltsevich DG, Kuznetsov NS, Kazaryan AM, Lysenko MA. Pheochromocytoma surgery: epidemiologic peculiarities in children. *World J Surg* 28: 592-596, 2004.
11. Werbel SS, Ober KP. Pheochromocytoma: update on diagnosis, localization, and management. *Med Clin North Am* 79: 131-153, 1995.
12. Barvo EL. Evolving concepts in pathophysiology, diagnosis, and treatment of pheochromocytoma. *Endocr Rev* 15: 356-368, 1994.
13. Russell WJ, Metcalfe IR, Tonkin AL, Frewin DB. The preoperative management of pheochromocytoma. *Anaesth Intensive Care* 26: 196-200, 1998.
14. Tso MO, Jampol LM. Pathophysiology of hypertensive retinopathy. *Ophthalmology* 89: 1132-1145, 1982.