



Contents lists available at ScienceDirect

Free Radical Biology & Medicine

journal homepage: www.elsevier.com/locate/freeradbiomed

Original Contribution

Viscolin reduces VCAM-1 expression in TNF- α -treated endothelial cells via the JNK/NF- κ B and ROS pathwayChan-Jung Liang^{a,1}, Shu-Huei Wang^{a,1}, Yung-Hsiang Chen^b, Shih-Sheng Chang^c, Tong-Long Hwang^d, Yann-Lii Leu^d, Ying-Chih Tseng^e, Chi-Yuan Li^{c,*}, Yuh-Lien Chen^{a,**}^a Department of Anatomy and Cell Biology, College of Medicine, National Taiwan University, Taipei, Taiwan, Republic of China^b Graduate Institute of Integrated Medicine, China Medical University, Taichung, Taiwan, Republic of China^c Graduate Institute of Clinical Medical Sciences, China Medical University, Taichung, Taiwan, Republic of China^d Graduate Institute of Natural Products, College of Medicine, Chang Gung University, Taoyuan, Taiwan, Republic of China^e Department of Obstetrics and Gynecology, Hsinchu Cathay General Hospital, Hsinchu, Taiwan, Republic of China

ARTICLE INFO

Article history:

Received 3 January 2011

Revised 10 June 2011

Accepted 18 June 2011

Available online xxx

Keywords:

Viscolin

Adhesion molecules

Reactive oxygen species

Inflammation

Mitogen-activated protein kinases

Free radicals

ABSTRACT

Viscolin, a major active component in a chloroform extract of *Viscum coloratum*, has antioxidative and anti-inflammatory properties. We focused on its effects on the expression of vascular cell adhesion molecule-1 (VCAM-1) in tumor necrosis factor- α (TNF- α)-treated human umbilical vein endothelial cells (HUVECs). The TNF- α -induced expression of VCAM-1 was significantly reduced by respectively 38 ± 7 or $34 \pm 16\%$ when HUVECs were pretreated with 10 or 30 μ M viscolin, as shown by Western blotting, and was also significantly reduced by pretreatment with the antioxidants *N*-acetylcysteine, diphenylene iodonium chloride, and apocynin. Viscolin also reduced TNF- α -induced VCAM-1 mRNA expression and promoter activity, decreased reactive oxygen species (ROS) production and nicotinamide adenine dinucleotide phosphate oxidase activity, and significantly reduced the binding of monocytes to TNF- α -stimulated HUVECs. The attenuation of TNF- α -induced VCAM-1 expression and cell adhesion was partly mediated by a decrease in JNK phosphorylation. Furthermore, viscolin reduced VCAM-1 expression in the aorta of TNF- α -treated mice in vivo. Taken together, these data show that viscolin inhibits TNF- α -induced JNK phosphorylation, nuclear translocation of NF- κ B p65, and ROS generation and thereby suppresses VCAM-1 expression, resulting in reduced adhesion of leukocytes. These results also suggest that viscolin may prevent the development of atherosclerosis and inflammatory responses.

© 2011 Elsevier Inc. All rights reserved.

Vascular inflammation is a critical risk factor in the initiation and development of cardiovascular disorders, such as atherosclerosis [1]. Activation of the endothelium at the inflammatory site results in leukocyte adhesion to the endothelium and subsequent transmigration of leukocytes into the subendothelial space, an early event in atherosclerosis [2]. The leukocyte adhesion is primarily mediated by adhesion molecules expressed on the surface of the endothelium. Previous studies have shown that vascular cell adhesion molecule 1 (VCAM-1), but not intercellular cell adhesion molecule-1, mediates leukocyte recruitment to early lesions of atherosclerosis and seems to be the dominant adhesion molecule on the endothelial surface of the vascular wall in the initiation of atherosclerosis [3,4]. In addition, levels of VCAM-1 expression have been suggested to be closely associated with the generation of reactive oxygen species (ROS) [5]. Thus, inhibition of VCAM-1 expression and ROS generation might be a

useful therapeutic strategy for treating vascular inflammation and cardiovascular diseases [6].

Herbal medicines have been widely used in Asian countries for many centuries and the active components of herbs and their structural backbones might provide a useful platform for the development of effective pharmacological agents. *Viscum coloratum* Nakai, a traditional Chinese herbal medicine, has been used for a long time to treat inflammatory diseases, such as rheumatism and atherosclerosis. However, its active components and pharmacological effects have not been extensively studied. *V. coloratum* inhibits superoxide anion generation by human neutrophils [7] and viscolin (4',4''-dihydroxy-2',3',6',3''-tetramethoxy-1,3-diphenylpropane), a new chalcone from *V. coloratum*, inhibits human neutrophil superoxide anion and elastase release [8]. Although previous studies have shown that viscolin has antioxidative and anti-inflammatory effects on human neutrophils [7-9], it is unclear whether it has anti-inflammatory and antioxidative effects on human vascular endothelial cells. Furthermore, the regulation of adhesion molecule expression involves a complex array of intracellular signaling pathways including mitogen-activated protein kinases (MAPKs), transcriptional factors, and ROS [10-13]. Although these multiple signaling molecules have

* Corresponding author.

** Corresponding author. Fax: +886 2 33931713.

E-mail addresses: cyl@ndmctsgh.edu.tw (C.-Y. Li), ylichen@ntu.edu.tw (Y.-L. Chen).¹ These authors contributed equally to this work.

received considerable attention [10–13], little is known about the effects of viscolin on adhesion molecule expression and the mechanisms of these effects, and a better understanding of this might provide important insights into the prevention of atherogenesis and inflammation. We therefore tested the ability of viscolin to modulate the expression of adhesion molecules, MAPKs, and transcriptional factors in tumor necrosis factor- α (TNF- α)-treated human umbilical vein endothelial cells (HUVECs). In addition, we examined its effects on VCAM-1 expression in TNF- α -treated mice. Our results showed that viscolin reduced VCAM-1 expression both in vitro and in vivo and that this effect is partly mediated by inhibition of JNK phosphorylation, NF- κ B activation, and ROS generation. Viscolin also significantly inhibited the adhesion of the human monocytic cell line U937 to TNF- α -treated HUVECs.

Materials and methods

Materials

Polyclonal rabbit IgG against human VCAM-1, GAPDH, phospho-p38, or histone (H1) and horseradish peroxidase (HRP)-conjugated goat anti-mouse IgG or anti-rabbit IgG antibodies were purchased from Santa Cruz Biotechnology (Santa Cruz, CA, USA). Polyclonal rabbit IgG against human phospho-ERK1/2 or phospho-JNK was purchased from Cell Signaling (Beverly, MA, USA). Monoclonal rabbit against human p65 and phospho-p65 antibodies were purchased from GeneTex (Irvine, CA, USA). Diphenylene iodonium chloride (DPI), PD98058, SP600125, and SB203580 were purchased from Biomol (Plymouth Meeting, PA, USA). Apocynin (APO) was purchased from ChromaDex (Irvine, CA, USA). The Amplex red hydrogen peroxide/peroxidase assay kit and Trizol reagent were purchased from Invitrogen (Carlsbad, CA, USA). Recombinant human TNF- α was purchased from PeproTech (Rocky Hill, NJ, USA). *N*-acetylcysteine (NAC) was purchased from Sigma-Aldrich (St. Louis, MO, USA). Polyvinylidene difluoride (PVDF) membranes were purchased from Millipore (Billerica, MA, USA).

Extraction and purification of viscolin

Viscolin was purified as described previously [7,8]. In brief, dried stems of *V. coloratum* Nakai (family Loranthaceae) (471 g) were extracted six times with 1 L of methanol and the combined extracts evaporated and partitioned to yield chloroform and aqueous extracts. The chloroform extract (10.5 g) was subjected to column chromatography on a silica gel column and eluted with chloroform and methanol step gradients to obtain the active extract, PPE-SVC (CHCl₃:MeOH 9:1) (4.7 g). PPE-SVC was rechromatographed on a silica gel column and eluted with a gradient of *n*-hexane and acetone to give a chalcone derivative, viscolin (Fig. 1, 53.6 mg).

Cell culture

Primary cultures of HUVECs were prepared as previously described [14]. The cells were grown in medium 199 containing penicillin-streptomycin (1%), endothelial cell growth supplement (30 μ g/ml),

and fetal bovine serum (10%) at 37 °C in a humidified atmosphere of 95% air, 5% CO₂ and were used between passages 2 and 5.

RNA extraction and reverse transcriptase-polymerase chain reaction (RT-PCR)

Total RNA was extracted using Trizol reagent (Invitrogen) according to the manufacturer's protocol. The reverse-transcriptase reaction was carried out using M-MLV reverse transcriptase (Invitrogen). Complementary DNA was generated by addition of 1 μ g of total RNA to a reaction mixture containing 0.5 μ g/ μ l oligo-deoxythymidine, 20 mM dNTP, 0.1 M dithiothreitol, 250 mM Tris-HCl, pH 8.3, 375 mM KCl, and 15 mM MgCl₂ and reaction at 37 °C for 90 min. The oligonucleotide primers used were 5'-GGAACCTTGACGCTTACAGTGACA-3' (forward) and 5'-CAAGTCTACATATACCCCAAG-3' (reverse) for VCAM-1 and 5'-GTAACCGTTGAACCCATT-3' (forward) and 5'-CCATCCAATCGGTAGTAGCG-3' (reverse) for 18S subunit ribosomal RNA. The amplification profile was 1 cycle of initial denaturation at 94 °C for 5 min and 30 cycles of denaturation at 94 °C for 1 min, primer annealing at 62 °C for 1 min, and extension at 72 °C for 5 min. PCR products were analyzed on ethidium bromide-stained 2% agarose gels.

Preparation of cell lysates and Western blot analysis

To prepare cell lysates, the cells were lysed for 1 h at 4 °C in 20 mM Tris-HCl, 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1% Triton X-100, 1 mM phenylmethylsulfonyl fluoride, pH 7.4, and then the lysate was centrifuged at 4000 g for 30 min at 4 °C and the supernatant retained. Western blot analyses were performed as described previously [15]. Briefly, samples of cell lysate (20 μ g of protein) were subjected to 10% sodium dodecyl sulfate (SDS)-PAGE and transferred to PVDF membranes, which were then treated with 3% nonfat milk in 0.1 M phosphate buffer for 1 h at room temperature to block nonspecific binding of antibody. The membranes were then incubated overnight at 4 °C with rabbit antibodies against human phospho-JNK, human phospho-ERK1/2, or human phospho-p38, all 1:1000 in phosphate-buffered saline (PBS), and then for 1 h at room temperature with HRP-conjugated goat anti-rabbit IgG antibodies (1:2000 in PBS; Santa Cruz Biotechnology), bound antibodies being detected using the chemiluminescence reagent Plus (NEN, Boston, MA, USA). The intensity of each band was quantified using a densitometer. Antibodies against GAPDH (1:5000; Santa Cruz Biotechnology) were used as loading controls.

VCAM-1 luciferase activity assay

The VCAM-1-luc plasmid was constructed by cloning the human VCAM-1 promoter (a region spanning bp -1716 to -119) into the pGL3-basic vector (Promega, Madison, WI, USA) and was used to transfect mouse embryonic fibroblast cells (NIH 3T3 cells) using Lipofectamine 2000 reagent (Invitrogen). To measure promoter activity, the cells were disrupted by sonication in lysis buffer (Promega), and then, after centrifugation at 13,000 g at 4 °C for 10 min, aliquots of the supernatants were tested for luciferase activity using the luciferase assay system (Promega). The luciferase activity was then normalized to the amount of cellular protein.

Immunocytochemical localization of NF- κ B p65

To localize NF- κ B expression in situ, confluent HUVECs (controls or cells treated for 24 h with different drugs) on slides were incubated in the absence or presence of 10 ng/ml TNF- α for 30 min, fixed in 4% paraformaldehyde in PBS, pH 7.4, for 15 min at 4 °C, and then reacted for 1 h at room temperature with rabbit anti-human NF- κ B p65 antibodies (1:100 dilution in PBS; GeneTex). After washes, the slides were incubated for 1 h at 37 °C with fluorescein isothiocyanate-




Fig. 1. Chemical structure of viscolin and its putative functional groups (boxed).

188	conjugated goat anti-rabbit IgG antibodies (Sigma) and viewed on a	246
189	fluorescence microscope.	247
190	<i>Electrophoretic mobility-shift assay (EMSA)</i>	248
191	The preparation of nuclear protein extracts and the EMSA	
192	conditions have been described previously [15]. Nuclear proteins	
193	were extracted using NE-PER reagent (Pierce, Rockford, IL, USA)	
194	according to the manufacturer's protocol. The AP-1 and NF- κ B binding	
195	activity of equal amounts (10 μ g) of nuclear protein was analyzed	
196	using a LightShift chemiluminescence EMSA kit (Pierce). The	
197	synthetic double-stranded oligonucleotides used as the probes in	
198	the gel-shift assay were 5'-AGTTGAGGGACTTCCAGGC-3' and 3'-	
199	TCAACTCCCTGAAAGGGTCCG-5' for NF- κ B and 5'-CGTTGATGAGT-	
200	CAGCCGAA-3' and 3'-CGAACTACTAGTCGGCCTT-5' for AP-1.	
201	<i>Detection of ROS (O₂⁻ and H₂O₂) production</i>	
202	The effect of viscolin on superoxide anion (O ₂ ⁻) and H ₂ O ₂	
203	production by HUVECs was determined by a fluorimetric assay	
204	using dihydroethidium (DHE) and Amplex red as the probe,	
205	respectively [16]. Confluent HUVECs were incubated with or without	
206	30 μ M viscolin for 24 h or 10 μ M DPI for 2 h. HUVECs were incubated	
207	with 20 μ M DHE for 20 min or with 50 μ M Amplex red/HRP for 10 min	
208	at 37 °C, and then 10 ng/ml TNF- α was added to the well for the	
209	indicated time. The fluorescence density (relative fluorescence units)	
210	was detected at 588 nm/630 nm and 544 nm/590 nm for excitatio-	
211	n/emission, respectively, for ethidium corresponding to O ₂ ⁻ and	
212	resorufin to H ₂ O ₂ , using a multidetection reader (SpectraMax M5;	
213	Molecular Devices, Sunnyvale, CA, USA).	
214	<i>Measurement of O₂⁻ production</i>	
215	The assay for the production of O ₂ ⁻ was based on the superoxide	
216	dismutase-inhibited reduction of ferricytochrome c and performed as	
217	described previously [17]. O ₂ ⁻ generation was measured after	
218	addition of 160 μ M NADPH to 800 μ l of relaxation buffer containing	
219	4 \times 10 ⁶ cell equivalents of membrane extract, 1.2 \times 10 ⁷ cell equiva-	
220	lents of cytosol, 2 μ M GTP- γ -S, 0.5 mg/ml ferricytochrome c, and	
221	100 μ M SDS. To facilitate the assembly of NADPH oxidase components,	
222	all constituents (excluding NADPH) were incubated at room temper-	
223	ature for 3 min, then any test drug was added and the mixture	
224	incubated for 1 min at room temperature, then NADPH was added and	
225	the mixture incubated for 10 min at 37 °C. Changes in absorbance at	
226	550 nm due to reduction of ferricytochrome c were monitored.	
227	<i>Plasma membrane preparation, NADPH oxidase activity assay, and</i>	
228	<i>Western blot analysis of p47^{phox}</i>	
229	The cytosolic and plasma membrane fractions were prepared as	
230	described previously, with modification [18]. Briefly, HUVECs were	
231	lysed in lysis buffer A (20 mM Tris-HCl, 10 mM EGTA, 2 mM EDTA,	
232	2 mM dithiothreitol, 1 mM phenylmethylsulfonyl fluoride (PMSF),	
233	25 μ g/ml aprotinin, and 10 μ g/ml leupeptin). Cell lysates were centri-	
234	fuged at 16,000 g for 20 min at 4 °C. The supernatant was collected and	
235	designated the cytosolic fraction. The pellets were resuspended in lysis	
236	buffer B (0.5% sodium dodecyl sulfate, 1% NP-40, 1 mM Na ₃ VO ₄ , 1 mM	
237	NaF, 1 mM PMSF, 25 μ g/ml aprotinin, and 10 μ g/ml leupeptin). Western	
238	blot analysis for p47 ^{phox} was performed on the plasma membrane	
239	fractions as described above, using a monoclonal mouse antibody	
240	against p47 ^{phox} (BD Biosciences Pharmingen, San Jose, CA, USA). For	
241	NADPH oxidase assay, HUVECs were lysed in lysis buffer containing	
242	20 mM monobasic potassium phosphate (pH 7.0), 1 mM EGTA,	
243	10 μ g/ml aprotinin, 0.5 μ g/ml leupeptin, 0.5 mM PMSF. Plasma mem-	
244	brane fractions were measured in a lucigenin chemiluminescence assay	
245	using 1 mM lucigenin (Sigma) and 5 mM NADPH (Sigma) as described	
	previously [18]. Chemiluminescence as relative light units was	246
	measured in a microtiter luminometer (SpectraMax M5; Molecular	247
	Devices) as an indicator of enzyme activity.	248
	<i>Endothelial cell–leukocyte adhesion assay</i>	249
	U937 cells, originally derived from a human histiocytic lymphoma	250
	and obtained from the American Type Culture Collection (Rockville,	251
	MD, USA) and grown in RPMI 1640 medium (M.A. Bioproducts,	252
	Walkersville, MD, USA), were labeled for 1 h at 37 °C with BCECF/AM	253
	(10 mM; Boehringer Mannheim, Mannheim, Germany). Labeled U937	254
	cells (10 ⁶) were added to HUVECs (10 ⁶) in a 48-well plate and	255
	incubation continued for 1 h, then the nonadherent cells were	256
	removed by two gentle washes with PBS and the number of bound	257
	U937 cells was evaluated by fluorescence microscopy.	258
	<i>Mouse model and immunohistochemical staining</i>	259
	Male 8-week-old C57BL6 mice ($n = 24$), weighing between 25 and	260
	35 g, were purchased from the National Taiwan University (Taipei,	261
	Taiwan). All procedures involving experimental animals were	262
	performed in accordance with the guidelines for animal care of the	263
	National Taiwan University and complied with the <i>Guide for the Care</i>	264
	<i>and Use of Laboratory Animals</i> , NIH publication No. 86–23, revised	265
	1985. The mice were randomly divided into four groups, which were	266
	to be treated with DMSO, TNF- α , TNF- α plus viscolin, or viscolin. The	267
	mice were injected intraperitoneally (ip) with viscolin (10 mg/kg/day	268
	in 50 μ l of DMSO) or DMSO (50 μ l) for 5 days and then were left	269
	untreated or were injected ip with TNF- α (10 μ g/kg/day) for the next	270
	3 days. They were then anesthetized by ip injection of 30–40 mg/kg	271
	pentobarbital and sacrificed, and the thoracic aorta was dissected out,	272
	immersion-fixed in 4% buffered paraformaldehyde, paraffin-embed-	273
	ded, and cross-sectioned for immunohistochemistry. To determine	274
	the level of expression of VCAM-1 in aortic walls and whether it was	275
	associated with endothelial cells, two serial sections were examined	276
	by immunostaining for, respectively, von Willebrand factor (vWF;	277
	marker for endothelial cells) or VCAM-1. The first section was	278
	incubated sequentially for 1 h at 37 °C with mouse monoclonal anti-	279
	human vWF antibody (1:50 dilution; Neomarkers, Fremont, CA, USA)	280
	and 1 h at room temperature with HRP-conjugated goat anti-mouse	281
	IgG antibodies (1:200 dilution in PBS; Sigma) and bound antibody	282
	visualized using 3,3'-diaminobenzidine (Sigma–Aldrich). The second	283
	section was incubated with rabbit antibodies against human VCAM-1	284
	(1:100; Santa Cruz Biotechnology) at 4 °C for 1 h, washed with PBS,	285
	and then incubated with HRP-conjugated second antibody and then	286
	with the same chromogen as above.	287
	<i>Statistical analysis of data</i>	288
	All values are presented as the mean \pm SEM and were analyzed using	289
	Student's <i>t</i> test. Statistical significance was determined as $P < 0.05$.	290
	Results	291
	<i>Viscolin reduces VCAM-1 mRNA and protein expression in TNF-α-treated</i>	292
	<i>HUVECs</i>	293
	When the cytotoxicity of TNF- α or viscolin for HUVECs was assessed	294
	by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide	295
	assay after 24 h of incubation, cell viability was not affected by the	296
	presence of 10 ng/ml TNF- α or 1–30 μ M viscolin (data not shown).	297
	TNF- α (10 ng/ml) induced significant VCAM-1 protein expression in	298
	HUVECs, which peaked at 6 h (Fig. 2A). As shown in Fig. 2B, when HUVECs	299
	were pretreated for 24 h with 1, 3, 10, or 30 μ M viscolin before incubation	300
	with 10 ng/ml TNF- α for 6 h, TNF- α -induced VCAM-1 expression was	301
	reduced respectively to 91 \pm 14, 96 \pm 16, 38 \pm 7, or 34 \pm 16% of control	302




Fig. 2. Viscolin inhibits the TNF- α -induced increase in VCAM-1 mRNA and protein levels in HUVECs. (A) HUVECs were treated with TNF- α (10 ng/ml) for the indicated times, then the protein levels in the cell lysates were measured on Western blots. (B) HUVECs were incubated with the indicated concentrations of viscolin for 24 h and then with 10 ng/ml TNF- α for 6 h in the continued presence of the same concentration of viscolin, and VCAM-1 protein in cell lysates was measured by Western blot. GAPDH was used as the loading control. (C) Analysis of VCAM-1 mRNA levels in untreated HUVECs or HUVECs preincubated with or without 10 μ M viscolin for 24 h and then incubated with 10 ng/ml TNF- α for 6 h. Total RNA was analyzed by RT-PCR after normalization to 18S levels. (D) NIH 3 T3 cells were transfected with a luciferase plasmid containing the VCAM-1 promoter for 24 h and then were incubated with or without 10 μ M viscolin for 24 h before the addition of TNF- α (10 ng/ml) for a further 6 h. In (A–D), the data are expressed as a fold value compared to the control value and are the means \pm SEM for three separate experiments. * P <0.05 compared to the untreated cells. † P <0.05 compared to the TNF- α -treated cells.

303 levels, the reductions caused by the two highest concentrations being
304 significant. In all subsequent experiments, unless otherwise specified,
305 10 ng/ml TNF- α and 30 μ M viscolin were used.

306 To determine whether the effects of TNF- α alone or together with
307 viscolin on VCAM-1 expression were exerted at the transcriptional
308 level, VCAM-1 mRNA levels were measured by RT-PCR. As shown in
309 Fig. 2C, unstimulated HUVECs produced low amounts of VCAM-1
310 mRNA, and 6 h treatment with TNF- α resulted in a marked increase in
311 levels. This increase was markedly inhibited by 24 h preincubation
312 with 10 μ M viscolin (36 \pm 2% inhibition).

313 The effect of viscolin on VCAM-1 gene transcription was confirmed
314 using the luciferase gene activity assay. The VCAM-1 luciferase
315 reporter gene was transfected into NIH 3 T3 cells, which have a high
316 transfection efficiency, rather than the very hard to transfect primary
317 HUVECs [19], which were then stimulated with TNF- α for 6 h. Fig. 2D
318 shows that TNF- α treatment stimulated VCAM-1 luciferase activity
319 and that preincubation of the cells for 24 h with viscolin significantly
320 reduced the effect of TNF- α by 35 \pm 4%. These results suggest that
321 viscolin significantly inhibited TNF- α -induced VCAM-1 expression at
322 the transcriptional level.

323 *The viscolin-induced reduction in TNF- α -induced VCAM-1 expression is*
324 *partly dependent on inhibition of JNK phosphorylation*

325 Because TNF- α -induced inflammation involves the secretion of
326 inflammatory cytokines via the MAPK pathways [20], we next investi-

327 gated whether TNF- α -induced VCAM-1 expression was mediated by
328 activation of MAPKs. As shown in Figs. 3A–C, TNF- α induced transient
329 phosphorylation of ERK1/2, JNK, and p38 in HUVECs, with the maximal
330 response being seen within 15 min, followed by a decline to the basal level
331 within 60 min. In addition, pretreatment for 1 h with the indicated
332 concentrations of PD98059 (an ERK1/2 inhibitor), SP600125 (a JNK
333 inhibitor), or SB203580 (a p38 inhibitor) inhibited the TNF- α -induced
334 VCAM-1 expression seen at 6 h of TNF- α treatment (Figs. 3D–F). These
335 results suggest that TNF- α -induced VCAM-1 expression is mediated by
336 activation of MAPKs.

337 To determine the potential targets that were negatively regulated
338 by viscolin, the cells were preincubated with viscolin for 24 h and then
339 incubated with TNF- α for 15 min. As shown in Figs. 3G–I, pretreat-
340 ment with viscolin significantly inhibited TNF- α -induced JNK phos-
341 phosphorylation by 36 \pm 4% (Fig. 3H), but had no significant effect on
342 ERK1/2 and p38 phosphorylation. These results suggest that viscolin
343 inhibits TNF- α -induced VCAM-1 expression partly by inhibiting
344 TNF- α -induced JNK phosphorylation.

345 *Viscolin decreases NF- κ B activation and NF- κ B p65 nuclear translocation*
346 *in TNF- α -treated HUVECs*

347 Because the VCAM-1 gene promoter contains consensus binding
348 sites for AP-1 and NF- κ B [21,22], we investigated whether viscolin
349 inhibited TNF- α -induced VCAM-1 expression via an effect on these
350 transcription factors. Gel-shift assays were performed to determine the




Fig. 3. The viscolin-mediated reduction in TNF- α -induced VCAM-1 expression is partly dependent on inhibition of JNK phosphorylation. (A–C) HUVECs were treated with 10 ng/ml TNF- α for the indicated times, then the cell lysate was analyzed for MAPK phosphorylation by Western blot using antibodies against (A) p-ERK1/2, (B) p-JNK, (C) or p-p38. (D–F) The cells were preincubated for 1 h with the indicated concentrations of (D) PD98059 (ERK1/2 inhibitor), (E) SP600125 (JNK inhibitor), or (F) SB203580 (p38 inhibitor) and then were treated with TNF- α for 6 h and the cell lysates were analyzed for VCAM-1 expression by Western blot. (G–I) Western blot analysis showing the effects of viscolin treatment on the phosphorylation of (G) p-ERK1/2, (H) p-JNK, or (I) p-p38 in TNF- α -treated HUVECs. HUVECs were incubated for 24 h with or without 30 μ M viscolin, then the cells were incubated with 10 ng/ml of TNF- α for 15 min and aliquots of cell lysates containing equal amounts of protein subjected to immunoblotting with the indicated antibodies. The data are expressed as a fold of the control value and are the means \pm SEM for three separate experiments. GAPDH was used as the loading control. * P <0.05 compared to the untreated cells. † P <0.05 compared to the TNF- α -treated cells.

351 effects of viscolin on AP-1 and NF- κ B activation in TNF- α -treated
 352 HUVECs. As shown in Figs. 4A and B, low basal levels of AP-1 and NF- κ B
 353 binding activity were detected in untreated control cells and binding
 354 was significantly increased by 30 min treatment with TNF- α . Pretreat-
 355 ment with viscolin for 24 h had no effect on TNF- α -induced AP-1
 356 activation, but blocked the increase in NF- κ B binding activity. To
 357 determine whether NF- κ B activation was involved in the pretransla-
 358 tional effects of viscolin on VCAM-1 expression, we examined NF- κ B p65
 359 protein levels in the nuclei of TNF- α -treated HUVECs by Western blot
 360 and immunofluorescence staining. Western blot (Fig. 4C) showed that
 361 higher levels of p65 and phospho-p65 were found in the nuclei of
 362 TNF- α -stimulated HUVECs compared to control HUVECs and that
 363 viscolin pretreatment significantly reduced the expression of p65 and p-
 364 p65. Consistent with the Western blot findings, HUVECs stimulated with
 365 TNF- α for 30 min showed marked NF- κ B p65 staining in the nuclei (T on

Fig. 4D) by immunofluorescence staining, whereas viscolin-pretreated
 366 cells (24 h; T + Vis) showed weaker nuclear NF- κ B expression, but
 367 stronger staining in the cytoplasm. The NF- κ B p65 translocation induced
 368 by TNF- α was also significantly inhibited by pretreatment for 24 h with
 369 NAC (10 mM), DPI (10 μ M), or APO (100 μ M) (Figs. 4C and D).
 370 Furthermore, the stimulatory effect of TNF- α on VCAM-1 levels was
 371 blocked by co-incubation with 0–10 μ M parthenolide, an NF- κ B
 372 inhibitor (Fig. 4E). These results suggest that viscolin inhibits the
 373 TNF- α -induced VCAM-1 expression by inhibiting NF- κ B activation. 374

Viscolin inhibits TNF- α -induced ROS production, NADPH oxidase
 375 activity, and p47^{phox} translocation in HUVECs 376

Because previous studies have shown that viscolin inhibits super-
 377 oxide anion production in human neutrophils [7,8], we investigated 378




Fig. 4. The viscolin-induced downregulation of VCAM-1 expression in TNF- α -stimulated HUVECs is mediated by inhibition of NF- κ B activation and NF- κ B p65 nuclear translocation. (A, B) Nuclear extracts prepared from untreated cells or from cells with or without 24 h pretreatment with 30 μ M viscolin and then incubated with 10 ng/ml TNF- α for 30 min were tested for (A) AP-1 or (B) NF- κ B DNA binding activity by EMSA. (C, D) Western blot and immunofluorescence staining for NF- κ B p65. HUVECs were preincubated for 24 h with 30 μ M viscolin or for 2 h with 10 mM NAC, 10 μ M DPI, or 100 μ M APO and then were treated with 10 ng/ml TNF- α for 30 min. Representative results from three separate experiments are shown. Bar, 100 μ m. (E) Cells were co-incubated for 24 h with 0–10 μ M parthenolide (Par; NF- κ B inhibitor) and 10 ng/ml TNF- α , then cell lysates were prepared and assayed for VCAM-1 by Western blot. The data are expressed as a fold of the control value and are the means \pm SEM for three separate experiments. GAPDH was used as the loading control. * P < 0.05 compared to the untreated cells. † P < 0.05 compared to the TNF- α -treated cells.

379 whether it had an antioxidant effect on TNF- α -treated HUVECs. First, we
 380 examined its effect on TNF- α -induced $O_2^{\cdot-}$ and H_2O_2 production using
 381 DHE and Amplex red as the probe, respectively. As shown in Figs. 5A and
 382 B, TNF- α induced $O_2^{\cdot-}$ and H_2O_2 production in a time-dependent
 383 manner. Twenty-four hours pretreatment with viscolin dramatically
 384 decreased TNF- α -induced $O_2^{\cdot-}$ and H_2O_2 production by 45 ± 1 and $22 \pm$
 385 2%, respectively, as did 2 h preincubation with the NADPH oxidase
 386 inhibitor DPI (Figs. 5C and D). These results suggest that viscolin has a
 387 potent antioxidant activity on TNF- α -induced ROS production and that
 388 this may be mediated through inhibition of NADPH oxidase activity.
 389 Because NADPH oxidase contains membrane-bound components (NOX
 390 and p22^{phox}) and cytosolic components (p40^{phox}, p47^{phox}, p67^{phox}, and
 391 Rac) [23], the membrane and cytosolic subunits of NADPH oxidase
 392 isolated from unstimulated HUVECs were assembled using SDS and
 393 incubated with viscolin or APO in the presence of NADPH. As shown in
 394 Fig. 5E, NADPH treatment induced marked superoxide anion produc-
 395 tion, which was significantly reduced by addition of viscolin or APO for

2 min before assay. Furthermore, HUVECs were treated with 10 ng/ml
 TNF- α for 20 min, and then the membrane fraction was assayed for
 NADPH oxidase activity. As shown in Fig. 5F, TNF- α addition resulted in
 a significant increase in NADPH oxidase activity, which was inhibited by
 viscolin treatment. We then determined whether this effect of viscolin
 was associated with translocation of p47, as this translocation
 mechanism was been reported to play an important role in the
 activation of NADPH oxidase [23]. Stimulation of HUVECs with
 10 ng/ml TNF- α for 20 min increased membrane p47^{phox} expression
 compared with the untreated cells, whereas 24 h pretreatment with
 viscolin resulted in a decrease in the membrane p47^{phox} content in
 TNF- α -treated HUVECs (Fig. 5G), suggesting an effect on the translo-
 cation of p47^{phox} from the cytoplasm to the membrane.

Because several lines of evidence have indicated that ROS production
 is the mediator inducing VCAM-1 expression [24,25], the role of ROS
 production in TNF- α -induced VCAM-1 expression was investigated. As
 shown in Figs. 5H–J, 2 h pretreatment with the antioxidant NAC or the




Fig. 5. Viscolin reduces TNF- α -induced ROS production, NADPH oxidase activity, and p47^{phox} translocation in HUVECs. (A, B) HUVECs were incubated with DHE and Amplex red–HRP, and then 10 ng/ml TNF- α was added to the well for the indicated times, and ethidium and resorufin fluorescence was measured for the generation of O₂^{•-} and H₂O₂, respectively. (C, D) The cells were pretreated with 30 μ M viscolin for 24 h or with 10 μ M DPI for 2 h before addition of 10 ng/ml TNF- α for 15 min. The methods for the measurement of O₂^{•-} and H₂O₂ were described under Materials and methods. (E) The cytosolic and membrane fractions from unstimulated HUVECs were incubated with the reaction mixture for NADPH oxidase assembly as described under Materials and methods and then were left untreated or were incubated for 2 min with 30 μ M viscolin or 100 μ M APO. Then ferricytochrome c reduction was measured at 550 nm. (F) Control cells or cells pretreated with 30 μ M viscolin for 24 h or 10 μ M DPI for 2 h were incubated with 10 ng/ml TNF- α for 20 min, then the plasma membrane protein was assayed with a superoxide-dependent lucigenin chemiluminescence assay. (G) Western blot of p47^{phox} levels in the membrane fractions of HUVECs pretreated with viscolin for 24 h or with 10 μ M DPI for 2 h and then stimulated with 10 ng/ml TNF- α for 20 min. (H–J) The cells were incubated for 2 h with the indicated concentrations of (H) 0–10 mM NAC, (I) 0–10 μ M DPI, (J) or 0–200 μ M APO and then were treated with TNF- α for 6 h and the cell lysates were analyzed for VCAM-1 expression by Western blot. (K) The cells were preincubated for 2 h with 10 mM NAC, 10 μ M DPI, or 100 μ M APO and then were treated with TNF- α for 15 min and the cell lysates were analyzed for JNK phosphorylation by Western blotting. Values are presented as the means \pm SEM. * P <0.05 compared to the untreated cells. † P <0.05 compared to the TNF- α -treated cells.

413 NADPH oxidase inhibitor DPI or APO significantly attenuated TNF- α -
 414 induced VCAM-1 expression in a concentration-dependent manner. In
 415 addition, as shown in Fig. 5K, 2 h pretreatment with antioxidants (NAC,
 416 DPI, or APO) partly inhibited TNF- α -induced JNK phosphorylation, the
 417 effects being similar to those of viscolin. These results suggest that
 418 NADPH oxidase-derived ROS production plays a critical role in TNF- α -
 419 induced VCAM-1 expression.

420 Viscolin reduces the adhesion of monocytes to TNF- α -treated HUVECs

421 To explore the effects of viscolin on the endothelial cell–leukocyte
 422 interaction, we examined the adhesion of U937 cells to TNF- α -
 423 activated HUVECs. As shown in Fig. 6, control confluent HUVECs (C)
 424 incubated with U937 cells for 1 h showed minimal binding, but

adhesion was substantially increased when the HUVECs were
 425 pretreated with TNF- α for 6 h (T). Pretreatment of HUVECs with
 426 viscolin for 24 h (T + Vis) reduced the number of U937 cells adherent
 427 to TNF- α -treated HUVECs by 46 \pm 4% compared to TNF- α alone. The
 428 involvement of VCAM-1 in the adhesion of U937 cells to TNF- α -
 429 treated HUVECs was examined by pretreatment of the cells with anti-
 430 VCAM-1 antibody. When HUVECs were pretreated with 1 (T + VCAM-
 431 1 Ab-1) or 2 μ g/ml (T + VCAM-1 Ab-2) anti-VCAM-1 antibody for 1 h
 432 and then incubated with TNF- α , the binding of U937 cells to HUVECs
 433 was significantly lower than that to non-antibody-treated TNF- α -
 434 stimulated cells, showing that VCAM-1 plays a major role in the
 435 adhesion of U937 cells to TNF- α -treated HUVECs. The adherence of
 436 TNF- α -treated U937 cells to HUVECs was also inhibited by 1 h
 437 pretreatment with 10 μ M PD98059 (T + PD), SP600125 (T + SP), 438




Fig. 6. Viscolin reduces the adhesion of U937 cells to TNF- α -stimulated HUVECs. Cells were left untreated or were pretreated for 24 h with 30 μ M viscolin; or for 1 h with 1 or 2 μ g/ml anti-VCAM-1 antibodies; or for 1 h with 10 μ M PD98059, SP600125, SB203580, or parthenolide; or for 2 h with 10 mM NAC, 10 μ M DPI, or 100 μ M APO. Then they were incubated with 10 ng/ml TNF- α for 6 h in the continued presence of the inhibitor. (A) Representative fluorescence photomicrographs showing the effects on TNF- α -induced adhesion of fluorescein-labeled U937 cells to HUVECs. C, untreated cells. Bar, 100 μ m. (B) The number of U937 cells bound per high power field was counted. The data are expressed as the means \pm SEM for three separate experiments. * P <0.05 compared to the untreated cells. † P <0.05 compared to the TNF- α -treated cells.




Fig. 7. Immunohistochemical staining for vWF or VCAM-1 expression in serial sections of thoracic aortas from mice. Mice were treated with DMSO (C), TNF- α , TNF- α + viscolin, or viscolin alone as described under Materials and methods, then serial sections were stained for vWF (endothelial cell marker) or VCAM-1. The lumen is uppermost in all sections. The reaction product and the internal elastic membrane are indicated by an arrowhead and an arrow, respectively. Bar, 50 μ m.

439 SB203580 (T + SB), or parthenolide (T + Par). Similarly, the
440 adherence of U937 cells to TNF- α -treated HUVECs was also inhibited
441 by 2 h pretreatment with antioxidants (NAC, DPI, or APO).

442 Viscolin reduces VCAM-1 protein expression in the thoracic aorta in 443 TNF- α -injected mice

444 To determine the effect of viscolin on VCAM-1 expression in vivo,
445 mice were injected with viscolin for 5 days before injection with
446 TNF- α for 3 days, then immunohistochemical staining was performed
447 to detect the expression of VCAM-1 on serial sections of thoracic aorta,
448 using vWF as an endothelial cell marker. As shown in Fig. 7, in the
449 control (C) and viscolin-treated (Vis) groups, no VCAM-1 staining was
450 seen on the vascular wall, whereas in the TNF- α -treated group
451 (TNF- α), strong VCAM-1 staining was seen on the luminal surface. In
452 contrast, preadministration of viscolin resulted in weak VCAM-1
453 staining in the TNF- α -treated animals (TNF- α + Vis).

454 Discussion

455 In this study, we demonstrated that viscolin treatment reduced
456 VCAM-1 expression both in vitro in TNF- α -stimulated HUVECs and in
457 vivo in the thoracic aorta of TNF- α -treated mice. Viscolin also
458 inhibited the binding of the human monocytic cell line U937 to
459 TNF- α -stimulated HUVECs. These effects were inhibited by SP600125,

a JNK inhibitor, or parthenolide, an NF- κ B inhibitor, showing that they
460 were partly mediated through inhibition of JNK phosphorylation and
461 NF- κ B activation. In addition, viscolin attenuated the increase in
462 VCAM-1 mRNA expression and VCAM-1 promoter activity induced by
463 TNF- α . Furthermore, viscolin had a scavenging effect on the
464 generation of ROS as well as on the decreased NADPH oxidase activity.
465

466 Viscolin, isolated from *V. coloratum*, was chosen for testing, as *V.*
467 *coloratum* has long been used in traditional Chinese medicine to treat
468 inflammatory diseases. Antioxidative and anti-inflammatory actions
469 are two of the pharmacological properties proposed to underlie its
470 beneficial effects [7–9]. A partially purified fraction from the
471 chloroform extract of *V. coloratum* (PPE-SVC) has been shown to
472 inhibit the generation of superoxide anions by formyl-L-methionyl-L-
473 leucyl-L-phenylalanine (fMLP)-activated human neutrophils, and
474 purified viscolin, a major active component of PPE-SVC, inhibits
475 the generation of superoxide anion and the release of elastase in fMLP-
476 activated human neutrophils [7]. Viscolin suppresses ROS and nitric
477 oxide generation in leukocytes and microglial cells and, in addition,
478 attenuates proinflammatory cytokine production [9]. This study is the
479 first to report that viscolin strongly reduces levels of VCAM-1 mRNA
480 and protein in TNF- α -treated HUVECs. The present results also show
481 that viscolin reduced TNF- α -induced VCAM-1 promoter activity.

482 Our results demonstrated that TNF- α induced time-dependent
483 phosphorylation of MAPKs (ERK1/2, JNK, and p38) and that the
484 increases in VCAM-1 expression and U937 cell adhesion induced by
485 TNF- α were inhibited by PD98059, SP600125, or SB203580. These
486 results show that activation of MAPKs is necessary for TNF- α -induced

VCAM-1 expression in HUVECs. Consistent with these findings, TNF- α -induced VCAM-1 expression in human tracheal smooth muscle cells requires activation of MAPKs [26]. Furthermore, our results demonstrated that viscolin inhibited the TNF- α -induced phosphorylation of JNK, but not that of ERK1/2 or p38 (Fig. 3), suggesting that the inhibitory effect of viscolin on VCAM-1 expression is mediated, in part, by JNK inhibition. Because a previous study showed that ROS regulate both protein kinases and protein phosphatases [27], one of our future aims is to determine the protein phosphatases involved in the dephosphorylation of JNK that are regulated by viscolin. In addition, our results also showed that viscolin inhibited the TNF- α -induced increase in VCAM-1 mRNA levels. Although we cannot rule out the possibility that viscolin may affect the stability of VCAM-1 mRNA, viscolin was found to inhibit the TNF- α -induced promoter activity of VCAM-1 (Fig. 2D). These results suggest that viscolin attenuates VCAM-1 expression induced by TNF- α , at least in part, through a transcriptional mechanism.

Several lines of evidence indicate that TNF- α induces ROS production in endothelial cells [10,24,25,28]. Consistent with these previous results, our study showed that it rapidly induced ROS production and that this was inhibited by the NADPH oxidase inhibitors DPI and APO. These results suggest that TNF- α induces ROS production via activation of NADPH oxidase. ROS seems to be a second messenger in the TNF- α -induced signal transduction pathway that regulates VCAM-1 expression [10,24,25,28]. In our study, antioxidants (NAC, DPI, or APO) inhibited the TNF- α -induced increase in VCAM-1 expression (Figs. 5H–J) and U937 cell adhesion (Fig. 6A), showing that ROS mediated the effects of TNF- α on VCAM-1 expression. In addition, preincubation with viscolin effectively attenuated the ROS production induced by TNF- α in HUVECs. Moreover, pretreatment with antioxidants (NAC, DPI, APO) inhibited TNF- α -induced JNK phosphorylation to a similar extent compared to viscolin. These results suggest that viscolin inhibits TNF- α -induced VCAM-1 expression via its antioxidative properties. Furthermore, we demonstrate that viscolin inhibited NADPH oxidase activity and p47 expression in the membrane fraction of TNF- α -treated HUVECs. Because of their chemical structure, a benzene ring with adjacent methoxy–hydroxyl groups, flavonoids are potent inhibitors of NADPH oxidase activity [29]. As the chemical structure of viscolin (Fig. 1) is similar to that of flavonoids, it may have a similar inhibitory effect on NADPH oxidase activity. Future studies are necessary to clarify the role of viscolin on NOX activity, as DPI and apocynin have been reported to inhibit NOX activity as well as affecting other reactive species and enzymes.

The VCAM-1 gene promoter contains consensus binding sites for AP-1 and NF- κ B [21,22]. Our results showed that the binding activity of NF- κ B and AP-1 was activated by TNF- α and that pretreatment with viscolin significantly inhibited the TNF- α -induced increase in binding activity of NF- κ B, but not that of AP-1. In addition, several reports have shown that natural products with antioxidant activity inhibit the TNF- α -induced activation of redox-sensitive NF- κ B [10,24,25,28]. Pretreatment with an NF- κ B inhibitor suppressed the TNF- α -induced increase in VCAM-1 expression and U937 cell adhesion, suggesting that viscolin attenuates VCAM-1 expression via a reduction in NF- κ B binding activity. Our results showed that viscolin and the antioxidants NAC, DPI, and APO significantly attenuated NF- κ B binding activity and NF- κ B p65 translocation and that these effects may be due to its antioxidative activity. Viscolin has anti-inflammatory and antioxidative properties, based on the above findings. Because atherosclerosis is a chronic inflammatory disease [1,2], viscolin may be beneficial for the prevention of inflammation and atherosclerosis.

In conclusion, our study demonstrates that viscolin reduces VCAM-1 expression under inflammatory conditions both in vitro and in vivo. Our results show that the inhibitory effect on VCAM-1 expression is partly mediated by inhibition of JNK phosphorylation, NF- κ B activation, and ROS production. Our results demonstrate the anti-inflammatory and antioxidative effects of viscolin, an active component of *V. coloratum*, on endothelial cells and suggest that this

compound may provide a chemical backbone for the development of therapeutic agents.

Acknowledgments

This work was supported by research grants from the National Science Council (NSC-99-2320-B-002-0220MY3) and the Cooperative Research Program of the NTU and CMUCM (99F0080-303), Taiwan, Republic of China.

References

- Ross, R. Atherosclerosis—an inflammatory disease. *N. Engl. J. Med.* **340**:115–126; 1999.
- Rao, R. M.; Yang, L.; Garcia-Cardena, G.; Luscinskas, F. W. Endothelial-dependent mechanisms of leukocyte recruitment to the vascular wall. *Circ. Res.* **101**:234–247; 2007.
- Cybulsky, M. I.; Gimbrone Jr., M. A. Endothelial expression of a mononuclear leukocyte adhesion molecule during atherogenesis. *Science* **251**:788–791; 1991.
- Cybulsky, M. I.; Iiyama, K.; Li, H.; Zhu, S.; Chen, M.; Iiyama, M.; Davis, V.; Gutierrez-Ramos, J. C.; Connelly, P. W.; Milstone, D. S. A major role for VCAM-1, but not ICAM-1, in early atherosclerosis. *J. Clin. Invest.* **107**:1255–1262; 2001.
- Hordijk, P. L. Endothelial signalling events during leukocyte transmigration. *FEBS J.* **273**:4408–4415; 2006.
- Preiss, D. J.; Sattar, N. Vascular cell adhesion molecule-1: a viable therapeutic target for atherosclerosis? *Int. J. Clin. Pract.* **61**:697–701; 2007.
- Leu, Y. L.; Hwang, T. L.; Chung, Y. M.; Hong, P. Y. The inhibition of superoxide anion generation in human neutrophils by *Viscum coloratum*. *Chem. Pharm. Bull. (Tokyo)* **54**:1063–1066; 2006.
- Hwang, T. L.; Leu, Y. L.; Kao, S. H.; Tang, M. C.; Chang, H. L. Viscolin, a new chalcone from *Viscum coloratum*, inhibits human neutrophil superoxide anion and elastase release via a cAMP-dependent pathway. *Free Radic. Biol. Med.* **41**:1433–1441; 2006.
- Su, C. R.; Shen, Y. C.; Kuo, P. C.; Leu, Y. L.; Damu, A. G.; Wang, Y. H.; Wu, T. S. Total synthesis and biological evaluation of viscolin, a 1,3-diphenylpropane as a novel potent anti-inflammatory agent. *Bioorg. Med. Chem. Lett.* **16**:6155–6160; 2006.
- Yoon, J. J.; Lee, Y. J.; Kim, J. S.; Kang, D. G.; Lee, H. S. Protective role of betulinic acid on TNF- α -induced cell adhesion molecules in vascular endothelial cells. *Biochem. Biophys. Res. Commun.* **391**:96–101; 2010.
- Collins, T.; Read, M. A.; Neish, A. S.; Whitley, M. Z.; Thanos, D.; Maniatis, T. Transcriptional regulation of endothelial cell adhesion molecules: NF- κ B and cytokine-inducible enhancers. *FASEB J.* **9**:899–909; 1995.
- Kim, H. J.; Park, K. G.; Yoo, E. K.; Kim, Y. H.; Kim, Y. N.; Kim, H. S.; Kim, H. T.; Park, J. Y.; Lee, K. U.; Jang, W. G.; Kim, J. G.; Kim, B. W.; Lee, I. K. Effects of PGC-1 α on TNF- α -induced MCP-1 and VCAM-1 expression and NF- κ B activation in human aortic smooth muscle and endothelial cells. *Antioxid. Redox Signal.* **9**:301–307; 2007.
- Griendling, K. K.; Sorescu, D.; Lassegue, B.; Ushio-Fukai, M. Modulation of protein kinase activity and gene expression by reactive oxygen species and their role in vascular physiology and pathophysiology. *Arterioscler. Thromb. Vasc. Biol.* **20**:2175–2183; 2000.
- Jaffe, E. A.; Nachman, R. L.; Becker, C. G.; Minick, C. R. Culture of human endothelial cells derived from umbilical veins: identification by morphologic and immunologic criteria. *J. Clin. Invest.* **52**:2745–2756; 1973.
- Lin, S. J.; Shyue, S. K.; Shih, M. C.; Chu, T. H.; Chen, Y. H.; Ku, H. H.; Chen, J. W.; Tam, K. B.; Chen, Y. L. Superoxide dismutase and catalase inhibit oxidized low-density lipoprotein-induced human aortic smooth muscle cell proliferation: role of cell-cycle regulation, mitogen-activated protein kinases, and transcription factors. *Atherosclerosis* **190**:124–134; 2007.
- Xu, S.; He, T.; Vokurkova, M.; Touyz, R. M. Endothelial cells negatively modulate reactive oxygen species generation in vascular smooth muscle cells: role of thioredoxin. *Hypertension* **54**:427–433; 2009.
- Yu, H. P.; Hsieh, P. W.; Chang, Y. J.; Chung, P. J.; Kuo, L. M.; Hwang, T. L. DSM-RX78, a new phosphodiesterase inhibitor, suppresses superoxide anion production in activated human neutrophils and attenuates hemorrhagic shock-induced lung injury in rats. *Biochem. Pharmacol.* **78**:983–992; 2009.
- Chen, Y. L.; Hu, C. S.; Lin, F. Y.; Chen, Y. H.; Sheu, L. M.; Ku, H. H.; Shiao, M. S.; Chen, J. W.; Lin, S. J. Salvianolic acid B attenuates cyclooxygenase-2 expression in vitro in LPS-treated human aortic smooth muscle cells and in vivo in the apolipoprotein E-deficient mouse aorta. *J. Cell. Biochem.* **98**:618–631; 2006.
- Tanner, F. C.; Carr, D. P.; Nabel, G. J.; Nabel, E. G. Transfection of human endothelial cells. *Cardiovasc. Res.* **35**:522–528; 1997.
- Muslin, A. J. MAPK signalling in cardiovascular health and disease: molecular mechanisms and therapeutic targets. *Clin. Sci.* **115**:203–218; 2008.
- Iademarco, M. F.; McQuillan, J. J.; Rosen, G. D.; Dean, D. C. Characterization of the promoter for vascular cell adhesion molecule-1 (VCAM-1). *J. Biol. Chem.* **267**:16323–16329; 1992.
- Cybulsky, M. I.; Fries, J. W.; Williams, A. J.; Sultan, P.; Eddy, R.; Byers, M.; Shows, T.; Gimbrone Jr., M. A.; Collins, T. Gene structure, chromosomal location, and basis for alternative mRNA splicing of the human VCAM1 gene. *Proc. Natl. Acad. Sci. U. S. A.* **88**:7859–7863; 1991.
- Gabor, C.; Taylor, W. R.; Pagano, P. J. NOX and inflammation in the vascular adventitia. *Free Radic. Biol. Med.* **47**:1254–1266; 2009.

- 631 [24] Kim, S. R.; Bae, Y. H.; Bae, S. K.; Choi, K. S.; Yoon, K. H.; Koo, T. H.; Jang, H. O.; Yun, I.;
632 Kim, K. W.; Kwon, Y. G.; Yoo, M. A.; Bae, M. K. Visfatin enhances ICAM-1 and
633 VCAM-1 expression through ROS-dependent NF- κ B activation in endothelial
634 cells. *Biochim. Biophys. Acta* **1783**:886–895; 2008.
- 635 [25] Park, J. K.; Kim, H. K.; Kim, M. J.; Kim, H. J.; Lee, E. N.; Oh, S. O.; Baek, S. Y.; Kim, B. S.;
636 Choi, Y. W.; Yoon, S. Inhibition of TNF- α -induced ROS generation, VCAM-1
637 expression and NF- κ B activation by chloroform extract of aged black garlic in
638 human umbilical vein endothelial cells. *FASEB J.* **24**:960; 2010.
- 639 [26] Lee, C. W.; Lin, W. N.; Lin, C. C.; Luo, S. F.; Wang, J. S.; Pouyssegur, J.; Yang, C. M.
640 Transcriptional regulation of VCAM-1 expression by tumor necrosis factor- α
641 in human tracheal smooth muscle cells: involvement of MAPKs, NF- κ B, p300, 641
642 and histone acetylation. *J. Cell. Physiol.* **207**:174–186; 2006.
- [27] Chiarugi, P. PTPs versus PTKs: the redox side of the coin. *Free. Radic. Res.* **39**: 643
644 353–364; 2005.
- [28] Cominacini, L.; Pasini, A.; Garbin, U.; Evangelista, S.; Crea, A. E.; Tagliacozzi, D.; Nava, C.; 645
646 Davoli, A.; LoCasio, V. Zofenopril inhibits the expression of adhesion molecules on
647 endothelial cells by reducing reactive oxygen species. *Am. J. Hypertens.* **15**:891–895; 2002.
- [29] Mladenka, P.; Zatloukalov, L.; Filipsky, T.; Hrdina, R. Cardiovascular effects of 648
649 flavonoids are not caused only by direct antioxidant activity. *Free Radic. Biol. Med.* 649
650 **49**:963–975; 2010.

UNCORRECTED PROOF