



# EFFECTS OF TAI-CHI CHUAN ON THE CONTROL OF BODY'S CENTRE OF MASS MOTION DURING OBSTACLE-CROSSING IN THE ELDERLY



Tsung-Jung Ho (何宗融)<sup>1,2</sup>, Sheng-Chang Chen(陳聖昌)<sup>3</sup>, Chu-Fen Chang(張祝芬)<sup>3</sup>,

Tsong-Rong Jang(張聰榮)<sup>4</sup>, Jaung-Geng Lin(林昭庚)<sup>1</sup>, Shinn-Zong Lin(林欣榮)<sup>2</sup>, Tung-Wu Lu(呂東武)<sup>3</sup>

<sup>1</sup>College of Chinese Medicine, China Medical University, Taiwan

<sup>2</sup>China Medical University Beigang Hospital, Taiwan

<sup>3</sup>Institute of Biomedical Engineering, National Taiwan University, Taiwan

<sup>4</sup>National Taiwan College of Physical Education, Taichung, Taiwan.

Corresponding author: Tsung-Jung Ho, email: tjho@mail.cmu.edu.tw

Keywords: Taichi, Gait analysis, Inclination angle, Centre of Pressure

## Introduction

Tripping over obstacles is a common problem among the elderly, which often leads to physical injuries. Previous studies have shown the positive effects of Tai Chi Chuan (TCC) exercises on muscle power, flexibility, endurance, dexterity, physical fitness, and balance (Wolf et al., 1996; Li et al., 2004). These effects are beneficial for the control of the body stability and prevention of falls. The aim of this study was to investigate the effects of Tai-Chi Chun and obstacle heights on the inclination angles of the center of mass (COM) relative to the center of foot pressure (COP) during obstacle-crossing.

## Method

Ten TCC practitioners (age: 71±5.3 years, height: 163±6.7 cm, mass: 58.7±6.5 kg, TCC experience: 25±9.9 years) and fifteen healthy controls (age: 72±6 years, height: 160±5.7 cm, mass: 58±10.4 kg) crossed obstacles of heights of 10%, 20% and 30% of their leg lengths while the COM and COP position data were measured using a 3D motion analysis system and forceplates, respectively. COM-COP inclination angles at the instances when the swing toe was above the obstacle for both leading and trailing limb (T1 and T2) were extracted and statistically analyzed using a mixed analysis of variance with one between-subject factor (subject group) and one within-subject factor (obstacle height).

## Results and Discussion

On the anteroposterior (A/P) COM-COP inclination angles, significant group effects were found at T1 (p<0.001) and T2 (p<0.001). Greater anterior COM-COP inclination angles in the control group indicated that they needed more effort to maintain A/P stability. Significant height effects were found at T1 (p<0.001) and T2 (p<0.001) and A/P COM-COP inclination angles were decreased with increasing obstacle height, except for the TCC group at T1. These results suggest that the non-TCC older people tended to keep their COM close to COP to increase the body's stability when the obstacle height increased. However, there were no effects on the mediolateral (M/L) COM-COP inclination angles at T1 and T2.




Figure 1 means and standard deviations of the A(+)/P(-) COM-COP inclination angles at T1 (a) and T2 (b), and these of the M(+)/L(-) angles at T1 (c) and T2 (d). (an arrow: a decreasing trend with increasing obstacle height; an asterisk: a significant difference between groups.)

## Conclusion

The differences of the control strategies adopted during crossing between the TCC and control groups in the sagittal plane may be related to the strengthening effects of one leg standing commonly practiced in the TCC. The findings will be helpful for future studies on kinematics and control strategies adopted during obstacle-crossing.

## References

- Wolf et al. (1996). Reducing frailty and falls in older persons: an investigation of Tai Chi and computerized balance training. *J Am Geriatr Soc.* 44, 489-97.
- Li et al. (2004). Tai Chi: improving functional balance and predicting subsequent falls in older persons. *Med Sci Sports Exerc.* 36, 2046-52

## Acknowledgement

This study was supported by NSC 91-2320-B-244-001.

