

Metric Characterizations of α -Well-Posedness for a System of Mixed Quasivariational-Like Inequalities in Banach Spaces

L. C. Ceng¹ and Y. C. Lin²

¹Professor, Department of Mathematics, Shanghai Normal University, and Scientific Computing Key Laboratory of Shanghai Universities, Shanghai 200234, China. This research was partially supported by the National Science Foundation of China (11071169), Innovation Program of Shanghai Municipal Education Commission (09ZZ133) and Leading Academic Discipline Project of Shanghai Normal University (DZL707). Email: zenglc@hotmail.com

²Corresponding author. Professor, Department of Occupational Safety and Health, College of Public Health, China Medical University, Taichung 404, Taiwan. This research was partially supported by grant NSC99-2115-M-039-001- from the National Science Council of Taiwan. E-mail: yclin@mail.cmu.edu.tw

Abstract. The purpose of this paper is to investigate the problems of the well-posedness for a system of mixed quasivariational-like inequalities in Banach spaces. First, we generalize the concept of α -well-posedness to the system of mixed quasivariational-like inequalities, which includes symmetric quasi-equilibrium problems as a special case. Second, we establish some metric characterizations of α -well-posedness for the system of mixed quasivariational-like inequalities. Under some suitable conditions, we prove that the α -well-posedness is equivalent to the existence and uniqueness of solution for the system of mixed quasivariational-like inequalities. The corresponding concept of α -well-posedness in the generalized sense is also considered for the system of mixed quasivariational-like inequalities having more than one solution. The results presented in this paper generalize and improve some known results in the literature.

Keywords: System of mixed quasivariational-like inequalities; Set-valued mapping; α -well-posedness; α -approximating sequence; Metric characterization.

AMS subject classifications. 49J40; 49K40; 90C31

1. Introduction

The classical notion of well-posedness for a minimization problem (MP) is due to Tykhonov [1], which has already been known as the Tykhonov well-posedness. The so-called Tykhonov well-posedness means the existence and uniqueness of solution, and the convergence of every minimizing sequence toward the unique solution. Taking into account that in many practical situations the solution may not be unique for a minimization problem, one naturally introduced the concept of well-posedness in the generalized sense, which means the existence of minimizers and the convergence of some subsequence of every minimizing sequence toward a minimizer. Obviously, the concept of well-posedness is inspired by numerical methods producing optimizing sequences for optimization problems. In the following years, the well-posedness has received much attention because it plays a crucial role in the stability theory for optimization problems. A large number of results about well-posedness have appeared in the literature; see, e.g., [12,20-26,37], where Refs. [12,20,22,24,37] are for the class of scalar optimization problems, Refs. [21,23,25,39] for the class of vector optimization problems.

On the other hand, the concept of well-posedness has been generalized to other related problems, such as variational inequalities [13,14,16,17,19,26,27,31-33,38,40], Nash equilibrium problems [18,19,35,36], inclusion problems [13,16,34,41] and fixed point problems [6,7,13,16,34]. An initial notion of well-posedness for variational inequalities is due to Lucchetti and Patrone [33]. They introduced the notion of well-posedness for variational inequalities and proved some related results by means of Ekeland's variational principle. Since then, many authors have been devoted to generating the concept of well-posedness from the minimization problem to various variational inequalities. Lignola and Morgan [32] introduced the parametric well-posedness for a family of variational inequalities. Lignola [17] further introduced two concepts of well-posedness and L -well-posedness for quasivariational-like inequalities and derived some metric characterizations of well-posedness. At the same time, Del Prete et al. [31] introduced the concept of α -well-posedness for a class of variational inequalities. Recently, Fang et al. [16] generalized the concept of well-posedness to a class of mixed variational inequalities in Hilbert spaces. They obtained some metric characterizations of its well-posedness and established the links with the well-posedness of inclusion problems and fixed point problems. Furthermore, Ceng and Yao [13] generalized the results of Fang et al. [16] to a class of generalized mixed variational inequalities in Hilbert spaces. Ceng et al. [14] investigated the well-posedness for a class of mixed quasivariational-like inequalities in Banach spaces. For the well-posedness of variational inequalities with functional constraints, we refer to Huang and Yang [26] and Huang et al. [27]. In 2006, Lignola and Morgan [18] presented the notion of α -well-posedness for the Nash equilibrium problem and gave some metric characterizations of this type well-posedness. Petrusel et al. [7] and Fuster et al. [6] discussed the well-posedness of fixed point problems for multivalued mappings in metric spaces.

It is obvious that the equilibrium problem plays a very important role in the establishment of a general mathematical model for a wide range of practical problems, which includes as special cases optimization problems, Nash equilibria problems, fixed point problems, variational inequality problems and complementarity problems (see, e.g., [2,30]), and has been

studied extensively and intensively. It is well known that each equilibrium problem can equivalently be transformed into a minimizing problem by using gap function and some numerical methods have been extended to solve of the equilibrium problem (see, e.g., [29]). This fact motivates the researchers to study the well-posedness for equilibrium problems. Recently, Fang et al. [15] introduced the concepts of parametric well-posedness for equilibrium problems and derived some metric characterizations for these types of well-posedness. For the well-posedness of equilibrium problems with functional constraints, we refer the readers to [28]. In 2009, Long and Huang [8] generalized the concept of α -well-posedness to symmetric quasi-equilibrium problems in Banach spaces, which includes equilibrium problems, Nash equilibrium problems, quasivariational inequalities, variational inequalities, and fixed point problems as special cases. Under some suitable conditions, they established some metric characterizations of α -well-posedness for symmetric quasi-equilibrium problems. Moreover, they gave some examples to illustrate their results. Their results represent the generalization and improvement of some previous known results in the literature, for instance, [13-18]. It is worth pointing out that up to the publication of [8] there are no results concerned with the problems of the well-posedness for symmetric quasi-equilibrium problems in Banach spaces.

In this paper, we consider and study the problems of the well-posedness for a system of mixed quasivariational-like inequalities in Banach spaces. First, we generalize the concept of α -well-posedness to the system of mixed quasivariational-like inequalities, which includes symmetric quasi-equilibrium problems as a special case. Second, some metric characterizations of α -well-posedness for the system of mixed quasivariational-like inequalities are given under very mild conditions. Furthermore, it is also proven that under quite appropriate conditions the α -well-posedness is equivalent to the existence and uniqueness of solution for the system of mixed quasivariational-like inequalities. At the same time, the corresponding concept of α -well-posedness in the generalized sense is also considered for the system of mixed quasivariational-like inequalities having more than one solution. In addition, we give some examples to illustrate our results. The results presented in this paper generalize and improve Long and Huang's results in [8].

2. Preliminaries

Throughout this paper, unless specified otherwise, let X and Y be two real Banach spaces, let their dual spaces be denoted by X^* and Y^* respectively, let the duality pairing between X and X^* and the one between Y and Y^* be denoted by the same $\langle \cdot, \cdot \rangle$. We write $x_n \rightharpoonup x$ to indicate that the sequence $\{x_n\}$ converges weakly to x . $x_n \rightarrow x$ implies that $\{x_n\}$ converges strongly to x . Let $C \subset X$ and $D \subset Y$ be two nonempty closed and convex subsets. Let $S : C \times D \rightarrow 2^C$ and $T : C \times D \rightarrow 2^D$ be two set-valued mappings, let $A : C \times D \rightarrow X^*$, $B : C \times D \rightarrow Y^*$, $\hat{\eta} : C \times C \rightarrow X$ and $\bar{\eta} : D \times D \rightarrow Y$ be four single-valued mappings, and let $f, g : C \times D \rightarrow \mathbf{R}$ be two real-valued functions. Suppose that α is a nonnegative real number and $N = \{1, 2, \dots\}$.

In this paper, we consider the system of mixed quasivariational-like inequalities (in short, SMQVLI), which is to find a point $(x_0, y_0) \in C \times D$ such that

$$\begin{cases} x_0 \in S(x_0, y_0) \text{ and } \langle A(x_0, y_0), \hat{\eta}(x_0, z) \rangle + f(x_0, y_0) - f(z, y_0) \leq 0, & \forall z \in S(x_0, y_0), \\ y_0 \in T(x_0, y_0) \text{ and } \langle B(x_0, y_0), \bar{\eta}(y_0, w) \rangle + g(x_0, y_0) - g(x_0, w) \leq 0, & \forall w \in T(x_0, y_0). \end{cases} \quad (2.1)$$

Remark 2.1. Whenever $A = 0$, $B = 0$, $\hat{\eta} = 0$ and $\bar{\eta} = 0$, the problem (2.1) reduces to the following symmetric quasi-equilibrium problem (in short, SQEP) of finding a point $(x_0, y_0) \in C \times D$ such that

$$\begin{cases} x_0 \in S(x_0, y_0) \text{ and } f(x_0, y_0) \leq f(z, y_0), & \forall z \in S(x_0, y_0), \\ y_0 \in T(x_0, y_0) \text{ and } g(x_0, y_0) \leq g(x_0, w), & \forall w \in T(x_0, y_0). \end{cases} \quad (2.2)$$

This problem was first considered by Noor and Oettli [38], which includes equilibrium problems [2], Nash equilibrium problems [3], quasivariational inequalities [4], variational inequalities [5] and fixed point problems [6,7] as special cases. It is worth mentioning that Noor and Oettli [38] only established the existence of solutions for SQEP (2.2). Subsequently, Long and Huang [8] investigated the α -well-posedness for SQEP (2.2) in Banach spaces.

Denote by Γ the solution set of SMQVLI (2.1). In the following we introduce the notions of α -approximating sequence and α -well-posedness for SMQVLI (2.1).

Definition 2.1. A sequence $\{(x_n, y_n)\} \subset C \times D$ is called an α -approximating sequence for SMQVLI (2.1) if there exists a sequence $\varepsilon_n > 0$ with $\varepsilon_n \rightarrow 0$ such that

$$\begin{cases} d(x_n, S(x_n, y_n)) \leq \varepsilon_n, \text{ i.e., } x_n \in B(S(x_n, y_n), \varepsilon_n), & \forall n \in N, \\ d(y_n, T(x_n, y_n)) \leq \varepsilon_n, \text{ i.e., } y_n \in B(T(x_n, y_n), \varepsilon_n), & \forall n \in N, \end{cases}$$

and

$$\begin{cases} \langle A(x_n, y_n), \hat{\eta}(x_n, z) \rangle + f(x_n, y_n) - f(z, y_n) \leq \varepsilon_n + \frac{\alpha}{2} \|x_n - z\|^2, & \forall z \in S(x_n, y_n), \forall n \in N, \\ \langle B(x_n, y_n), \bar{\eta}(y_n, w) \rangle + g(x_n, y_n) - g(x_n, w) \leq \varepsilon_n + \frac{\alpha}{2} \|y_n - w\|^2, & \forall w \in T(x_n, y_n), \forall n \in N, \end{cases}$$

where $B(S(x, y), \varepsilon)$ denotes the ball of radius ε around $S(x, y)$, that is, the set $\{m \in X : d(S(x, y), m) = \inf_{b \in S(x, y)} \|m - b\| \leq \varepsilon\}$. Whenever $\alpha = 0$, we say that the sequence $\{(x_n, y_n)\}$ is an approximating sequence for SMQVLI (2.1).

We remark that, if $A = 0$, $B = 0$, $\hat{\eta} = 0$ and $\bar{\eta} = 0$, the notions of α -approximating sequence and approximating sequence for SMQVLI (2.1) reduce to the ones of α -approximating sequence and approximating sequence for SQEP (2.2) in [8, Definition 2.1], respectively.

Definition 2.2. SMQVLI (2.1) is said to be α -well-posed if it has a unique solution (x_0, y_0) and every α -approximating sequence $\{(x_n, y_n)\}$ converges strongly to (x_0, y_0) . Whenever $\alpha = 0$, we say that SMQVLI (2.1) is well-posed.

We remark that, if $A = 0$, $B = 0$, $\hat{\eta} = 0$ and $\bar{\eta} = 0$, the notions of α -well-posedness and well-posedness for SMQVLI (2.1) reduce to the ones of α -well-posedness and well-posedness for SQEP (2.2) in [8, Definition 2.2], respectively.

Definition 2.3. SMQVLI (2.1) is said to be α -well-posed in the generalized sense if the solution set F of SMQVLI (2.1) is nonempty and every α -approximating sequence $\{(x_n, y_n)\}$ has a subsequence which converges strongly to some element of F . Whenever $\alpha = 0$, we say that SMQVLI (2.1) is well-posed in the generalized sense.

We remark that, if $A = 0$, $B = 0$, $\hat{\eta} = 0$ and $\bar{\eta} = 0$, the notions of α -well-posedness in the generalized sense and well-posedness in the generalized sense for SMQVLI (2.1) reduce to the ones of α -well-posedness in the generalized sense and well-posedness in the generalized sense for SQEP (2.2) in [8, Definition 2.3], respectively.

In order to investigate the α -well-posedness for SMQVLI (2.1), we need the following definitions.

Definition 2.4 (see [9]). The Painleve-Kuratowski limits of a sequence $\{H_n\} \subset X$ are defined by

$$\liminf_n H_n = \{y \in X : \exists y_n \in H_n, n \in N, \text{ with } \lim_n y_n = y\},$$

$$\limsup_n H_n = \{y \in X : \exists n_k \uparrow +\infty, n_k \in N, \exists y_{n_k} \in H_{n_k}, k \in N, \text{ with } \lim_k y_{n_k} = y\}.$$

Definition 2.5 (see [9]). A set-valued mapping F from a topological space (W, τ) to a topological space (Z, σ) is called

(i) (τ, σ) -closed if for every $x \in K$, for every sequence $\{x_n\}$ τ -converging to x , and for every sequence $\{y_n\}$ σ -converging to a point y , such that $y_n \in F(x_n)$, one has $y \in F(x)$, i.e.,

$$F(x) \supset \limsup_n F(x_n).$$

(ii) (τ, σ) -lower semicontinuous if for every $x \in K$, for every sequence $\{x_n\}$ τ -converging to x , and for every $y \in F(x)$, there exists a sequence $\{y_n\}$ σ -converging to y , such that $y_n \in F(x_n)$ for n sufficiently large, i.e.,

$$F(x) \subset \liminf_n F(x_n).$$

(iii) (τ, σ) -subcontinuous on K , if for every sequence $\{x_n\}$ τ -converging in K , every sequence $\{y_n\}$, such that $y_n \in F(x_n)$, has a σ -convergent subsequence.

Definition 2.6 (see [9]). Let V be a nonempty subset of X . The measure of non-compactness μ of the set V is defined by

$$\mu(V) = \inf\{\varepsilon > 0 : V \subset \bigcup_{i=1}^n V_i, \text{ diam} V_i < \varepsilon, \quad i = 1, 2, \dots, n\},$$

where diam means the diameter of a set.

Definition 2.7 (see [9]). Let (X, d) be a metric space and Let U, V be nonempty subsets of X . The Hausdorff metric $\mathcal{H}(\cdot, \cdot)$ between U and V is defined by

$$\mathcal{H}(U, V) = \max\{e(U, V), e(V, U)\},$$

where $e(U, V) = \sup_{u \in U} d(u, V)$ with $d(u, V) = \inf_{v \in V} \|u - v\|$. Let $\{U_n\}$ be a sequence of nonempty subsets of X . We say that U_n converges to U in the sense of Hausdorff metric if $\mathcal{H}(U_n, U) \rightarrow 0$. It is easy to see that $e(U_n, U) \rightarrow 0$ iff $d(u_n, U) \rightarrow 0$ for all section $u_n \in U_n$. For more details on this topic, we refer the readers to [9].

Now, we prove the following lemma.

Lemma 2.1. Suppose that set-valued mappings S and T are nonempty convex-valued, the function $f(\cdot, y)$ is convex on C for any $y \in D$, and the function $g(x, \cdot)$ is convex on D for any $x \in C$. Then $(x_0, y_0) \in \Gamma$ if and only if the following two conditions hold:

$$x_0 \in S(x_0, y_0), \langle A(x_0, y_0), \hat{\eta}(x_0, z) \rangle + f(x_0, y_0) - f(z, y_0) \leq \frac{\alpha}{2} \|x_0 - z\|^2, \quad \forall z \in S(x_0, y_0), \quad (2.3)$$

$$y_0 \in T(x_0, y_0), \langle B(x_0, y_0), \bar{\eta}(y_0, w) \rangle + g(x_0, y_0) - g(x_0, w) \leq \frac{\alpha}{2} \|y_0 - w\|^2, \quad \forall w \in T(x_0, y_0), \quad (2.4)$$

where both $\hat{\eta} : C \times C \rightarrow X$ and $\bar{\eta} : D \times D \rightarrow Y$ are affine in the second variable such that $\hat{\eta}(x, x) = 0$ and $\bar{\eta}(y, y) = 0$ for all $(x, y) \in C \times D$.

Proof. The necessity is obvious. For the sufficiency, suppose that (2.3) and (2.4) hold. Now let us show that $(x_0, y_0) \in \Gamma$. Indeed, let $z \in S(x_0, y_0)$ and for any $t \in [0, 1]$, $z_t = tz + (1 - t)x_0$. Since $S(x_0, y_0)$ is convex, $z_t \in S(x_0, y_0)$ and so

$$\langle A(x_0, y_0), \hat{\eta}(x_0, z_t) \rangle + f(x_0, y_0) - f(z_t, y_0) \leq \frac{\alpha}{2} \|x_0 - z_t\|^2, \quad \forall t \in (0, 1].$$

Also, since $f(\cdot, y)$ is convex for any $y \in D$ and $\hat{\eta} : C \times C \rightarrow X$ is affine in the second variable with $\hat{\eta}(x, x) = 0$, $\forall x \in X$, we have

$$\begin{aligned} & t\{\langle A(x_0, y_0), \hat{\eta}(x_0, z) \rangle + f(x_0, y_0) - f(z, y_0)\} \\ &= \langle A(x_0, y_0), t\hat{\eta}(x_0, z) + (1 - t)\hat{\eta}(x_0, x_0) \rangle + f(x_0, y_0) - (tf(z, y_0) + (1 - t)f(x_0, y_0)) \\ &\leq \langle A(x_0, y_0), \hat{\eta}(x_0, z_t) \rangle + f(x_0, y_0) - f(z_t, y_0) \\ &\leq \frac{\alpha}{2} \|x_0 - z_t\|^2 \\ &= \frac{\alpha}{2} t^2 \|x_0 - z\|^2, \quad \forall t \in (0, 1]. \end{aligned}$$

Thus, dividing by t in the above inequality, we have

$$\langle A(x_0, y_0), \hat{\eta}(x_0, z) \rangle + f(x_0, y_0) - f(z, y_0) \leq \frac{\alpha}{2} t \|x_0 - z\|^2, \quad \forall t \in (0, 1], \quad z \in S(x_0, y_0). \quad (2.5)$$

By the similar argument,

$$\langle B(x_0, y_0), \bar{\eta}(y_0, w) \rangle + g(x_0, y_0) - g(x_0, w) \leq \frac{\alpha}{2} t \|y_0 - w\|^2, \quad \forall t \in (0, 1], \quad w \in T(x_0, y_0). \quad (2.6)$$

The combination of (2.5) and (2.6) implies, for t tending to zero, that (x_0, y_0) is a solution of SMQVLI (2.1). This completes the proof. \square

Corollary 2.1 (i.e., [8, Lemma 2.1]). Suppose that set-valued mappings S and T are nonempty convex-valued, the function $f(\cdot, y)$ is convex on C for any $y \in D$, and the function $g(x, \cdot)$ is convex on D for any $x \in C$. Then (x_0, y_0) solves SQEP (2.2) if and only if the following two conditions hold:

$$x_0 \in S(x_0, y_0), \quad f(x_0, y_0) \leq f(z, y_0) + \frac{\alpha}{2} \|x_0 - z\|^2, \quad \forall z \in S(x_0, y_0),$$

$$y_0 \in T(x_0, y_0), \quad g(x_0, y_0) \leq g(x_0, w) + \frac{\alpha}{2} \|y_0 - w\|^2, \quad \forall w \in T(x_0, y_0).$$

Proof. Put $A = 0$, $B = 0$, $\hat{\eta} = 0$ and $\bar{\eta} = 0$ in Lemma 2.1. Then, utilizing Lemma 2.1 we get the desired result. \square

3. Metric Characterizations of α -Well-Posedness for SMQVLI

In this section, we shall investigate some metric characterizations of α -well-posedness for SMQVLI (2.1).

For any $\varepsilon > 0$, the α -approximating solution set of SMQVLI (2.1) is defined by

$$\begin{aligned} \Omega_\alpha(\varepsilon) = \{ & (x_0, y_0) \in C \times D : \\ & x_0 \in B(S(x_0, y_0), \varepsilon), \quad \langle A(x_0, y_0), \hat{\eta}(x_0, z) \rangle + f(x_0, y_0) - f(z, y_0) \leq \varepsilon + \frac{\alpha}{2} \|x_0 - z\|^2, \quad \forall z \in S(x_0, y_0), \\ & y_0 \in B(T(x_0, y_0), \varepsilon), \quad \langle B(x_0, y_0), \bar{\eta}(y_0, w) \rangle + g(x_0, y_0) - g(x_0, w) \leq \varepsilon + \frac{\alpha}{2} \|y_0 - w\|^2, \quad \forall w \in T(x_0, y_0) \}. \end{aligned}$$

Theorem 3.1. SMQVLI (2.1) is α -well-posed if and only if the solution set Γ of SMQVLI (2.1) is nonempty and

$$\lim_{\varepsilon \rightarrow 0} \text{diam} \Omega_\alpha(\varepsilon) = 0. \quad (3.1)$$

Proof. Suppose that SMQVLI (2.1) is α -well-posed. Then, Γ is a singleton, and $\Omega_\alpha(\varepsilon) \neq \emptyset$ for any $\varepsilon > 0$, since $\Gamma \subset \Omega_\alpha(\varepsilon)$. Suppose by contraction that

$$\lim_{\varepsilon \rightarrow 0} \text{diam} \Omega_\alpha(\varepsilon) > \beta > 0.$$

Then there exists $\varepsilon_n > 0$ with $\varepsilon_n \rightarrow 0$, and $(x_n, y_n), (\bar{x}_n, \bar{y}_n) \in \Omega_\alpha(\varepsilon_n)$ such that

$$\|(x_n, y_n) - (\bar{x}_n, \bar{y}_n)\| > \beta, \quad \forall n \in \mathbb{N},$$

where the norm $\|\cdot\|$ in the product space $X \times Y$ is defined as follows

$$\|(u, v) - (\bar{u}, \bar{v})\| = \sqrt{\|u - \bar{u}\|^2 + \|v - \bar{v}\|^2}, \quad \forall (u, v), (\bar{u}, \bar{v}) \in X \times Y. \quad (3.2)$$

(It is not difficult to verify that $X \times Y$ is a Banach space in terms of the last norm.)

Since $(x_n, y_n), (\bar{x}_n, \bar{y}_n) \in \Omega_\alpha(\varepsilon_n)$ and SMQVLI (2.1) is α -well-posed, the sequences $\{(x_n, y_n)\}$ and $\{(\bar{x}_n, \bar{y}_n)\}$, which are both α -approximating sequences for SMQVLI (2.1), converge strongly to the unique solution (x_0, y_0) , and this leads to a contraction. Therefore, (3.1) holds.

Conversely, let (3.1) hold and $\{(x_n, y_n)\} \subset C \times D$ be any α -approximating sequence for SMQVLI (2.1). Then, there exists a sequence $\varepsilon_n > 0$ with $\varepsilon_n \rightarrow 0$ such that

$$d(x_n, S(x_n, y_n)) \leq \varepsilon_n, \text{ and } \langle A(x_n, y_n), \hat{\eta}(x_n, z) \rangle + f(x_n, y_n) - f(z, y_n) \leq \varepsilon_n + \frac{\alpha}{2} \|x_n - z\|^2, \forall z \in S(x_n, y_n),$$

$$d(y_n, T(x_n, y_n)) \leq \varepsilon_n, \text{ and } \langle B(x_n, y_n), \bar{\eta}(y_n, w) \rangle + g(x_n, y_n) - g(x_n, w) \leq \varepsilon_n + \frac{\alpha}{2} \|y_n - w\|^2, \forall w \in T(x_n, y_n).$$

This implies that $\{(x_n, y_n)\} \subset \Omega_\alpha(\varepsilon_n)$, $\forall n \in N$. Since the solution set Γ of SMQVLI (2.1) is nonempty, we can take two elements in Γ arbitrarily, denoted by (x_0, y_0) and (\bar{x}_0, \bar{y}_0) , respectively. Note that $\Gamma \subset \Omega_\alpha(\varepsilon)$ for all $\varepsilon > 0$. Hence both (x_0, y_0) and (\bar{x}_0, \bar{y}_0) lie in $\Omega_\alpha(\varepsilon_n)$ for all $n \geq 1$. This fact together with (3.1) yields

$$\|(x_n, y_n) - (x_0, y_0)\| \leq \text{diam}\Omega_\alpha(\varepsilon_n) \rightarrow 0 \text{ and } \|(x_n, y_n) - (\bar{x}_0, \bar{y}_0)\| \leq \text{diam}\Omega_\alpha(\varepsilon_n) \rightarrow 0. \quad (3.3)$$

Utilizing (3.3) and the uniqueness of the limit we conclude that $(x_0, y_0) = (\bar{x}_0, \bar{y}_0)$. This means that Γ is a singleton. Thus, it is known that SMQVLI (2.1) has the unique solution (x_0, y_0) and $\{(x_n, y_n)\}$ converges strongly to (x_0, y_0) . This shows that SMQVLI (2.1) is α -well-posed. This completes the proof. \square

Corollary 3.1 (i.e., [8, Theorem 3.1]). SQEP (2.2) is α -well-posed if and only if the solution set Γ of SQEP (2.2) is nonempty and

$$\lim_{\varepsilon \rightarrow 0} \text{diam}M_\varepsilon = 0,$$

where

$$M_\varepsilon = \{(x_0, y_0) \in C \times D : \begin{aligned} &x_0 \in B(S(x_0, y_0), \varepsilon), f(x_0, y_0) - f(z, y_0) \leq \varepsilon + \frac{\alpha}{2} \|x_0 - z\|^2, \forall z \in S(x_0, y_0), \\ &y_0 \in B(T(x_0, y_0), \varepsilon), g(x_0, y_0) - g(x_0, w) \leq \varepsilon + \frac{\alpha}{2} \|y_0 - w\|^2, \forall w \in T(x_0, y_0) \end{aligned}\}$$

Proof. Put $A = 0$, $B = 0$, $\hat{\eta} = 0$ and $\bar{\eta} = 0$ in Theorem 3.1. Then, utilizing Theorem 3.1 we get the desired result. \square

In the sequel the following concept will be needed to apply to our main results.

Definition 3.1. Let C be a nonempty, closed convex subset of X . A single-valued mapping $\eta : C \times C \rightarrow X$ is said to be Lipschitz continuous if there exists a constant $\lambda > 0$ such that

$$\|\eta(x, y)\| \leq \lambda \|x - y\|, \quad \forall x, y \in C.$$

We remark that, whenever $X = H$ a Hilbert space and $C = K$ a nonempty closed convex subset of H , the Lipschitz continuous mapping $\eta : K \times K \rightarrow H$ has been introduced and considered in Ansari and Yao [10]. In their main result for the existence of solutions and convergence of iterative algorithm (i.e., [10, Theorem 3.1]), the Lipschitz continuous mapping $\eta : K \times K \rightarrow H$ satisfies the following conditions:

- (a) $\eta(x, y) + \eta(y, x) = 0$ for all $x, y \in K$,
- (b) $\eta(x, y) = \eta(x, z) + \eta(z, y)$ for all $x, y, z \in K$,
- (c) $\eta(\cdot, \cdot)$ is affine in the first variable,
- (d) for each fixed $y \in K$, $x \mapsto \eta(y, x)$ is sequentially continuous from the weak topology to the weak topology (in short, (w, w) -continuous).

Inspired by the above restrictions imposed on the Lipschitz continuous mapping η , we give the following theorem.

Theorem 3.2. Assume that the following conditions hold:

- (i) set-valued mappings S and T are nonempty convex-valued, (s, w) -closed, (s, s) -lower semicontinuous and (s, w) -subcontinuous on $C \times D$;
- (ii) single-valued mappings A and B are (s, w^*) -continuous on $C \times D$;
- (iii) single-valued mappings $\hat{\eta}$ and $\bar{\eta}$ are Lipschitz continuous with constants $\hat{\lambda}$ and $\bar{\lambda}$ respectively, such that
 - (a) $\hat{\eta}(x_1, x_3) = \hat{\eta}(x_1, x_2) + \hat{\eta}(x_2, x_3)$ for all $x_1, x_2, x_3 \in C$ and $\bar{\eta}(y_1, y_3) = \bar{\eta}(y_1, y_2) + \bar{\eta}(y_2, y_3)$ for all $y_1, y_2, y_3 \in D$,
 - (b) $\hat{\eta}(\cdot, \cdot)$ and $\bar{\eta}(\cdot, \cdot)$ both are affine in the second variable;
 - (iv) functions f and g are continuous on $C \times D$;
 - (v) for any $y \in D$, the function $f(\cdot, y)$ is convex on C ; for any $x \in C$, the function $g(x, \cdot)$ is convex on D .

Then, SMQVLI (2.1) is α -well-posed if and only if

$$\Omega_\alpha(\varepsilon) \neq \emptyset, \quad \forall \varepsilon > 0, \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0} \text{diam} \Omega_\alpha(\varepsilon) = 0. \quad (3.4)$$

Proof. First, utilizing condition (iii) (a), we can readily obtain that

$$\hat{\eta}(x_1, x_1) = 0, \quad \text{and} \quad \hat{\eta}(x_1, x_2) = -\hat{\eta}(x_2, x_1), \quad \forall x_1, x_2 \in C;$$

$$\bar{\eta}(y_1, y_1) = 0, \quad \text{and} \quad \bar{\eta}(y_1, y_2) = -\bar{\eta}(y_2, y_1), \quad \forall y_1, y_2 \in D.$$

The necessity has been proved in Theorem 3.1. For the sufficiency, let condition (3.4) hold. Let $\{(x_n, y_n)\} \subset C \times D$ be any α -approximating sequence for SMQVLI (2.1). Now let us show that Γ is a singleton and $\{(x_n, y_n)\}$ converges strongly to the unique element of Γ . As a matter of fact, since $\{(x_n, y_n)\}$ be α -approximating sequence for SMQVLI (2.1), there exists a sequence $\varepsilon_n > 0$ with $\varepsilon_n \rightarrow 0$ such that

$$d(x_n, S(x_n, y_n)) \leq \varepsilon_n, \quad \text{and} \quad \langle A(x_n, y_n), \hat{\eta}(x_n, z) \rangle + f(x_n, y_n) - f(z, y_n) \leq \varepsilon_n + \frac{\alpha}{2} \|x_n - z\|^2, \quad \forall z \in S(x_n, y_n),$$

$d(y_n, T(x_n, y_n)) \leq \varepsilon_n$, and $\langle B(x_n, y_n), \bar{\eta}(y_n, w) \rangle + g(x_n, y_n) - g(x_n, w) \leq \varepsilon_n + \frac{\alpha}{2} \|y_n - w\|^2$, $\forall w \in T(x_n, y_n)$.

This means $\{(x_n, y_n)\} \subset \Omega_\alpha(\varepsilon_n)$, $\forall n \in N$. It follows from (3.4) that $\{(x_n, y_n)\}$ is a Cauchy sequence in Banach space $(X \times Y, \|\cdot\|)$ and hence converges strongly to a point $(x_0, y_0) \in X \times Y$. By the definition of the norm $\|\cdot\|$ in Banach space $(X \times Y, \|\cdot\|)$, we deduce that

$$\|x_n - x_0\| \leq \sqrt{\|x_n - x_0\|^2 + \|y_n - y_0\|^2} = \|(x_n, y_n) - (x_0, y_0)\| \rightarrow 0,$$

$$\|y_n - y_0\| \leq \sqrt{\|x_n - x_0\|^2 + \|y_n - y_0\|^2} = \|(x_n, y_n) - (x_0, y_0)\| \rightarrow 0.$$

On account of the closedness of C and D we conclude from $\{x_n\} \subset C$ and $\{y_n\} \subset D$ that $x_n \rightarrow x_0 \in C$ and $y_n \rightarrow y_0 \in D$. In order to show $(x_0, y_0) \in \Gamma$, we start to prove that

$$d(x_0, S(x_0, y_0)) \leq \liminf_n d(x_n, S(x_n, y_n)) = \lim_n \varepsilon_n = 0.$$

Indeed, suppose that the left inequality does not hold. Then there exists a positive number γ such that

$$\liminf_n d(x_n, S(x_n, y_n)) < \gamma < d(x_0, S(x_0, y_0)),$$

or equivalently, there exist an increasing sequence $\{n_k\}$ and a sequence $\{z_k\}$, $z_k \in S(x_{n_k}, y_{n_k})$, $\forall k \in N$ such that

$$\|x_{n_k} - z_k\| < \gamma, \quad \forall k \in N.$$

Since the set-valued mapping S is (s, w) -closed and (s, w) -subcontinuous, the sequence $\{z_k\}$ has a subsequence, denoted still by $\{z_k\}$, converging weakly to a point $z_0 \in S(x_0, y_0)$. From the weak lower semicontinuity of the norm, it follows that

$$\gamma < d(x_0, S(x_0, y_0)) \leq \|x_0 - z_0\| \leq \liminf_k \|x_{n_k} - z_k\| < \gamma,$$

which leads to a contradiction. Thus we must have $d(x_0, S(x_0, y_0)) = 0$ and hence $x_0 \in S(x_0, y_0)$. Similarly, we can prove $y_0 \in T(x_0, y_0)$.

To complete the proof, we take a point $z \in S(x_0, y_0)$ arbitrarily. Since S is (s, s) -lower semicontinuous, there exists a sequence $\{z_n\}$ converging strongly to z , such that $z_n \in S(x_n, y_n)$ for n sufficiently large. Furthermore, utilizing condition (iii) (a) and the Lipschitz continuity of $\hat{\eta}$ we deduce that

$$\begin{aligned} \|\hat{\eta}(x_n, z_n) - \hat{\eta}(x_0, z)\| &= \|\hat{\eta}(x_n, z_n) - \hat{\eta}(x_0, z_n) + \hat{\eta}(x_0, z_n) - \hat{\eta}(x_0, z)\| \\ &\leq \|\hat{\eta}(x_n, z_n) - \hat{\eta}(x_0, z_n)\| + \|\hat{\eta}(x_0, z_n) - \hat{\eta}(x_0, z)\| \\ &= \|\hat{\eta}(x_n, z_n) + \hat{\eta}(z_n, x_0)\| + \|\hat{\eta}(x_0, z_n) + \hat{\eta}(z, x_0)\| \\ &= \|\hat{\eta}(x_n, x_0)\| + \|\hat{\eta}(z, z_n)\| \\ &\leq \hat{\lambda}(\|x_n - x_0\| + \|z_n - z\|) \rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned}$$

Since A is (s, w^*) -continuous, it is known that $A(x_n, y_n)$ converges weak*ly to $A(x_0, y_0)$, that is, for each $x \in X$, the real sequence $\{\langle A(x_n, y_n), x \rangle\}$ converges to the real number $\langle A(x_0, y_0), x \rangle$. This implies that $\{\langle A(x_n, y_n), x \rangle\}$ is a bounded sequence of real numbers for each $x \in X$. Thus

$\{A(x_n, y_n)\}$ is bounded in the norm topology according to the uniform boundedness principle [11], that is, $\sup_{n \geq 1} \|A(x_n, y_n)\| < \infty$.

Now observe that

$$\begin{aligned} & |\langle A(x_n, y_n), \hat{\eta}(x_n, z_n) \rangle - \langle A(x_0, y_0), \hat{\eta}(x_0, z) \rangle| \\ &= |\langle A(x_n, y_n), \hat{\eta}(x_n, z_n) \rangle - \langle A(x_n, y_n), \hat{\eta}(x_0, z) \rangle + \langle A(x_n, y_n), \hat{\eta}(x_0, z) \rangle - \langle A(x_0, y_0), \hat{\eta}(x_0, z) \rangle| \\ &\leq |\langle A(x_n, y_n), \hat{\eta}(x_n, z_n) - \hat{\eta}(x_0, z) \rangle| + |\langle A(x_n, y_n) - A(x_0, y_0), \hat{\eta}(x_0, z) \rangle| \\ &\leq \|A(x_n, y_n)\| \|\hat{\eta}(x_n, z_n) - \hat{\eta}(x_0, z)\| + |\langle A(x_n, y_n) - A(x_0, y_0), \hat{\eta}(x_0, z) \rangle| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned}$$

Consequently, it follows from condition (iv) that

$$\begin{aligned} \langle A(x_0, y_0), \hat{\eta}(x_0, z) \rangle + f(x_0, y_0) - f(z, y_0) &= \lim_n \{ \langle A(x_n, y_n), \hat{\eta}(x_n, z_n) \rangle + f(x_n, y_n) - f(z_n, y_n) \} \\ &\leq \lim_n (\varepsilon_n + \frac{\alpha}{2} \|x_n - z_n\|^2) \\ &= \frac{\alpha}{2} \|x_0 - z\|^2, \end{aligned}$$

for all $z \in S(x_0, y_0)$. Analogously, we have

$$\langle B(x_0, y_0), \bar{\eta}(y_0, w) \rangle + g(x_0, y_0) - g(x_0, w) \leq \frac{\alpha}{2} \|y_0 - w\|^2, \quad \forall w \in T(x_0, y_0).$$

It follows from Lemma 2.1 that $(x_0, y_0) \in \Gamma$. Therefore, SMQVLI (2.1) is α -well-posed. This completes the proof. \square

Corollary 3.2 (i.e., [8, Theorem 3.2]). Assume that the following conditions hold:

- (i) set-valued mappings S and T are nonempty convex-valued, (s, w) -closed, (s, s) -lower semicontinuous and (s, w) -subcontinuous on $C \times D$;
- (ii) functions f and g are continuous on $C \times D$;
- (iii) for any $y \in D$, the function $f(\cdot, y)$ is convex on C ; for any $x \in C$, the function $g(x, \cdot)$ is convex on D .

Then, SQEP (2.2) is α -well-posed if and only if

$$M_\varepsilon \neq \emptyset, \quad \forall \varepsilon > 0, \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0} \text{diam} M_\varepsilon = 0.$$

To illustrate Theorem 3.2, we give the following two examples.

Example 3.1. Let $X = Y = \mathbf{R}$ and $C = D = \mathbf{R}^+ (= [0, +\infty))$. Let $S(x, y) = [0, x]$, $T(x, y) = [0, y]$, $A(x, y) = B(x, y) = -(x - y)^2$, $\hat{\eta}(x, z) = x - z$, $\bar{\eta}(y, w) = y - w$, $f(x, y) = x^2 - y^2$ and $g(x, y) = y^2 - x^2$ for all $x, z \in C$ and $y, w \in D$. Obviously, the conditions (i)-(v) of Theorem 3.2 are satisfied. Note that

$$\begin{aligned} & \{(x, y) \in C \times D : d(x, S(x, y)) \leq \varepsilon, \langle A(x, y), \hat{\eta}(x, z) \rangle + f(x, y) - f(z, y) \leq \varepsilon + \frac{\alpha}{2} \|x - z\|^2, \\ & \quad \forall z \in S(x, y)\} \\ &= \{(x, y) \in C \times D : d(x, S(x, y)) \leq \varepsilon, -(x - y)^2(x - z) + x^2 - z^2 \leq \varepsilon + \frac{\alpha}{2}(x - z)^2, \forall z \in S(x, y)\} \\ &= \{(x, y) \in C \times D : d(x, S(x, y)) \leq \varepsilon, -(x - y)^2(x - z) - (2 + \alpha)(z - \frac{\alpha x}{2 + \alpha})^2 + \frac{4}{2 + \alpha}x^2 - 2\varepsilon \leq 0, \\ & \quad \forall z \in S(x, y)\} \\ &= [0, \sqrt{\frac{(2 + \alpha)\varepsilon}{2}}] \times \mathbf{R}, \end{aligned}$$

and

$$\begin{aligned}
& \{(x, y) \in C \times D : d(y, T(x, y)) \leq \varepsilon, \langle B(x, y), \bar{\eta}(y, w) \rangle + g(x, y) - g(x, w) \leq \varepsilon + \frac{\alpha}{2} \|y - w\|^2, \\
& \quad \forall w \in T(x, y)\} \\
&= \{(x, y) \in C \times D : d(y, T(x, y)) \leq \varepsilon, -(x - y)^2(y - w) + y^2 - w^2 \leq \varepsilon + \frac{\alpha}{2}(y - w)^2, \forall w \in T(x, y)\} \\
&= \{(x, y) \in C \times D : d(y, T(x, y)) \leq \varepsilon, -(x - y)^2(y - w) - (2 + \alpha)(w - \frac{\alpha y}{2 + \alpha})^2 + \frac{4}{2 + \alpha}y^2 - 2\varepsilon \leq 0, \\
& \quad \forall w \in T(x, y)\} \\
&= \mathbf{R} \times [0, \sqrt{\frac{(2 + \alpha)\varepsilon}{2}}].
\end{aligned}$$

It follows that

$$\Omega_\alpha(\varepsilon) = [0, \sqrt{\frac{(2 + \alpha)\varepsilon}{2}}] \times [0, \sqrt{\frac{(2 + \alpha)\varepsilon}{2}}]$$

and so $\text{diam}\Omega_\alpha \rightarrow 0$ as $\varepsilon \rightarrow 0$. By Theorem 3.2, SMQVLI (2.1) is α -well-posed.

Example 3.2. Let $X = Y = \mathbf{R}$ and $C = D = \mathbf{R}^+$ ($= [0, +\infty)$). Let $S(x, y) = [0, x]$, $T(x, y) = [0, y]$, $A(x, y) = B(x, y) = -(x - y)^2$, $\hat{\eta}(x, z) = x - z$, $\bar{\eta}(y, w) = y - w$ and $f(x, y) = g(x, y) = -xy$ for all $x, z \in C$ and $y, w \in D$. It is easy to see that the conditions (i)-(v) of Theorem 3.2 are satisfied, and $\Omega_\alpha(\varepsilon) = [0, +\infty) \times [0, +\infty)$. But, SMQVLI (2.1) is not α -well-posed, since $\text{diam}\Omega_\alpha(\varepsilon) \not\rightarrow 0$ as $\varepsilon \rightarrow 0$.

Whenever $\alpha = 0$, we have the following result.

Theorem 3.3. Assume that the following conditions hold:

- (i) set-valued mappings S and T are nonempty convex-valued, (s, w) -closed, (s, s) -lower semicontinuous and (s, w) -subcontinuous on $C \times D$;
- (ii) single-valued mappings A and B are (s, w^*) -continuous on $C \times D$;
- (iii) single-valued mappings $\hat{\eta}$ and $\bar{\eta}$ are Lipschitz continuous with constants $\hat{\lambda}$ and $\bar{\lambda}$ respectively, such that for all $x_1, x_2, x_3 \in C$ and $y_1, y_2, y_3 \in D$

$$\hat{\eta}(x_1, x_3) = \hat{\eta}(x_1, x_2) + \hat{\eta}(x_2, x_3) \quad \text{and} \quad \bar{\eta}(y_1, y_3) = \bar{\eta}(y_1, y_2) + \bar{\eta}(y_2, y_3);$$

- (iv) functions f and g are continuous on $C \times D$.

Then, SMQVLI (2.1) is well-posed if and only if

$$\Omega_0(\varepsilon) \neq \emptyset, \quad \forall \varepsilon > 0, \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0} \text{diam}\Omega_0(\varepsilon) = 0.$$

Corollary 3.3 (i.e., [8, Corollary 3.1]). Assume that the following conditions hold:

- (i) set-valued mappings S and T are nonempty convex-valued, (s, w) -closed, (s, s) -lower semicontinuous and (s, w) -subcontinuous on $C \times D$;
- (ii) functions f and g are continuous on $C \times D$.

Then, SQEP (2.2) is well-posed if and only if

$$M_\varepsilon \neq \emptyset, \quad \forall \varepsilon > 0, \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0} \text{diam}M_\varepsilon = 0.$$

The following theorem shows that under some suitable conditions, the α -well-posedness of SMQVLI (2.1) is equivalent to the existence and uniqueness of its solutions.

Theorem 3.4. Let X and Y be two finite-dimensional spaces. Suppose that the following conditions hold:

- (i) set-valued mappings S and T are nonempty convex-valued, closed, lower semicontinuous and subcontinuous on $C \times D$;
- (ii) single-valued mappings A and B are continuous on $C \times D$;
- (iii) single-valued mappings $\hat{\eta}$ and $\bar{\eta}$ are Lipschitz continuous with constants $\hat{\lambda}$ and $\bar{\lambda}$ respectively, such that
 - (a) $\hat{\eta}(x_1, x_3) = \hat{\eta}(x_1, x_2) + \hat{\eta}(x_2, x_3)$ for all $x_1, x_2, x_3 \in C$ and $\bar{\eta}(y_1, y_3) = \bar{\eta}(y_1, y_2) + \bar{\eta}(y_2, y_3)$ for all $y_1, y_2, y_3 \in D$,
 - (b) $\hat{\eta}(\cdot, \cdot)$ and $\bar{\eta}(\cdot, \cdot)$ both are affine in the second variable;
 - (iv) the functions f and g are continuous on $C \times D$;
 - (v) for any $y \in D$, the function $f(\cdot, y)$ is convex on C ; for any $x \in C$, the function $g(x, \cdot)$ is convex on D ;
 - (vi) $\Omega_\alpha(\varepsilon)$ is nonempty bounded for some $\varepsilon > 0$.

Then, SMQVLI (2.1) is α -well-posed if and only if SMQVLI (2.1) has a unique solution.

Proof. The necessary of the theorem is obvious. In order to show the sufficiency, let (x_0, y_0) be the unique solution of SMQVLI (2.1) and $\{(x_n, y_n)\}$ be any α -approximating sequence for SMQVLI (2.1). Then there exists a sequence $\varepsilon_n > 0$ with $\varepsilon_n \rightarrow 0$ such that,

$$d(x_n, S(x_n, y_n)) \leq \varepsilon_n, \text{ and } \langle A(x_n, y_n), \hat{\eta}(x_n, z) \rangle + f(x_n, y_n) - f(z, y_n) \leq \varepsilon_n + \frac{\alpha}{2} \|x_n - z\|^2, \forall z \in S(x_n, y_n),$$

$$d(y_n, T(x_n, y_n)) \leq \varepsilon_n, \text{ and } \langle B(x_n, y_n), \bar{\eta}(y_n, w) \rangle + g(x_n, y_n) - g(x_n, w) \leq \varepsilon_n + \frac{\alpha}{2} \|y_n - w\|^2, \forall w \in T(x_n, y_n).$$

which means $\{(x_n, y_n)\} \subset \Omega_\alpha(\varepsilon_n)$, $\forall n \in N$. Let $\varepsilon > 0$ be such that $\Omega_\alpha(\varepsilon)$ is nonempty bounded. Then there exists $n_0 \in N$ such that $\{(x_n, y_n)\} \subset \Omega_\alpha(\varepsilon_n) \subset \Omega_\alpha(\varepsilon)$ for all $n \geq n_0$. Thus, $\{(x_n, y_n)\}$ is bounded and so the sequence $\{(x_n, y_n)\}$ has a subsequence $\{(x_{n_k}, y_{n_k})\}$ which converges to (\tilde{x}, \tilde{y}) . Reasoning as in Theorem 3.2, one can prove that (\tilde{x}, \tilde{y}) solves SMQVLI (2.1). The uniqueness of the solution implies that $(x_0, y_0) = (\tilde{x}, \tilde{y})$, and so the whole sequence $\{(x_n, y_n)\}$ converges to (x_0, y_0) . Thus, SMQVLI (2.1) is α -well-posed. This completes the proof. \square

Example 3.3. Let $X = Y = \mathbf{R}$ and $C = D = \mathbf{R}^+$ ($= [0, +\infty)$). Let $S(x, y) = [0, x]$, $T(x, y) = [0, y]$, $A(x, y) = B(x, y) = -(x - y)^2$, $\hat{\eta}(x, z) = x - z$, $\bar{\eta}(y, w) = y - w$, $f(x, y) = x^2 - y^2$ and $g(x, y) = y^2 - x^2$ for all $x, z \in C$ and $y, w \in D$. Clearly, the conditions (i)-(vi) of Theorem 3.3 are satisfied, and SMQVLI (2.1) has a unique solution $(x_0, y_0) = (0, 0)$. By Theorem 3.3, SMQVLI (2.1) is α -well-posed.

Corollary 3.4 (i.e., [8, Theorem 3.3]) Let X and Y be two finite-dimensional spaces. Suppose that the following conditions hold:

- (i) set-valued mappings S and T are nonempty convex-valued, closed, lower semicontinuous and subcontinuous on $C \times D$;
 - (ii) the functions f and g are continuous on $C \times D$;
 - (iii) for any $y \in D$, the function $f(\cdot, y)$ is convex on C ; for any $x \in C$, the function $g(x, \cdot)$ is convex on D ;
 - (iv) M_ε is nonempty bounded for some $\varepsilon > 0$.
- Then, SQEP (2.2) is α -well-posed if and only if SQEP (2.2) has a unique solution.

4. Metric Characterizations of α -Well-Posedness in the Generalized Sense for SMQVLI

In this section, we derive some metric characterizations of α -well-posedness in the generalized sense for SMQVLI (2.1) by considering the non-compactness of approximate solution set.

Theorem 4.1. SMQVLI (2.1) is α -well-posed in the generalized sense if and only if the solution set Γ of SMQVLI (2.1) is nonempty compact and

$$e(\Omega_\alpha(\varepsilon), \Gamma) \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0. \quad (4.1)$$

Proof. Suppose that SMQVLI (2.1) is α -well-posed in the generalized sense. Then Γ is nonempty. To show the compactness of Γ , let $\{(x_n, y_n)\} \subset \Gamma$. Clearly, if $\{(x_n, y_n)\}$ is an approximation sequence of SMQVLI (2.1), then it is also α -approximation sequence. Since SMQVLI (2.1) is α -well-posed in the generalized sense, it contains a subsequence converging strongly to an element of Γ . Thus, Γ is compact. Now, we prove that (4.1) holds. Suppose by contradiction that there exist $\gamma > 0$, $0 < \varepsilon_n \rightarrow 0$, and $(x_n, y_n) \in \Omega_\alpha(\varepsilon_n)$ such that

$$d((x_n, y_n), \Gamma) \geq \gamma. \quad (4.2)$$

Being $\{(x_n, y_n)\} \subset \Omega_\alpha(\varepsilon_n)$, $\{(x_n, y_n)\}$ is an α -approximating sequence for SMQVLI (2.1). Since SMQVLI (2.1) is α -well-posed in the generalized sense, there exists a subsequence $\{(x_{n_k}, y_{n_k})\}$ of $\{(x_n, y_n)\}$ converging strongly to some element of Γ . This contradicts (4.2) and so (4.1) holds.

To prove the converse, suppose that Γ is nonempty compact and (4.1) holds. Let $\{(x_n, y_n)\}$ be an α -approximating sequence for SMQVLI (2.1). Then $\{(x_n, y_n)\} \subset \Omega_\alpha(\varepsilon_n)$, and so $e(\Omega_\alpha(\varepsilon_n), \Gamma) \rightarrow 0$. This implies that there exists a sequence $\{(z_n, w_n)\} \subset \Gamma$ such that

$$\|(x_n, y_n) - (z_n, w_n)\| \rightarrow 0.$$

where the norm $\|\cdot\|$ in the product space $X \times Y$ is defined as follows

$$\|(u, v) - (\bar{u}, \bar{v})\| = \sqrt{\|u - \bar{u}\|^2 + \|v - \bar{v}\|^2}, \quad \forall (u, v), (\bar{u}, \bar{v}) \in X \times Y.$$

(It is not hard to verify that $X \times Y$ is a Banach space in terms of the last norm.)

Since Γ is compact, there exists a subsequence $\{(z_{n_k}, w_{n_k})\}$ of $\{(z_n, w_n)\}$ converging strongly to $(x_0, y_0) \in \Gamma$. Hence the corresponding subsequence $\{(x_{n_k}, y_{n_k})\}$ of $\{(x_n, y_n)\}$ converges strongly to (x_0, y_0) . Therefore, SMQVLI (2.1) is α -well-posed in the generalized sense. \square

We give the following example to illustrate that the compactness condition of Γ is necessary.

Example 4.1. Let $X = Y = \mathbf{R}$ and $C = D = \mathbf{R}^+ (= [0, +\infty))$. Let $S(x, y) = [x, x+y]$, $T(x, y) = [y, x+y]$, $A(x, y) = B(x, y) = x^2 + y^2$, $\hat{\eta}(x, z) = x - z$, $\bar{\eta}(y, w) = y - w$ and $f(x, y) = g(x, y) = xy$ for all $x, z \in C$ and $y, w \in D$. Then $\Gamma = \Omega_\alpha(\varepsilon) = [0, +\infty) \times [0, +\infty)$. It is clear that $e(\Omega_\alpha(\varepsilon), \Gamma) \rightarrow 0$ as $\varepsilon \rightarrow 0$. It is easy to see that the diverging sequence $\{(n, n)\}_{n \in \mathbf{N}}$ is an α -approximating sequence, but it has no convergent subsequence. Therefore, SMQVLI (2.1) is not α -well-posed in the generalized sense.

Corollary 4.1 (i.e., [8, Theorem 4.1]). SQEP (2.2) is α -well-posed in the generalized sense if and only if the solution set Γ of SQEP (2.2) is nonempty compact and

$$e(M_\varepsilon, \Gamma) \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.$$

Theorem 4.2. Assume that the following conditions hold:

- (i) set-valued mappings S and T are nonempty convex-valued, (s, w) -closed, (s, s) -lower semicontinuous and (s, w) -subcontinuous on $C \times D$;
- (ii) single-valued mappings A and B are (s, w^*) -continuous on $C \times D$;
- (iii) single-valued mappings $\hat{\eta}$ and $\bar{\eta}$ are Lipschitz continuous with constants $\hat{\lambda}$ and $\bar{\lambda}$ respectively, such that
 - (a) $\hat{\eta}(x_1, x_3) = \hat{\eta}(x_1, x_2) + \hat{\eta}(x_2, x_3)$ for all $x_1, x_2, x_3 \in C$ and $\bar{\eta}(y_1, y_3) = \bar{\eta}(y_1, y_2) + \bar{\eta}(y_2, y_3)$ for all $y_1, y_2, y_3 \in D$,
 - (b) $\hat{\eta}(\cdot, \cdot)$ and $\bar{\eta}(\cdot, \cdot)$ both are affine in the second variable;
 - (iv) functions f and g are continuous on $C \times D$;
 - (v) for any $y \in D$, the function $f(\cdot, y)$ is convex on C ; for any $x \in C$, the function $g(x, \cdot)$ is convex on D .

Then, SMQVLI (2.1) is α -well-posed in the generalized sense if and only if

$$\Omega_\alpha(\varepsilon) \neq \emptyset, \quad \forall \varepsilon > 0, \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0} \mu(\Omega_\alpha(\varepsilon)) = 0. \quad (4.3)$$

Proof. Suppose that SMQVLI (2.1) is α -well-posed in the generalized sense. By the same argument as in Theorem 4.1, Γ is nonempty compact, and $e(\Omega_\alpha(\varepsilon), \Gamma) \rightarrow 0$ as $\varepsilon \rightarrow 0$. Clearly $\Omega_\alpha(\varepsilon) \neq \emptyset$ for any $\varepsilon > 0$, because $\Gamma \subset \Omega_\alpha(\varepsilon)$. Observe that for any $\varepsilon > 0$, we have

$$\mathcal{H}(\Omega_\alpha(\varepsilon), \Gamma) = \max\{e(\Omega_\alpha(\varepsilon), \Gamma), e(\Gamma, \Omega_\alpha(\varepsilon))\} = e(\Omega_\alpha(\varepsilon), \Gamma).$$

Since Γ is compact, $\mu(\Gamma) = 0$ and the following relation holds (see, e.g., [12]):

$$\mu(\Omega_\alpha(\varepsilon)) \leq 2\mathcal{H}(\Omega_\alpha(\varepsilon), \Gamma) + \mu(\Gamma) = 2\mathcal{H}(\Omega_\alpha(\varepsilon), \Gamma) = 2e(\Omega_\alpha(\varepsilon), \Gamma).$$

It follows that (4.3) holds.

Conversely, suppose that (4.3) holds. It is easy to prove that $\Omega_\alpha(\varepsilon)$, for any $\varepsilon > 0$, is closed. Note that $\Omega_\alpha(\varepsilon) \subset \Omega_\alpha(\varepsilon')$ whenever $\varepsilon < \varepsilon'$, their intersection $\Omega_\alpha = \bigcap_{\varepsilon > 0} \Omega_\alpha(\varepsilon)$ is nonempty compact and satisfies: $\lim_{\varepsilon \rightarrow 0} \mathcal{H}(\Omega_\alpha(\varepsilon), \Omega_\alpha) = 0$ ([9, p. 412]), where

$$\begin{aligned} \Omega_\alpha &= \{(x_0, y_0) \in C \times D : \\ &x_0 \in S(x_0, y_0), \langle A(x_0, y_0), \hat{\eta}(x_0, z) \rangle + f(x_0, y_0) - f(z, y_0) \leq \frac{\alpha}{2} \|x_0 - z\|^2, \forall z \in S(x_0, y_0), \\ &y_0 \in T(x_0, y_0), \langle B(x_0, y_0), \bar{\eta}(y_0, w) \rangle + g(x_0, y_0) - g(x_0, w) \leq \frac{\alpha}{2} \|y_0 - w\|^2, \forall w \in T(x_0, y_0)\}. \end{aligned}$$

By Lemma 2.1, we obtain that Ω_α coincides with the solution set Γ of SMQVLI (2.1). Thus, Γ is compact.

Let $\{(x_n, y_n)\}$ be any α -approximating sequence for SMQVLI (2.1). Then there exists a sequence $\varepsilon_n > 0$ with $\varepsilon_n \rightarrow 0$ such that

$$\begin{aligned} d(x_n, S(x_n, y_n)) &\leq \varepsilon_n, \text{ and } \langle A(x_n, y_n), \hat{\eta}(x_n, z) \rangle + f(x_n, y_n) - f(z, y_n) \leq \varepsilon_n + \frac{\alpha}{2} \|x_n - z\|^2, \forall z \in S(x_n, y_n), \\ d(y_n, T(x_n, y_n)) &\leq \varepsilon_n, \text{ and } \langle B(x_n, y_n), \bar{\eta}(y_n, w) \rangle + g(x_n, y_n) - g(x_n, w) \leq \varepsilon_n + \frac{\alpha}{2} \|y_n - w\|^2, \forall w \in T(x_n, y_n). \end{aligned}$$

which means $\{(x_n, y_n)\} \subset \Omega_\alpha(\varepsilon_n)$, $\forall n \in N$. It follows from (4.3) that there exists a sequence $\{(z_n, w_n)\} \subset \Gamma$ such that

$$\|(x_n, y_n) - (z_n, w_n)\| = d((x_n, y_n), \Gamma) \leq e(\Omega_\alpha(\varepsilon_n), \Gamma) = \mathcal{H}(\Omega_\alpha(\varepsilon_n), \Gamma) \rightarrow 0.$$

Since Γ is compact, there exists a subsequence $\{(z_{n_j}, w_{n_j})\}$ of $\{(z_n, w_n)\}$ converging strongly to $(x_0, y_0) \in \Gamma$. Hence, the corresponding subsequence $\{(x_{n_j}, y_{n_j})\}$ of $\{(x_n, y_n)\}$ converges strongly to (x_0, y_0) . Thus, SMQVLI (2.1) is α -well-posed in the generalized sense. \square

Example 4.2. Let $X = Y = \mathbf{R}$ and $C = D = [0, 1]$. Let $S(x, y) = [0, x]$, $T(x, y) = [0, y]$, $A(x, y) = B(x, y) = -(x - y)^2$, $\hat{\eta}(x, z) = x - z$, $\bar{\eta}(y, w) = y - w$ and $f(x, y) = g(x, y) = -xy$ for all $x, z \in C$ and $y, w \in D$. Obviously, the conditions (i)-(v) of Theorem 4.2 are satisfied, and $\Omega_\alpha(\varepsilon) = [0, 1] \times [0, 1]$. By Theorem 4.2, SMQVLI (2.1) is α -well-posed in the generalized sense.

Corollary 4.2 (i.e., [8, Theorem 4.2]). Assume that the following conditions hold:

- (i) set-valued mappings S and T are nonempty convex-valued, (s, w) -closed, (s, s) -lower semicontinuous and (s, w) -subcontinuous on $C \times D$;
- (ii) functions f and g are continuous on $C \times D$;
- (iii) for any $y \in D$, the function $f(\cdot, y)$ is convex on C ; for any $x \in C$, the function $g(x, \cdot)$ is convex on D .

Then, SQEP (2.2) is α -well-posed in the generalized sense if and only if

$$M_\varepsilon \neq \emptyset, \forall \varepsilon > 0, \text{ and } \lim_{\varepsilon \rightarrow 0} \mu(M_\varepsilon) = 0.$$

We now give a sufficient condition for the α -well-posedness in the generalized sense of SMQVLI (2.1) in finite-dimensional spaces.

Theorem 4.3. Let X and Y be two finite-dimensional spaces. Suppose that the following conditions hold:

- (i) set-valued mappings S and T are nonempty convex-valued, closed, lower semicontinuous and subcontinuous on $C \times D$;
- (ii) single-valued mappings A and B are continuous on $C \times D$;
- (iii) single-valued mappings $\hat{\eta}$ and $\bar{\eta}$ are Lipschitz continuous with constants $\hat{\lambda}$ and $\bar{\lambda}$ respectively, such that
 - (a) $\hat{\eta}(x_1, x_3) = \hat{\eta}(x_1, x_2) + \hat{\eta}(x_2, x_3)$ for all $x_1, x_2, x_3 \in C$ and $\bar{\eta}(y_1, y_3) = \bar{\eta}(y_1, y_2) + \bar{\eta}(y_2, y_3)$ for all $y_1, y_2, y_3 \in D$,
 - (b) $\hat{\eta}(\cdot, \cdot)$ and $\bar{\eta}(\cdot, \cdot)$ both are affine in the second variable;
 - (iv) functions f and g are continuous on $C \times D$;
 - (v) for any $y \in D$, the function $f(\cdot, y)$ is convex on C ; for any $x \in C$, the function $g(x, \cdot)$ is convex on D ;
 - (vi) $\Omega_\alpha(\varepsilon)$ is nonempty bounded for some $\varepsilon > 0$.

Then, SMQVLI (2.1) is α -well-posed in the generalized sense.

Proof. Let $\{(x_n, y_n)\}$ be any α -approximating sequence for SMQVLI (2.1). Then there exists a sequence $\varepsilon_n > 0$ with $\varepsilon_n \rightarrow 0$ such that

$$d(x_n, S(x_n, y_n)) \leq \varepsilon_n, \text{ and } \langle A(x_n, y_n), \hat{\eta}(x_n, z) \rangle + f(x_n, y_n) - f(z, y_n) \leq \varepsilon_n + \frac{\alpha}{2} \|x_n - z\|^2, \forall z \in S(x_n, y_n),$$

$$d(y_n, T(x_n, y_n)) \leq \varepsilon_n, \text{ and } \langle B(x_n, y_n), \bar{\eta}(y_n, w) \rangle + g(x_n, y_n) - g(x_n, w) \leq \varepsilon_n + \frac{\alpha}{2} \|y_n - w\|^2, \forall w \in T(x_n, y_n).$$

As proven in Theorem 3.3, $\{(x_n, y_n)\}$ is bounded. Then there exists a subsequence $\{(x_{n_j}, y_{n_j})\}$ of $\{(x_n, y_n)\}$ which converges to (x_0, y_0) . Reasoning as in Theorem 3.2, one can prove that (x_0, y_0) solves SMQVLI (2.1). Therefore, SMQVLI (2.1) is α -well-posed in the generalized sense. \square

The following example shows that the nonempty boundedness of $\Omega_\alpha(\varepsilon)$ is necessary for some $\varepsilon > 0$.

Example 4.3. Let $X = Y = \mathbf{R}$ and $C = D = \mathbf{R}^+ (= [0, +\infty))$. Let $S(x, y) = [0, x]$, $T(x, y) = [0, y]$, $A(x, y) = B(x, y) = -(x - y)^2$, $\hat{\eta}(x, z) = x - z$, $\bar{\eta}(y, w) = y - w$ and $f(x, y) = g(x, y) = -xy$ for all $x, z \in C$ and $y, w \in D$. Clearly, the conditions (i)-(v) of Theorem 4.3 are satisfied. But $\Omega_\alpha(\varepsilon) = [0, +\infty) \times [0, +\infty)$ is unbounded. Therefore, SMQVLI (2.1) is not α -well-posed in the generalized sense.

Corollary 4.3 (i.e., [8, Theorem 4.3]). Let X and Y be two finite-dimensional spaces. Suppose that the following conditions hold:

- (i) set-valued mappings S and T are nonempty convex-valued, closed, lower semicontinuous and subcontinuous on $C \times D$;
 - (ii) functions f and g are continuous on $C \times D$;
 - (iii) for any $y \in D$, the function $f(\cdot, y)$ is convex on C ; for any $x \in C$, the function $g(x, \cdot)$ is convex on D ;
 - (iv) M_ε is nonempty bounded for some $\varepsilon > 0$.
- Then, SQEP (2.2) is α -well-posed in the generalized sense.

5. Concluding Remarks

In this paper, we generalize the concept of α -well-posedness to the system of mixed quasivariational-like inequalities which includes as a special case symmetric quasi-equilibrium problems considered in Long and Huang [8]. It is well known that symmetric quasi-equilibrium problems include equilibrium problems, Nash equilibrium problems, quasivariational inequalities, variational inequalities and fixed point problems as special cases. It is worth emphasizing that the results presented in [8] generalize and improve some known results in the recent literature; see, e.g., [13-18].

Further, under some suitable conditions, we obtain some metric characterizations of α -well-posedness for the system of mixed quasivariational-like inequalities in Banach spaces. The results presented in this paper represent the supplement, improvement, generalization and development of Long and Huang's known results [8] in the following aspects:

(i) The symmetric quasi-equilibrium problem (SQEP) in [8] is extended to develop the more general problem, that is, the system of mixed quasivariational-like inequalities (SMQVLI) in the setting of Banach spaces. Moreover, the concept of α -well-posedness (resp., in the generalized sense) for SQEP is extended to develop the concept of α -well-posedness (resp., in the generalized sense) for SMQVLI.

(ii) Since the system of mixed quasivariational-like inequalities (SMQVLI) is more general and more complicated than the symmetric quasi-equilibrium problem (SQEP), the assumptions in our results are very different from the corresponding ones in [8]; see, for instance, the assumptions imposed on the single-valued mappings A , B and $\hat{\eta}$, $\bar{\eta}$.

(iii) The new technique of arguments are applied to deriving our main results. As a matter of fact, in the process of proving our main results, our arguments depend on the properties of the mappings $\hat{\eta}$, $\bar{\eta}$, the uniform boundedness principle for a family of linear continuous functionals, the completion of the Banach space $(X \times Y, \|\cdot\|)$, etc. For instance, see the proof of Theorem 3.2.

References

- [1] A.N. Tykhonov, On the stability of the functional optimization problem, USSR J. Comput. Math. Math. Phys. 6 (1966) 631-634.

- [2] E. Blum, W. Oettli, From optimization and variational inequalities to equilibrium problems, *Math. Stud.* 63 (1994) 123-145.
- [3] F. Facchinei, J.S. Pang, *Finite-Dimensional Variational Inequalities and Complementarity Problems*, Springer, New York, 2003.
- [4] C. Baiocchi, A. Capelo, *Variational and Quasivariational Inequalities: Applications to Free Boundary Problems*. Wiley, New York, 1984.
- [5] D. Kinderlehrer, G. Stampacchia, *An Introduction to Variational Inequalities and Their Applications*, Academic Press, New York, 1980.
- [6] E.L. Fuster, A. Petrusel, J.C. Yao, Iterated functions systems and well-posedness, *Chaos Solitons Fractals*. (2008). doi:10.1016/j.chaos.2008.06.019
- [7] A. Petrusel, I.A. Rus, J.C. Yao, Well-posedness in the generalized sense of the fixed point problems for multivalued operators, *Taiwanese J. Math.* 11 (2007) 903-914.
- [8] X.J. Long, N.J. Huang, Metric characterizations of α -well-posedness for symmetric quasi-equilibrium problems, *J. Global Optim.* 45 (2009) 459-471.
- [9] K. Kuratowski, *Topology*, Vols. 1 and 2, Academic Press, New York, 1968.
- [10] Q.H. Ansari, J.C. Yao, Iterative schemes for solving mixed variational-like inequalities, *J. Optim. Theory Appl.* 108 (3) (2001) 527-541.
- [11] S.K. Berberian, *Lectures in Functional Analysis and Operator Theory*, Springer-Verlag, Berlin, Germany, 1974.
- [12] A.L. Dontchev, T. Zolezzi, *Well-Posedness Optimization Problems*, *Lecture Notes in Math.* Vol. 1543, Springer, Berlin, 1993.
- [13] L.C. Ceng, J.C. Yao, Well-posedness of generalized mixed variational inequalities, inclusion problems and fixed point problems, *Nonlinear Anal. TMA* 69 (2008) 4585-4603.
- [14] L.C. Ceng, N. Hadjisavvas, S. Schaible, J.C. Yao, Well-posedness for mixed quasivariational-like inequalities, *J. Optim. Theory Appl.* 139 (2008) 109-125.
- [15] Y.P. Fang, R. Hu, N.J. Huang, Well-posedness for equilibrium problems and optimization problems with equilibrium constraints, *Comput. Math. Appl.* 55 (2008) 89-100.
- [16] Y.P. Fang, N.J. Huang, J.C. Yao, Well-posedness of mixed variational inequalities, inclusion problems and fixed point problems, *J. Global Optim.* 41 (2008) 117-133.
- [17] M.B. Lignola, Well-posedness and L -well-posedness for quasivariational inequalities, *J. Optim. Theory Appl.* 128 (2006) 119-138.
- [18] M.B. Lignola, J. Morgan, α -Well-posedness for Nash equilibria and for optimization problems with Nash equilibrium constraints, *J. Glob. Optim.* 36 (2006) 439-459.
- [19] M.B. Lignola, J. Morgan, Approximate solutions and α -well-posedness for variational inequalities and Nash equilibria, In: G. Zaccour (ed.), *Decision and Control in Management Science*, pp. 367-378. Kluwer Academic Publisher, Dordrecht, 2001.
- [20] E. Bednarczuck, J.P. Penot, Metrically well-set minimization problems, *Appl. Math. Optim.* 26 (1992) 273-285.
- [21] G.P. Crespi, A. Guerraggio, M. Rocca, Well-posedness in vector optimization problems and vector variational inequalities, *J. Optim. Theory Appl.* 132 (2007) 213-226.

- [22]M. Furi, A. Vignoli, About well-posed minimization problems for functionals in metric spaces, *J. Optim. Theory Appl.* 5 (1970) 225-229.
- [23]X.X. Huang, Pointwise well-posedness of perturbed vector optimization problems in a vector-valued variational principle, *J. Optim. Theory Appl.* 108 (2000) 671-686.
- [24]X.X. Huang, X.Q. Yang, Generalized Levitin-Polyak well-posedness in constrained optimization, *SIAM J. Optim.* 17 (2006) 243-258.
- [25]X.X. Huang, X.Q. Yang, Levitin-Polyak well-posedness of constrained vector optimization problems, *J. Glob. Optim.* 37 (2007) 287-304.
- [26]X.X. Huang, X.Q. Yang, Levitin-Polyak well-posedness of generalized variational inequalities with functional constraints, *J. Ind. Manag. Optim.* 3 (2007) 571-584.
- [27]X.X. Huang, X.Q. Yang, D.L. Zhu, Levitin-Polyak well-posedness of variational inequality problems with functional constraints, *J. Glob. Optim.* 44 (2009) 159-174.
- [28]X.J. Long, N.J. Huang, K.L. Teo, Levitin-Polyak well-posedness for equilibrium problems with functional constraints, *J. Inequal. Appl.* (2008), Article ID 657329, 14 pages.
- [29]G. Mastroeni, Gap function for equilibrium problems, *J. Glob. Optim.* 27 (2003) 411-426.
- [30]J. Li, N.J. Huang, J.K. Kim, On implicit vector equilibrium problems, *J. Math. Anal. Appl.* 283 (2003) 501-512.
- [31]I. Del Prete, M.B. Lignola, J. Morgan, New concepts of well-posedness for optimization Problems with variational inequality constraints, *J. Inequal. Pure Appl. Math.* 4 (1) (2003) (Article 5).
- [32]M.B. Lignola, J. Morgan, Well-posedness for optimization problems with constraints defined by variational inequalities having a unique solution, *J. Glob. Optim.* 16 (2000) 57-67.
- [33]R. Lucchetti, F. Patrone, A characterization of Tykhonov well-posedness for minimum problems, with applications to variational inequalities, *Numer. Funct. Anal. Optim.* 3 (1981) 461-476.
- [34]B. Lemaire, C. Ould Ahmed Salem, J.P. Revalski, Well-posedness by perturbations of variational problems, *J. Optim. Theory Appl.* 115 (2002) 345-368.
- [35]J. Morgan, Approximations and well-posedness in multicriteria games, *Ann. Oper. Res.* 137 (2005) 257-268.
- [36]J. Yu, H. Yang, C. Yu, Well-posed Ky Fan's point, quasi-variational inequality and Nash equilibrium problems, *Nonlinear Anal. TMA* 66 (2007) 777-790.
- [37]T. Zolezzi, Well-posedness criteria in optimization with application to the calculus of variations, *Nonlinear Anal. TMA* 25 (1995) 437-453.
- [38]M.A. Noor, W. Oettli, On general nonlinear complementarity problems and quasiequilibria, *Le Math.* XLIX (1994) 313-331.
- [39]Y. Yao, R. Chen, Y.-C. Liou, A unified implicit algorithm for solving the triple-hierarchical constrained optimization problem, *Math. & Comp. Model.*, in press.
- [40]Y.C. Lin, Finite-step Relaxed Hybrid Steepest-descent Methods for Variational Inequalities, *J. Inequal. Appl.* (2008), Article ID 598632, 13 pages.

[41]Y. Yao and N. Shahzad, New methods with perturbations for non-expansive mappings in Hilbert spaces, Fixed Point Theory and Appl., in press.