

Urological Research

Protective effect of Flos carthami extract against ethylene glycol-induced urolithiasis in rats

--Manuscript Draft--

Manuscript Number:	URES-D-11-00224R1
Full Title:	Protective effect of Flos carthami extract against ethylene glycol-induced urolithiasis in rats
Article Type:	Original Paper
Keywords:	calcium oxalate; Ethylene glycol; Traditional Chinese medicine; Flos carthami
Corresponding Author:	Wen-Chi Chen, M.D., Ph.D. Graduate Institute of Integrated Medicine Taichung, TAIWAN, REPUBLIC OF CHINA
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Graduate Institute of Integrated Medicine
Corresponding Author's Secondary Institution:	
First Author:	Wu-Chou Lin, M.D.
First Author Secondary Information:	
Order of Authors:	Wu-Chou Lin, M.D. Ming-Tsung Lai, M.D., Ph.D. Huey-Yi Chen, M.D., Ph.D. Chien-Yi Ho, M.D. Kee-Ming Man, M.D. Jui-Lung Shen, M.D., Ph.D. Yuan-Ju Lee, M.D. Fuu-Jen Tsai, M.D., Ph.D. Yung-Hsiang Chen, Ph.D. Wen-Chi Chen, M.D., Ph.D.
Order of Authors Secondary Information:	
Abstract:	<p>Flos carthami (FC), also known as <i>Carthamus tinctorius</i>, is a traditional Chinese herbal plant that has been prescribed since centuries for treating various symptoms related to blood circulation improvement. This study aimed to investigate the effects of FC on calcium oxalate (CaOx) formation in ethylene glycol (EG)-fed rats. A total of 50 male Sprague-Dawley rats were divided into the following 6 groups: group 1, as the normal control (n = 5); group 2 received gastric gavages of starch and 0.75% EG (placebo, n = 5) as a stone inducer; group 3 (n = 10) received EG and potassium citrate as positive controls; group 4 (n = 10) received 0.75% EG and 300 mg/day FC; group 5 (n = 10) was treated with EG and 600 mg/day FC; and group 6 (n = 10) received with EG and 1200 mg/day FC. For all experimental animals, 24-h urine and blood samples were analyzed at the beginning and end of the experiment. Kidney tissue was histopathologically examined using a polarized light microscope, and crystal deposits were evaluated by a semi-quantitative scoring method; these scores were significantly lower in the FC groups (600 and 1200 mg/day) than in the placebo group. Thus, FC administration appeared to inhibit the deposition of CaOx crystal EG-fed rats. We therefore consider that FC may be effective for preventing stone disease, albeit with certain side effects, such as a bleeding tendency. Further clinical trials are needed for evaluating its benefits and possible side effects.</p>

Response to Reviewers:

Please see attachment

Urological Research

Protective effect of *Flos carthami* extract against ethylene glycol-induced urolithiasis in rats

Wu-Chou Lin • Ming-Tsung Lai • Huey-Yi Chen • Chien-Yi Ho • Kee-Ming Man •
Jui-Lung Shen • Yuan-Ju Lee • Fuu-Jen Tsai • Yung-Hsiang Chen • Wen-Chi Chen

W.-C. Lin • M.-T. Lai • H.-Y. Chen • C.-Y. Ho • F.-J. Tsai • Y.-H. Chen (✉) • W.-C. Chen (✉)
Graduate Institute of Integrated Medicine, Institute of Chinese Medicine, School of Chinese Medicine,
College of Chinese Medicine, China Medical University, No. 91, Hsueh-Shih Road, Taichung 40402,
Taiwan
e-mail: yhchen@mail.cmu.edu.tw

W.-C. Chen
e-mail: wgchen@mail.cmu.edu.tw

W.-C. Lin • M.-T. Lai • H.-Y. Chen • C.-Y. Ho • F.-J. Tsai • Y.-H. Chen • W.-C. Chen
Departments of Urology, Family Medicine, Obstetrics and Gynecology, Medical Genetics and
Pediatrics, and Medical Research, China Medical University Hospital, Taichung, Taiwan

M.-T. Lai
Department of Pathology, Show Chwan Memorial Hospital, Changhua, Taiwan

K.-M. Man
Department of Anesthesiology, Tungs' Taichung MetroHarbor Hospital, Taichung, Taiwan

J.-L. Shen
Department of Dermatology, Taichung Veterans General Hospital, Taichung, Taiwan

Y.-J. Lee
Department of Urology, National Taiwan University Hospital, Taipei Taiwan

Urological Research

(Ms. No. URES-D-11-00224.R1)

Protective effect of *Flos carthami* extract against ethylene glycol-induced urolithiasis in rats

Wu-Chou Lin,^{1,2} Ming-Tsung Lai,^{1,3} Huey-Yi Chen,^{1,2} Chien-Yi Ho,^{1,2} Kee-Ming Man,^{2,4}

Jui-Lung Shen,⁵ Yuan-Ju Lee,⁶ Fuu-Jen Tsai,^{1,2} Yung-Hsiang Chen,^{1,2,*} and Wen-Chi Chen^{1,2,*}

¹ Departments of Urology, Family Medicine, Obstetrics and Gynecology, Medical Genetics and Pediatrics, and Medical Research, China Medical University Hospital, Taichung, Taiwan

² Graduate Institute of Integrated Medicine, Institute of Chinese Medicine, School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan

³ Department of Pathology, Show Chwan Memorial Hospital, Changhua, Taiwan

⁴ Department of Anesthesiology, Tungs' Taichung MetroHarbor Hospital, Taichung, Taiwan

⁵ Department of Dermatology, Taichung Veterans General Hospital, Taichung, Taiwan

⁶ Department of Urology, National Taiwan University Hospital, Taipei Taiwan

* Corresponding authors. Address: No. 91, Hsueh-Shih Road, Taichung, 40402, Taiwan.

Tel: +886-4-22053366-3500; fax: +886-4-22037690.

E-mail address: wgchen@mail.cmu.edu.tw; yhchen@mail.cmu.edu.tw

ABSTRACT

Flos carthami (FC), also known as *Carthamus tinctorius*, is a traditional Chinese herbal plant that has been prescribed since centuries for treating various symptoms related to blood circulation **improvement**. This study aimed to investigate the effects of FC on calcium oxalate (CaOx) formation in ethylene glycol (EG)-fed rats. A total of 50 male Sprague-Dawley rats were divided into the following 6 groups: group 1, as the normal control (n = 5); group 2 received gastric gavages of starch and 0.75% EG (placebo, n = 5) as a stone inducer; group 3 (n = 10) received EG and potassium citrate as positive controls; group 4 (n = 10) received 0.75% EG and 300 mg/day FC; group 5 (n = 10) was treated with EG and 600 mg/day FC; and group 6 (n = 10) received with EG and 1200 mg/day FC. For all experimental animals, 24-h urine and blood samples were analyzed at the beginning and end of the experiment. Kidney tissue was histopathologically examined using a polarized light microscope, and crystal deposits were evaluated by a semi-quantitative scoring method; these scores were significantly lower in the FC groups (600 and 1200 mg/day) than in the placebo group. Thus, FC administration appeared to inhibit the deposition of CaOx crystal EG-fed rats. We therefore consider that FC may be effective for preventing stone disease, albeit with certain side effects, such as a bleeding tendency. Further clinical trials are needed for evaluating its benefits and possible side effects.

Keywords: Calcium oxalate, Ethylene glycol, Traditional Chinese medicine, *Flos carthami*

Introduction

Nephrolithiasis has a high prevalence worldwide, including Taiwan [1-3]. Changing lifestyles, genetic factors, and climate-related changes are considered to gradually increase the prevalence and decrease the age of onset [4, 5]. However, long-term treatment strategies for preventing the recurrence of urinary stones are few; these include the administration of potassium citrate (K-citrate) [6]. Currently, several herbal drugs have been used in the treatment of nephrolithiasis. Reported herbal treatments for urolithiasis include single-plant as well as polyherbal formulae such as Wulingsan, Zhulingtang [7, 8], Takusha [9], *Herniaria hirsuta* extract [10], *Phyllanthus niruri* [11], and other kampo medicines [12-14]. Most of these studies have demonstrated several degrees of inhibition of calcium oxalate (CaOx) crystal formation.

We previously performed a quick screening for antilithic Chinese herbal drugs in a *Drosophila* animal model [15, 16]. Eighty candidate herbal extracts with possible effects on the kidneys were chosen from a database. *Flos carthami* (FC, also known as *Carthami tinctorius*, plant family name: Asteraceae) has been found to have the highest efficacy for the inhibition of CaOx crystal deposition in *Drosophila* Malpighian tubules (unpublished data). Therefore, we chose FC for further studying its effects on the inhibition of CaOx crystallization. FC contains carthamin, neocarthamine, and kaempferol 3-rhamnoglucoside; in China, it has been used in the treatment of urological diseases since centuries [17]. Based on

1 an official website of our government and literatures, FC was initially used to improve blood
2
3
4 circulation based on its anti-coagulation effects [18, 19]. FC is also used in amelioration of
5
6
7 pain, dilation of coronary arteries, lowering blood pressure, and anti-inflammatory effects.
8
9
10 Moderate and low doses of FC can improve blood flow [19]. However, a high dose of FC
11
12
13 enhances the risk of bleeding tendencies [20]. Li et al. reported that FC decreased blood
14
15
16 viscosity, plasma viscosity, and erythrocyte aggregation in a rat model with blood stasis [21].
17
18
19 However, few animal and clinical studies have been conducted to confirm the effects of FC on
20
21
22 kidney stone disease. The purpose of this study was to investigate the effects of FC on CaOx
23
24
25 formation in ethylene glycol (EG)-fed rats. Therefore, we focused on the *in vivo* effects of FC
26
27
28 in male SD rats by analyzing changes in body weight and the biochemical parameters of urine
29
30
31 and serum along with assessment of CaOx deposition in rat kidneys using a polarized light
32
33
34 microscope and a computer-assisted image scoring system.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Materials and methods

Experimental animals and designs

Fifty male Sprague-Dawley (SD) rats (six weeks of age), each weighing approximately 250 - 300 g, were purchased from BioLASCO Co (Taipei, Taiwan) and acclimated to a room temperature of 25°C with a 12 h light, 12 h dark cycle. They were housed in the animal breeding center of the China Medical University (Taichung, Taiwan) and fed standard commercial rat chow. The rats were randomly divided into six groups. Rats in the experimental groups (groups 2, 3, 4, 5, and 6) were given free access to drinking water containing 0.75% EG as a stone inducer throughout the entire 4-week experimental period

(According to the previous studies [7, 8], the CaOx crystals appeared after EG administration for 4 weeks. The incidence of crystal formation in EG group for 4 weeks was about 90%).

The commercial herbal powder of FC was furnished by the Koda pharmaceutical company (Taoyun, Taiwan). The quality control of FC was provided by the Koda pharmaceutical company (**Figure 1**). The HPLC fingerprint of FC powder was examined by the Medical and

Pharmaceutical Industry Technology and Development Center (Taiwan) (data not shown).

Briefly, reverse-phase C18 column (Cosmosil 5C18-MS-II Waters 4.6 × 250 mm, 5 μm, NACALAI), and a linear gradient of 0.1% H₃PO₄ and acetonitrile at a flow rate of 1 ml/min were used. Chromatographed samples were monitored at UV 203 nm. The metal contents of FC powder were analyzed; the result revealed cadmium ≤ 5.0 ppm, mercury ≤ 0.1 ppm,

1 arsenic \leq 2.0 ppm and lead \leq 10.0 ppm.
2
3

4 Rats in group 1 (n = 5) served as the normal control and were given de-ionized distilled
5
6 water and fed normal chow. Rats in group 2 (n = 5) were treated with gastric gavages of 60
7
8 mg starch and 0.75% EG (placebo) as a stone inducer once daily. Rats in group 3 (n = 10)
9
10 were given EG-added drinking water and daily gastric gavages of K-citrate (500 mg/day). The
11
12 doses of FC were determined by our pilot study and translated from the dosage used for
13
14 human. The dose of scientific herbal medicine for human is 3 – 10 g/day. After conversing of
15
16 animal doses to human equivalent dose based on body surface area [22], 300 mg/day were
17
18 determined for the study. We further used 600 and 1200 mg/day to examine the high-dose
19
20 response. Rats in group 4 (n = 10) were given EG-added drinking water and treated with daily
21
22 300 mg FC of gastric gavages. Rats in group 5 (n = 10) were given EG-added drinking water
23
24 and treated with daily 600 mg FC of gastric gavages. Rats in group 6 (n = 10) were given
25
26 EG-added drinking water and treated with daily 1,200 mg FC of gastric gavages. Various
27
28 treatments were administered to the rats in the same volume (1 - 2 ml). Body weights of rats
29
30 were recorded before and after the experiment. All rats were sacrificed at the end of the
31
32 experiment under ether anesthesia. Both kidneys from each rat were harvested and weighed.
33
34 Right-sided kidneys were fixed in formalin, embedded in paraffin and stained with
35
36 hematoxylin and eosin Y solution. We then examined the histological sections of kidneys for
37
38 crystal deposits with a polarized light microscope.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Analysis of serum and urine

Twenty-four hour urine samples were collected twice, once at the beginning and once at the end of the experiment. For collection of 24 h urine samples, we placed rats individually in metabolic cages. The food and water were given to the rats during 24 h urine collection. Serum was obtained for biochemical analysis at the beginning and end of the experiment. We drew blood samples from the tails at the beginning of the experiment, and from the heart at the end. Urine and serum calcium, free calcium, phosphorus, and pH values, and known risk factor parameters such as sodium, potassium, magnesium and citrate for stone disease, were measured with a Hitachi-7150 and a Roche-Omnicon analyzer. Prothrombin time (PT) and activated partial thromboplastin time (APTT) were measured. We checked serum aspartate transaminase (AST), alanine transaminase (ALT), uric acid, and urinary micro-albumin at the ending of the experiment.

Evaluation of the severity of renal crystal deposition

We used a polarized light microscope to measure the severity of crystal deposition as previously described. In brief, spotlight the bi-refringent crystals of CaOx with a magnification factor of 100 to evaluate the score. A subjective, semi-quantitative scoring system was used with four grades ranging from 0 to 3+ (where 0 = none, 1+ = few, 2+ = several, and 3+ = many crystal deposits) to evaluate the degree of crystal deposition as previously described [10-12]. The scoring system was according to our computer-assisted image-scoring system

1 *ImageScoring* [7, 8]. This software offers randomized and blinding selection of digital images
2 spontaneously. We then randomly took a digital image of one of the eight regions under a
3 polarized light microscope (100×) using a digital camera mounted onto the microscope. Six
4 independent coworkers used this computer-assisted system to score every individual picture in
5 the database. Finally, all of the personal score files were collected, merged, and sent for
6 statistical analysis.
7
8
9
10
11
12
13
14
15
16
17

18 Statistical methods

19
20
21 Data regarding rat weights and parameters of urine and serum (presented as mean ± S.D.)
22 were analyzed by one-way ANOVA. The severity of renal crystal deposition was expressed as
23 an ordinal scale using the semi-quantitative scoring method described above. The crystal
24 deposit scores were analyzed by the non-parametric Kruskal-Wallis test for inter-group
25 comparison, and by the Wilcoxon rank sum test (Mann-Whitney U) for pair wise comparison.
26
27
28
29
30
31
32
33
34
35
36

37 A *P* value of < 0.05 denotes the presence of statistical significance. The *Statistical Package*
38 *for the Social Science* (SPSS® for Windows, release 11.5) was used for statistical analysis.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Results

There were no differences in the baseline biochemical data among the 6 groups. No significant differences in body weight, urine volume, pH, micro-albumin, creatinine, uric acid, Na^+ , K^+ , Cl^+ , Mg^{2+} , and Ca^{2+} ions were observed (**Table 1**). At the end of the 4-week experiment, no significant differences were observed for body weight among the 6 groups.

At the end of the experimental period, the urine volume in group 2 was significantly increased. By contrast, the urine volume in group 6 was significantly decreased after 1200 mg/day FC administration. Serum creatinine was significantly decreased in the group 3. Serum ALT and AST levels were significantly increased in EG-treated group, whereas FC administration reduced the ALT and AST levels, suggesting a potential hepatoprotective effect of FC. All other urine and serum biochemical data were similar among the 6 groups.

Histological studies revealed that no calcifications in the kidneys of the normal control rats. As expected, several crystal deposits were found in the renal cortex and medulla of the EG-fed rats. Under polarized light microscopy, the crystals exhibited a birefringent appearance as shown in **Figure 2**, wherein multiple crystals were found both in the renal cortex and the medulla. Analyses of the crystal deposit scores revealed that the scores of the K-citrate- and FC-fed groups (600 mg/day and 1200 mg/day) were significantly lower than those of the placebo group (**Figure 3**). Overall, FC administration significantly inhibited CaOx crystal deposition in the kidneys of the experimental rats.

Discussion

The results of the present study were consistent with our hypothesis that FC would inhibit CaOx crystal formation since the severity of renal crystal deposition and the proportion of crystal formation in the 600- and 1200-mg/day FC treatment groups were significantly lower than those in the placebo group. To the best of our knowledge, this is the first evidence for the inhibitory effect of FC on CaOx crystal formation in EG-fed rats. Based on these preliminary results, we aim to perform further investigations of the prophylactic effects of FC on nephrolithiasis in a larger sample size and different dosages. At least, we can no longer assume that the dose of FC equal to the human daily dose per unit of body weight reveal obvious nephrolithic prophylaxis.

Administering EG to rats is the most common and effective method for inducing CaOx renal stones *in vivo*, and our present study successfully induced crystal formation using this model [23, 24]. Other methods for experimentally inducing CaOx renal stones include the addition of hydroxy-L-proline (HLP) to drinking water or food, implantation of osmotic mini-pumps filled with oxalate, and intraperitoneal administration of oxalate or HLP [25, 26]. However, it has been shown that if rats are made significantly hyperoxaluric via intraperitoneal injections of sodium oxalate or oral intake of EG, CaOx crystalluria occurs rather than the formation of papillary plaques or stones [15, 26, 27]. EG has also been criticized because it induces metabolic acidosis, and some of its metabolites are nephrotoxic

1 in nature [28]. Furthermore, both oxalate and CaOx crystals are known to damage renal
2
3
4 epithelial cells. Therefore, it is difficult to distinguish the effects of EG and its metabolites
5
6
7 from those induced by oxalate and CaOx crystals. However, Green et al. reported that
8
9
10 metabolic acidosis does not always occur if renal function is preserved [29]. Khan et al.
11
12
13 reported that HLP, a common ingredient of many diets, is a hyperoxaluria-inducing agent.
14
15
16 They further reported that CaOx crystal deposition in animals administered HLP was similar
17
18
19 to that observed in other animal hyperoxaluria models of CaOx, although HLP was less
20
21
22 nephrotoxic [26]. Therefore, we aim to compare the differences of the effects of FC between
23
24
25 HLP and EG-induced nephrolithiasis models in our next investigation.
26
27
28

29 Many medicinal plants are used to treat urolithiasis worldwide [30-32], of which *Alisma*
30
31
32 *orientalis* and *Poria cocos* Wolf have been shown to inhibit the kidney stone formation
33
34
35 process [9, 33-35]. *Alisma orientalis* (also known as Takusha) strongly suppresses each step
36
37
38 of crystal formation, growth, and aggregation of CaOx crystals *in vitro*; furthermore, the renal
39
40
41 calcium content has been shown to decrease in *Alisma*-treated rats [33, 36]. In addition,
42
43
44 several studies have shown that this herb significantly decreased the formation of CaOx
45
46
47 deposits [35, 37, 38]. All of these reports suggest that herbal medicines may be a useful
48
49
50 strategy for preventing renal stones. However, it is important to note that FC seems to prolong
51
52
53 the PTT and APTT. Since FC was initially used to improve blood circulation in the traditional
54
55
56 Chinese medicine, it should be noticed when used in patients with abnormal function of
57
58
59
60
61
62
63
64
65

1 coagulation, although there were no apparent bleeding complications in the animals treated
2
3
4 with the FC extract.
5

6
7 The present study has some limitations. The antilithic mechanism of FC remains unclear.
8
9 Some key urine numbers are not provided, including urine oxalate and citrate. Without this
10
11 sort of basic information, we have little idea what the extract may or may not be doing. In
12
13 addition, if the extract change EG metabolism to oxalate or oxalate handling, does it inhibit
14
15 crystallization, and its cellular responses are unclear. A complete characterization of the model
16
17 and effects of the extract on urinary components would interpret these results; appropriate
18
19 evaluation must be further established.
20
21

22
23
24 The software (*Image Scoring*) that we created has proved to be a helpful tool because it
25
26 could easily reduce the time-consuming process of reviewing a multitude of images.
27
28 Furthermore, the automatically randomized selection process diminished examination bias
29
30 and offered a reliable method for blindness toward the samples. Moreover, the format of the
31
32 scoring files was compatible with the statistical package used; thus, no data transfer was
33
34 needed, preventing the possibility of transfer errors [7, 8].
35
36
37
38
39
40
41

42
43 In summary, our results revealed that FC administration led to a significant reduction in
44
45 CaOx crystal formation as compared to that in the placebo group. The herbal medicine FC
46
47 appeared to effectively prevent kidney stone formation, indicating that this herb might be an
48
49 effective antilithic drug. The exact mechanism underlying its effects remains unknown, and
50
51 further experimental and clinical studies are required to elucidate the chemical constituents of
52
53 the extracts and the mechanism(s) responsible for its pharmacological activities.
54
55
56
57
58
59
60
61

1
2
3
4 **Acknowledgments**
5
6

7 This study was supported in part by grants from the Department of Health
8
9
10 (DOH101-TD-B-111-004), the National Science Council of Taiwan (NSC
11
12
13 98-2314-B-039-023-MY3 and NSC 100-2320-B-039-008-MY2), and China Medical
14
15
16 University Hospital (DMR-100-068). We thank Dr. Chou-Huang Tsai for his helping in the
17
18
19 process of computer aid scoring system (*ImageScoring*) and we thank Chiao-Hui Chang,
20
21
22 Yu-Ning Lin, Yun-Chen Hsu, and Chi-Hsiang Wei for experimental assistance.
23
24
25
26
27
28

29 **Conflict of interest**
30
31

32 The authors declare that there are no conflicts of interest.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 **Figure Legends**

2
3
4 **Figure 1.** Thin layer chromatography of *Flos carthami* (Chinese name: Hong Hua) compared
5
6
7 with standard.
8
9

10
11
12
13 **Figure 2.** Represented paraffin sections viewed under polarized light of rat kidneys in each
14
15
16 groups. Arrows indicate the crystals in the cortex, tubular, and medulla ($\times 100$).
17
18
19

20
21
22
23 **Figure 3.** Quantitative analysis of crystal deposits scores. Values represent mean \pm SD.
24

25
26 *Values are significantly different ($p < 0.05$) from the control group, #Values are significantly
27
28
29 different ($P < 0.05$) from the EG + starch group.
30
31

References

1. Lee YH, Huang WC, Tsai JY, Lu CM, Chen WC, Lee MH, Hsu HS, Huang JK, Chang LS (2002) Epidemiological studies on the prevalence of upper urinary calculi in Taiwan. *Urol Int* 68:172-177
2. Bansal AD, Hui J, Goldfarb DS (2009) Asymptomatic nephrolithiasis detected by ultrasound. *Clin J Am Soc Nephrol* 4:680-684
3. Sowers MR, Jannausch M, Wood C, Pope SK, Lachance LL, Peterson B (1998) Prevalence of renal stones in a population-based study with dietary calcium, oxalate, and medication exposures. *Am J Epidemiol* 147:914-920
4. Goldfarb DS (2003) Increasing prevalence of kidney stones in the United States. *Kidney Int* 63:1951-1952
5. Brikowski TH, Lotan Y, Pearle MS (2008) Climate-related increase in the prevalence of urolithiasis in the United States. *Proc Natl Acad Sci U S A* 105:9841-9846
6. Lee YH, Huang WC, Tsai JY, Huang JK (1999) The efficacy of potassium citrate based medical prophylaxis for preventing upper urinary tract calculi: a midterm followup study. *J Urol* 161:1453-1457
7. Tsai CH, Chen YC, Chen LD, Pan TC, Ho CY, Lai MT, Tsai FJ, Chen WC (2008) A traditional Chinese herbal antilithic formula, *Wulingsan*, effectively prevents the renal deposition of calcium oxalate crystal in ethylene glycol-fed rats. *Urol Res* 36:17-24
8. Tsai CH, Pan TC, Lai MT, Lee SC, Chen ML, Jheng JR, Chen WC (2009) Prophylaxis of experimentally induced calcium oxalate nephrolithiasis in rats by *Zhulingtang*, a traditional Chinese herbal formula. *Urol Int* 82:464-471
9. Yasui T, Fujita K, Sato M, Sugimoto M, Iguchi M, Nomura S, Kohri K (1999) The effect of takusha, a kampo medicine, on renal stone formation and osteopontin expression in a rat urolithiasis model. *Urol Res* 27:194-199
10. Atmani F, Khan SR (2000) Effects of an extract from *Herniaria hirsuta* on calcium oxalate crystallization in vitro. *BJU Int* 85:621-625
11. Freitas AM, Schor N, Boim MA (2002) The effect of *Phyllanthus niruri* on urinary inhibitors of calcium oxalate crystallization and other factors associated with renal stone formation. *BJU Int* 89:829-834
12. Liu QL, Sato S, Kishikawa T, Matsuzaki H, Yamanaka N (2001) Effectiveness of a traditional Chinese medicine, *Wulingsan*, in suppressing the development of nephrocalcinosis induced by a high phosphorus diet in young rats. *Med Electron Microsc* 34:103-114
13. Koide T, Yamaguchi S, Utsunomiya M, Yoshioka T, Sugiyama K (1995) The inhibitory effect of kampo extracts on *in vitro* calcium oxalate crystallization and *in vivo* stone formation in an animal model. *Int J Urol* 2:81-86
14. Al-Ghamdi SS, Al-Ghamdi AA, Shammah AA (2007) Inhibition of calcium oxalate

- nephrotoxicity with *Cymbopogon schoenanthus* (Al-Ethkher). Drug Metab Lett 1:241-244
15. Chen YH, Liu HP, Chen HY, Tsai FJ, Chang CH, Lee YJ, Lin WY, Chen WC (2011) Ethylene glycol induces calcium oxalate crystal deposition in Malpighian tubules: a *Drosophila* model for nephrolithiasis/urolithiasis. Kidney Int 80:369-377
 16. Knauf F, Preisig PA (2011) *Drosophila*: a fruitful model for calcium oxalate nephrolithiasis? Kidney Int 80:327-329
 17. Li Y, Che Q (1998) Studies on chemical components of *Carthamus tinctorius* petals. Yao Xue Xue Bao 33:626-628
 18. Sun M, Zhang JJ, Shan JZ, Zhang H, Jin CY, Xu S, Wang YL (2009) Clinical observation of Danhong Injection (herbal TCM product from *Radix Salviae miltiorrhizae* and *Flos Carthami tinctorii*) in the treatment of traumatic intracranial hematoma. Phytomedicine 16:683-689
 19. Lu Y, Hu YL, Kong XF, Wang DY (2008) Selection of component drug in activating blood flow and removing blood stasis of Chinese herbal medicinal formula for dairy cow mastitis by hemorheological method. J Ethnopharmacol 116:313-317
 20. Zhang S, Xu C (1997) Progress in the clinical uses of *flos Carthami*. Zhongguo Zhong Yao Za Zhi 22:439-440, 442
 21. Li HX, Han SY, Wang XW, Ma X, Zhang K, Wang L, Ma ZZ, Tu PF (2009) Effect of the carthamins yellow from *Carthamus tinctorius* L. on hemorheological disorders of blood stasis in rats. Food Chem Toxicol 47:1797-1802
 22. Reagan-Shaw S, Nihal M, Ahmad N (2008) Dose translation from animal to human studies revisited. The FASEB journal : official publication of the Federation of American Societies for Experimental Biology 22:659-661
 23. Lee YH, Huang WC, Chiang H, Chen MT, Huang JK, Chang LS (1992) Determinant role of testosterone in the pathogenesis of urolithiasis in rats. J Urol 147:1134-1138
 24. McMartin K (2009) Are calcium oxalate crystals involved in the mechanism of acute renal failure in ethylene glycol poisoning? Clin Toxicol (Phila) 47:859-869
 25. Khan SR, Glenton PA, Byer KJ (2007) Dietary oxalate and calcium oxalate nephrolithiasis. J Urol 178:2191-2196
 26. Khan SR, Glenton PA, Byer KJ (2006) Modeling of hyperoxaluric calcium oxalate nephrolithiasis: experimental induction of hyperoxaluria by hydroxy-L-proline. Kidney Int 70:914-923
 27. Khan SR, Johnson JM, Peck AB, Cornelius JG, Glenton PA (2002) Expression of osteopontin in rat kidneys: induction during ethylene glycol induced calcium oxalate nephrolithiasis. J Urol 168:1173-1181
 28. Jammalamadaka D, Raissi S (2010) Ethylene glycol, methanol and isopropyl alcohol intoxication. American J Med Sci 339:276-281
 29. Green ML, Hatch M, Freel RW (2005) Ethylene glycol induces hyperoxaluria without

metabolic acidosis in rats. Am J Physiol Renal Physiol 289:F536-543

- 1
2 30. Selvam R, Kalaiselvi P, Govindaraj A, Bala Murugan V, Sathish Kumar AS (2001)
3 Effect of *A. lanata* leaf extract and *Vediuppu chunnam* on the urinary risk factors of
4 calcium oxalate urolithiasis during experimental hyperoxaluria. Pharmacol Res
5 43:89-93
6
- 7
8 31. Araujo Viel T, Diogo Domingos C, da Silva Monteiro AP, Riggio Lima-Landman MT,
9 Lapa AJ, Souccar C (1999) Evaluation of the antiurolithiatic activity of the extract of
10 *Costus spiralis* Roscoe in rats. J Ethnopharmacol 66:193-198
11
- 12 32. Gohel MD, Wong SP (2006) Chinese herbal medicines and their efficacy in treating
13 renal stones. Urol Res 34:365-372
14
- 15 33. Cao ZG, Liu JH, Zhou SW, Wu W, Yin CP, Wu JZ (2004) The effects of the active
16 constituents of *Alisma orientalis* on renal stone formation and bikunin expression in
17 rat urolithiasis model. Zhonghua Yi Xue Za Zhi 84:1276-1279
18
- 19 34. Yoshimura K, Miyake O, Okuyama A, Yoshioka T, Honda M, Yamaguchi S, Koide T
20 (1998) Effect of chorei-to and gorei-san on calcium oxalate crystallization in human
21 urine. Hinyokika Kiyo 44:13-16
22
- 23 35. Miyaoka R, Monga M (2009) Use of traditional Chinese medicine in the management
24 of urinary stone disease. Int Braz J Urol 35:396-405
25
- 26 36. Cao ZG, Liu JH, Radman AM, Wu JZ, Ying CP, Zhou SW (2003) An experimental
27 study of effect of different extracts of *Alisma orientalis* on urinary calcium oxalate
28 stones formation in rats. Zhongguo Zhong Yao Za Zhi 28:1072-1075
29
- 30 37. Butterweck V, Khan SR (2009) Herbal medicines in the management of urolithiasis:
31 alternative or complementary? Planta Med 75:1095-1103
32
- 33 38. Atmani F (2003) Medical management of urolithiasis, what opportunity for
34 phytotherapy? Front Biosci 8:s507-514
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Figure 2









Figure 3




Table 1. Biochemical data of rats

	Group 1	Group 2	Group 3	Group 4	Group 5	Group 6
	Control	EG + starch	EG + K-citrate (500 mg/day)	EG + FC (300 mg/day)	EG + FC (600 mg/day)	EG + FC (1200 mg/day)
Baseline						
Body weight (g)	304.7 ± 12.4	313.2 ± 19.0	302.6 ± 12.3	303.1 ± 17.4	305.4 ± 16.5	302.9 ± 19.6
Urine						
Volume (ml/24 h)	21.2 ± 7.9	25.4 ± 4.9	25.5 ± 11.5	22.8 ± 6.4	20.4 ± 6.	19.2 ± 3.0
pH	7.0 ± 0.1	7.1 ± 0.3	7.0 ± 0.1	7.0 ± 0.2	7.1 ± 0.2	7.2 ± 0.3
Micro-albumin (mg/l)	1.0 ± 1.6	3.1 ± 4.2	3.4 ± 3.6	1.9 ± 2.4	3.5 ± 2.4	2.2 ± 2.4
Creatinine (mg/dl)	39.6 ± 13.7	43.0 ± 11.2	51.1 ± 18.8	51.9 ± 17.2	53.6 ± 17.2	58.5 ± 13.6
Uric acid (mg/dl)	10.6 ± 3.3	9.2 ± 2.4	8.2 ± 2.7	6.9 ± 2.3	11.2 ± 2.3	13.5 ± 3.5
Na⁺ (mmol/l)	98.0 ± 17.2	85.7 ± 17.8	111.9 ± 31.6	112.3 ± 34.0	91.9 ± 36.0	98.6 ± 37.9
Mg²⁺ (mmol/l)	20.3 ± 10.4	30.5 ± 15.4	34.0 ± 18.7	29.3 ± 20.4	25.3 ± 20.4	32.1 ± 10.9
Ca²⁺ (mmol/l)	0.51 ± 0.09	0.34 ± 0.04	0.53 ± 0.15	0.54 ± 0.06	0.40 ± 0.15	0.53 ± 0.13
Serum						
pH	7.6 ± 0.2	7.5 ± 0.1	7.4 ± 2.2	7.5 ± 0.1	7.5 ± 0.1	7.5 ± 0.1
Creatinine (mg/dl)	0.63 ± 0.10	0.63 ± 0.11	0.58 ± 0.11	0.63 ± 0.41	0.52 ± 0.09	0.52 ± 0.09
Uric acid (mg/dl)	9.7 ± 0.3	8.9 ± 1.4	8.0 ± 1.6	8.1 ± 1.5	8.1 ± 1.5	7.9 ± 1.7
Na⁺ (mmol/l)	120.4 ± 3.9	121.9 ± 2.5	123.1 ± 2.2	111.6 ± 13.1	119.0 ± 4.9	113.3 ± 9.1
K⁺ (mmol/l)	17.6 ± 1.6	16.9 ± 4.1	16.9 ± 1.8	16.7 ± 2.2	15.2 ± 2.5	15.5 ± 2.4
Cl⁻ (mmol/l)	102.6 ± 2.6	103.5 ± 5.2	100.3 ± 1.2	93.7 ± 12.0	97.5 ± 5.5	94.6 ± 7.2
Mg²⁺ (mmol/l)	3.7 ± 0.15	3.1 ± 1.0	3.2 ± 0.3	3.2 ± 1.0	3.1 ± 0.5	3.2 ± 0.7
Ca²⁺ (mmol/l)	0.82 ± 0.09	0.78 ± 0.11	0.87 ± 0.06	0.69 ± 0.17	0.79 ± 0.72	0.74 ± 0.14
4 weeks						
Body weight (g)	439.6 ± 27.5*	441.0 ± 34.0*	405.8 ± 20.3*	417.2 ± 47.4*	423.0 ± 40.8*	409.5 ± 46.3*
Left kidney weight (g)	1.9 ± 0.3	1.9 ± 0.4	1.8 ± 0.2	1.9 ± 0.3	1.7 ± 0.2	1.7 ± 0.1
Right kidney weight (g)	2.0 ± 0.2	1.9 ± 0.2	1.8 ± 0.4	1.8 ± 0.3	1.8 ± 0.3	1.7 ± 0.2
Urine						
Volume (ml/24 h)	26.0 ± 6.0	43.2 ± 15.9 [#]	30.5 ± 13.6	37.0 ± 11.8	27.8 ± 8.3	24.8 ± 8.3 [†]
pH	7.2 ± 0.3	7.0 ± 0.2	7.1 ± 0.4	7.0 ± 0.5	7.0 ± 0.5	7.1 ± 0.4
Micro-albumin (mg/l)	0.5 ± 0.6	2.1 ± 4.2	1.2 ± 3.2	1.1 ± 1.9	1.3 ± 3.8	1.4 ± 2.6
Creatinine (mg/dl)	55.3 ± 16.6	41.9 ± 18.6	50.3 ± 20.2	47.2 ± 19.9	64.0 ± 23.3	58.9 ± 26.8

Uric acid (mg/dl)	7.2 ± 0.6	6.8 ± 3.1	7.3 ± 3.1	6.4 ± 2.8	8.5 ± 2.8	7.7 ± 3.4
Na⁺ (mmol/l)	108.9 ± 18.7	78.6 ± 27.7	72.9 ± 16.0	90.9 ± 27.4	109.7 ± 28.5	93.1 ± 41.3
Mg²⁺ (mmol/l)	21.6 ± 15.7	16.1 ± 5.4	14.1 ± 11.8	25.3 ± 7.2	17.6 ± 7.3	12.1 ± 9.6
Ca²⁺ (mmol/l)	0.49 ± 0.24	0.36 ± 0.15	0.43 ± 0.12	0.39 ± 0.26	0.63 ± 0.15	0.53 ± 0.25
Serum						
pH	7.3 ± 0.0	7.4 ± 0.1	7.3 ± 0.1	7.3 ± 0.1	7.3 ± 0.1	7.3 ± 0.0
Creatinine (mg/dl)	0.45 ± 0.07	0.51 ± 0.11	0.23 ± 0.06 [†]	0.56 ± 0.06	0.51 ± 0.04	0.52 ± 0.04
Uric acid (mg/dl)	2.1 ± 0.5	4.0 ± 1.2 [#]	5.5 ± 0.6	4.5 ± 1.4	3.5 ± 0.7	3.0 ± 0.8
Na⁺ (mmol/l)	137.8 ± 1.7	135.6 ± 5.3	135.2 ± 2.1	125.4 ± 38.2	137.4 ± 2.1	137.1 ± 1.2
K⁺ (mmol/l)	5.1 ± 0.7	8.1 ± 3.8	7.4 ± 1.3	6.6 ± 0.4	6.2 ± 1.3	6.5 ± 1.7
Cl⁻ (mmol/l)	100.0 ± 1.0	100.8 ± 1.0	100.1 ± 1.3	99.8 ± 1.6	100.1 ± 0.5	99.6 ± 0.8
Mg²⁺ (mmol/l)	2.2 ± 0.3	2.5 ± 0.5	2.7 ± 0.5	2.7 ± 0.6	2.0 ± 0.4	2.4 ± 0.4
Ca²⁺ (mmol/l)	1.3 ± 0.1	1.2 ± 0.2	1.2 ± 0.1	1.3 ± 0.1	1.3 ± 0.1	1.3 ± 0.1
Albumin (mg/dl)	4.0 ± 0.1	4.0 ± 0.9	4.0 ± 0.3	4.1 ± 0.2	4.0 ± 0.1	4.0 ± 0.2
AST (U/l)	80.8 ± 4.3	153.6 ± 39.7 [#]	185.2 ± 43.1	107.7 ± 25.1 [†]	95.1 ± 10.1 [†]	101.7 ± 20.1 [†]
ALT (U/l)	51.4 ± 5.8	74.6 ± 8.6 [#]	69.8 ± 7.9	64.5 ± 5.1	58.6 ± 7.6 [†]	52.8 ± 13.2 [†]
PT (sec)	10.2 ± 0.3	10.3 ± 0.1	27.9 ± 25.7	10.2 ± 0.3	17.7 ± 19.1	17.4 ± 19.2
APTT (sec)	19.4 ± 2.2	18.6 ± 0.9	75.5 ± 81.0	20.8 ± 3.1	45.5 ± 60.1	43.4 ± 60.3

Values are expressed as Mean ± SD

* p < 0.05, values are significantly different compared to the relative groups in baseline

p < 0.05, values are significantly different compared to control group

† p < 0.05, values are significantly different compared to “EG + starch” group.

Respond to Reviewers

[Click here to download Supplementary Material: Response0213_UR.pdf](#)