

The effects of *Fatsia polycarpa* Hayata extracts on anti-*Helicobacter pylori* activity

Hong-Han Shih¹ (施閔瀚), Chiu-Hsiang Su¹ (蘇千香), Yuan-Man Hsu¹ (徐媛曼),
Chang-Hung Chou² (周昌弘)

Department of Biological Science and Technology, College of Life Sciences, China Medical University, Taichung 404, Taiwan¹

Research Center for Biodiversity, China Medical University, Taichung 404, Taiwan²

Fatsia polycarpa Hayata has been used as an herbal medicine to treat ankylosing spondyloarthritis, oseteoarthritis, rheumatism, rheumatoid arthritis with accompanying reactive gout, osteochondrosis, synovitis, and tendinitis in Chinese medicine for many years. We analyzed the natural products isolated from *Fatsia polycarpa* Hayata by evaluating the anti-*H. pylori* activity and the effects of these compounds on the suppression of human gastric epithelial AGS cells. Fraction DM-24-6-3-1 exhibited the strongest antibacterial activity against *H. pylori*, with minimum bactericidal concentration of 4 µg/ml. Fraction DM-24-7-7 exhibited the strongest cytotoxicity against AGS cells, with IC₅₀ of 14.3 µg/ml. However, fraction DM-24-6-3-1 showed a weak cytotoxicity against AGS cells (158 µg/ml). Therefore, the active compound in this fraction will be further identified and might be valuable for treating *H. pylori* infection.