



Specific Expression of MAPK p38 β Dominates BAD^{ser112} Phosphorylation Contributes to TNF- α Resistance in Oral Cancer

Wei-Jen Ting¹, Wei-Wen Kuo², Da-Tian Bau³, Chih-Yang Huang^{1,4,*}
¹Graduate Institute of Basic Medical Science, China Medical University, Taiwan
²Department of Biological Science and Technology, China Medical University, Taiwan
³Chairman of Terry Fox Cancer Research Lab, Department of Medical Research, China Medical University Hospital, Taiwan
⁴Department of Health and Nutrition Biotechnology, Asia University, Taiwan

Backgrounds: When oral cancer is diagnosed, the three-year survival rate prediction of this patient is only 58% and can only be increased to 74% after surgery in Taiwan. Before the squamous cell carcinoma (SCC) formed, hyperlosia is an initial stage symptom induced by EGFR over expression in a long term inflammation. Thus, tumor necrosis factor- α (TNF- α) releasing in inflammation lose its original anti-tumor function and a TNF- α resistance might happen in many cases. In our previous study indicated p38 β MAPK over expression in oral cancer might be associated with TNF- α resistance through serine 112 of BAD phosphorylation, and which is a gatekeeper of BAD-mediated apoptosis.

Materials and Methods: In this research, a cell line T28 from 4-nitroquinoline-N-oxide (4-NQO) induced oral cancer in C57B mouse and human tongue squamous cell carcinoma cell line SCC4 were screened in this TNF- α resistance issue. All proteins from cell were analyzed by immune blot assay.

Results: TNF- α releasing is through p38 mitogen-activated protein kinases and TNF- α resistance exists in both T28 and SCC4 cell lines. Further, the serine 136 of BAD phosphorylation was promoted by p38 α MAPK isoform and the serine 112 and 155 of BAD phosphorylation were promoted by p38 β MAPK and also block the apoptosis caused by TNF- α . A p38 β MAPK inhibitor SB202190 (10 μ M) was used, the cell cycle arrested at G2 phase from 9.5% to 17.36% within 24h treatment in SCC4 cells.

Conclusion: Over expression of p38 β MAPK in oral cancer indeed caused TNF- α induced apoptosis resistance by BAD phosphorylation. And serine 112 of BAD is controlled by p38 β MAPK. Above these experimental evidences suggest that p38 β MAPK is a possible anticancer target in oral cancer therapy.




Figure 1. The relationships between TNF- α expressions and cell proliferation rate in normal / oral cancer cells. (A) p38 MAPK over expression induced TNF- α releasing and BAD phosphorylation in tumor cell lines rather than normal cells lines. (B) The cell proliferation rates in tumor cell lines are higher than normal cell lines. (Cell lines: H9C2 is a rat cardiomyoblast cell line as normal. N28 is none tumor cell line from C57B mouse. T8 and T28 are 4NQO induced oral cancer cell from C57B mouse, SCC4 and CE146T are human oral cancer cell lines.)




Figure 2. The role of phosphorylated BAD in TNF- α resistance. (A) After 24h 5ng/ml TNF- α treatment, Caspase 3 expression increased and phosphorylated BAD decreased in H9C2 cell. (B) The same 24 h 5ng/ml TNF- α treatment in SCC4 cell can not induce the BAD dependent apoptosis, and show the TNF- α resistance existed in this oral cancer cell line. (C) The cell survival percentage between H9C2 and SCC4 cell line in 24h TNF- α treatment. Data are expressed as mean \pm SE (n=3) and * = p < 0.01, ** = p < 0.05, *** = p < 0.001 as compared with the control group.




Figure 3. The p38 α / β MAPK related BAD phosphorylation. The p38 α / β MAPK specific inhibitor CMPD1 (10 μ M) and SB202190 (10 μ M) were used to block the p38 α / β expressions in human oral cancer cell line SCC4 for 24h, and the down stream BAD phosphorylation location (serine 112, 136, 155) decreased following the p38 α / β inhibitions. The SB202190 treatments remarkably lowered the p-BAD^{ser112} and p-BAD^{ser155} induced the caspase 3 cleavage in SCC4 cells.

Figure 4. The complex formed by p38 α / β MAPK and different BAD phosphorylation locations. After 30 min anisomycin treatments, the co-immunoprecipitation assay was used in complex detection between p38 α / β MAPK and different BAD phosphorylation location points (serine 112, 136, 155). The result shown that, p38 α combined with p-BAD^{ser155} only and p38 β combined with p-BAD^{ser112} and p-BAD^{ser155} in the formed complex.




Figure 5. The SB202190 inhibit the p-BAD^{ser112} expressions in TNF- α resistance oral cancer cells. After 24h TNF- α (human, 5ng) or SB202190 (10 μ M) co-treatments, the Laser Scanning Confocal Microscope imagination shown a partial p-BAD^{ser112} exist in cytoplasm (in degradation progress) and a partial p-BAD^{ser112} exist in nucleus. After TNF- α treatment for 24h, a large amount of p-BAD^{ser112} exist in nucleus. SB202190 (10 μ M) and TNF- α co-treatments for 24h, p-BAD^{ser112} exist in cytoplasm only.




Figure 6. The SB202190 treatments caused a cell cycle arrest in SCC4 cells. The cell cycle of human oral cancer cell SCC4 arrest while a p38 β specific inhibitor SB202190 (10 μ M) treatments from 0 to 24h. The cell cycle of SCC4 cell is major arrest at G2/M, from 9.50% (control) to 17.36% (SB202190, 10 μ M) within 24h.




Figure 7. p38 β MAPK and p-Bad^{ser112} was co-localized in the cytoplasm of the T28 and SCC4 oral cancer cells. (A) The confocal microscopy assay showed the evidences about the co-localization (indicated by arrows) of p38 β MAPK and p-BAD^{ser112} in T28 and (B) SCC4 cells. These co-localization expressions were enhanced within 30 min of 10 μ M anisomycin treatments, and inhibited by the SB202190 treatment in both T28 and SCC4 oral cancer cells.




Figure 8. The anti-tumor effects of p38 β MAPK inhibitor treatments in C27B mice. The T28 cell was xenografted into C57B mice for the oral cancer animal model. After 30 days induction, different concentrations of SB202190 (0, 10, 20 μ M) were used and directly injected into the tumor of the C57B mice. (A) H&E staining assay indicates the structures of oral cancer tissues were changed within 7 days injection. (B) DAPI and TUNEL staining assays indicate the nuclei location and the apoptosis cell DNA fragments. (C) The tumor volume was still increased without any treatment, but it reduced by SB202190 efficiently and shown a dose dependent manner. (D) The proteins analysis shown that the SB202190 injection caused the p38 β and p-BAD^{ser112} expressions reductions and increased the caspase-3 cleavage form in the oral cancer.




Figure 9. The p38 MAPK dominates TNF- α release and p38 β MAPK expression cause a TNF- α resistance through the downstream BAD phosphorylation. In several published research shown that the TNF- α releasing is dominated by p38 MAPK (p38 α major) activation by the outcome stress. At the same time, the stress could induce the p38 α MAPK provides a BAD phosphorylation at serine 155 and p38 β MAPK expression provides a BAD phosphorylation at serine 112 and 155. The serine 112 of BAD plays a gatekeeper role preventing BAD dependent apoptosis caused by the protein phosphatases (such as PP2A) dephosphorylation. Finally, the BAD phosphorylation provides a protection of human oral cancer cell from the stress caused TNF- α induced apoptosis and promotes oral cancer a TNF- α resistance ability.