

**Chronic Osteomyelitis is Associated with Increased Risk of New-Onset Atrial
Fibrillation: Evidence from a Nationwide Cohort of 23 Million People**

Lien-Cheng Hsiao^{1,2}, Chih-Hsin Muo³, Che-Yi Chou^{2,4}, Chun-Hung Tseng^{5,6},
Ming-Fong Chen¹, and Kuan-Cheng Chang^{1,2}

Affiliations: ¹Division of Cardiology, Department of Internal Medicine, China Medical University Hospital, Taichung, 40447, TAIWAN; ²Graduate Institute of Clinical Medical Science, China Medical University, Taichung 40402, TAIWAN; ³Institute of Environmental Health, College of Public Health, China Medical University, Taichung, 40402, TAIWAN; ⁴Department of Nephrology, China Medical University Hospital, Taichung, 40447, TAIWAN; ⁵Department of Neurology, China Medical University Hospital, Taichung, 40447, TAIWAN; ⁶School of Medicine, China Medical University, Taichung, 40402, TAIWAN

Running head: Chronic Osteomyelitis and Atrial Fibrillation

Potential Competing Interests: The authors have declared that no competing interests exist.

Word count: 2416

Address correspondence to:

Kuan-Cheng Chang, M.D., Ph.D, FESC.

Division of Cardiology, Department of Medicine, China Medical University Hospital

2, Yuh-Der Road, Taichung 40447, Taiwan

Tel: 886-4-22052121, ext. 2311 Fax: 886-4-23602467

E-mail: kuancheng.chang@gmail.com

Or

Chun-Hung Tseng, MD.

Department of Neurology, China Medical University Hospital, and School of

Medicine, China Medical University

No.91 Hsueh-Shih Road, Taichung, 40402, Taiwan

Tel: 886-4-2205-3366; Fax: 886-4-2234-4055

E-mail: d8333@mail.cmuh.org.tw